

STATISTICS WORKSHEET-1

Q1 to Q9 have only one correct answer. Choose the correct option to answer your question.

1. Bernoulli random variables take (only) the values 1 and 0.
 - a) **True**
 - b) False
2. Which of the following theorem states that the distribution of averages of iid variables, properly normalized, becomes that of a standard normal as the sample size increases?
 - a) **Central Limit Theorem**
 - b) Central Mean Theorem
 - c) Centroid Limit Theorem
 - d) All of the mentioned
3. Which of the following is incorrect with respect to use of Poisson distribution?
 - a) Modeling event/time data
 - b) **Modeling bounded count data**
 - c) Modeling contingency tables
 - d) All of the mentioned
4. Point out the correct statement.
 - a) The exponent of a normally distributed random variables follows what is called the log- normal distribution
 - b) Sums of normally distributed random variables are again normally distributed even if the variables are dependent
 - c) The square of a standard normal random variable follows what is called chi-squared distribution
 - d) **All of the mentioned**
5. _____ random variables are used to model rates.
 - a) Empirical
 - b) Binomial
 - c) **Poisson**
 - d) All of the mentioned
6. 10. Usually replacing the standard error by its estimated value does change the CLT.
 - a) True
 - b) **False**
7. 1. Which of the following testing is concerned with making decisions using data?
 - a) Probability
 - b) **Hypothesis**
 - c) Causal
 - d) None of the mentioned
8. 4. Normalized data are centered at _____ and have units equal to standard deviations of the original data.
 - a) **0**
 - b) 5
 - c) 1
 - d) 10
9. Which of the following statement is incorrect with respect to outliers?
 - a) Outliers can have varying degrees of influence
 - b) Outliers can be the result of spurious or real processes
 - c) **Outliers cannot conform to the regression relationship**
 - d) None of the mentioned

Q10 and Q15 are subjective answer type questions, Answer them in your own words briefly.

10. What do you understand by the term Normal Distribution?

Ans) Normal distribution, also known as the Gaussian distribution, is a probability distribution that is symmetric about the mean, showing that data near the mean are more frequent in occurrence than data far from the mean. In graph form, normal distribution will appear as a bell curve

- In a normal distribution the mean is zero and the standard deviation is 1. It has zero skew and a kurtosis of 3.
- Normal distributions are symmetrical, but not all symmetrical distributions are normal.

11. How do you handle missing data? What imputation techniques do you recommend?

Ans) Handling the missing data: When dealing with missing data, data scientists can use two primary methods to solve the error: **imputation or the removal of data**. The imputation method develops reasonable guesses for missing data. It's most useful when the percentage of missing data is low

Implementation techniques:

*Complete case analysis(CCA)

*Arbitrary value Implementation

*Frequent category implementation

12. What is A/B testing?

Ans) A/B testing is a basic randomized control experiment. It is a way to compare the two versions of a variable to find out which performs better in a controlled environment.

13. Is mean imputation of missing data acceptable practice?

Ans) Mean imputation is typically considered terrible practice since it ignores feature correlation. Consider the following scenario: we have a table with age and fitness scores, and an eight-year-old has a missing fitness score. If we average the fitness scores of people between the ages of 15 and 80, the eighty-year-old will appear to have a significantly greater fitness level than he actually does.


14. What is linear regression in statistics?

Ans) Linear regression quantifies the relationship between one or more predictor variable(s) and one outcome variable. Linear regression is commonly used for predictive analysis and modelling.

15. What are the various branches of statistics?

Ans) The two main branches of statistics are descriptive statistics and inferential statistics. Both of these are employed in scientific analysis of data and both are equally important for the student of statistics.

Answers in red color



FLIP ROBO