

Problem Sheet 4

discussion: week of Monday, 21.11.2022

- 4.1.** **a)** Show that the weights of the Newton-Cotes formulas satisfy $\sum_{i=0}^n w_i = 1$ (= length of the interval $[0, 1]$).
(Hint: apply the quadrature formula to a suitable function f .)
- b)** Show the symmetry property $w_{n-i} = w_i$, $i = 0, \dots, n$.
(Hint: use the symmetry of the points, i.e., $x_j = 1 - x_{n-j}$.)
- c)** Let $n = 2m$ be even. Consider the function $f = (x - 1/2)^{n+1}$, which is antisymmetric with respect to $1/2$. Show: $\int_0^1 f(x) dx = 0 = \hat{Q}_n^{cNC}(f) = \hat{Q}_n^{oNC}(f)$. Conclude that the quadrature formulas \hat{Q}_n^{cNC} and \hat{Q}_n^{oNC} are exact for polynomials of degree $n + 1$.
 In particular, the midpoint rule is exact for polynomials in \mathcal{P}_1 , and the Simpson rule is exact for polynomials in \mathcal{P}_3 .

- 4.2.** Let $C, \alpha > 0$ and consider the function $h \mapsto f(h) = Ch^\alpha$. Why is this function a straight line in a **loglog**-plot? What is its slope? Other popular plotting schemes are, **semilogx** and **semilogy**. Which one would you use to plot functions of the form $N \mapsto Ce^{-bN}$? How would you proceed if you suspect that a function $h \mapsto f(h)$ has the form $f(h) = Ce^{-b/h}$? What if you suspect $f(h) = Ce^{-b/h^2}$?

- 4.3.** We wish to show that the extrapolation of the composite trapezoidal rule is the composite Simpson rule. To that end, let $T(h)$ be the composite trapezoidal rule with step size $h = (b - a)/N$ and $S(h)$ be the composite Simpson rule with step size $h = (b - a)/N$. Use Romberg extrapolation with step sizes $h_i = (b - a)2^{-i}$, $i = 0, 1, \dots$,

- a)** Extrapolation of the composite trapezoidal rule (with step size h) has in column $m = 0$ the values $T(h_i)$. Show: in column $m = 1$ of the Neville scheme are the values

$$N_i := T(h_{i+1}) + \frac{1}{3}(T(h_{i+1}) - T(h_i))$$

- b)** Show: $N_i = S(h_i)$.

- 4.4.** Write a program with signature $I = \text{adapt}(f, a, b, \tau, h_{\min})$ that realizes an adaptive quadrature for $\int_a^b f(x) dx$. The quadrature should be based on the Simpson rule. τ is the desired (absolute) accuracy and h_{\min} the minimal interval length. To estimate the accuracy, compare the value of the Simpson rule $S_{\{a,b\}}(f)$ for the integration on $[a, b]$ with the value $S_{\{a,m\}}(f) + S_{\{m,b\}}(f)$ with $m = (a + b)/2$. Use your algorithm for the integration of the function

$$\begin{cases} \frac{1}{2}e^x & x < 1/3 \\ e^x & x \geq 1/3 \end{cases}$$

over $[0, 1]$. Use $\tau = h_{\min} = 2^{-j}$, $j = 0, \dots, 10$. Plot the error versus τ . What convergence do you observe? Why was $h_{\min} = \tau$ chosen?

- 4.5.** Develop an adaptive algorithm for the integration of functions over the rectangle $[a_x, b_x] \times [a_y, b_y]$. Base your algorithm on the midpoint rule, i.e., $Q_{[a,b] \times [c,d]}(f) = (b-a)(d-c)f((a+b)/2, (c+d)/2)$. *Hint: adapt the ideas of the 1d-adaptive algorithm of Problem 4.4. Test your adaptive algorithm for the integration over $[0, 1]^2$ of the following functions:*

$$f_1(x, y) = x^2 \quad \text{and} \quad f_2(x, y) = \begin{cases} 0 & x < y \\ 1 & x \geq y \end{cases}$$

Use the tolerances $\tau = 2^{-i}$, $i = 0, \dots, 15$, and make a convergence plot (quadrature error versus tolerance) in **loglog** scale.