

1. Some special nonlinear differential equations can be transformed to linear equations. Such examples are the *Bernoulli equations*, which have the form

$$u'(x) + P(x)u(x) = Q(x)u^n(x), \quad (1)$$

where  $n \neq 0, 1$  and  $P, Q$  are given functions.

- a) Show that the transformation  $v = u^{1-n}$  transforms the Bernoulli equation into the linear ODE

$$v'(x) + (1-n)P(x)v(x) = (1-n)Q(x).$$

- b) Use this to solve the Bernoulli equation

$$u' + 2xu = u^2.$$

a)  $v = u^{1-n}$

$$v' = (1-n) u^{-n} u' \rightarrow u' = \frac{v'}{(1-n)} u^n$$

in (1)

$$\frac{v'}{(1-n)} u^n + P u = Q u^n \quad \left| u^{-n} \right| \cdot (1-n)$$

$$v' + (1-n)P u^{1-n} = (1-n)Q \quad [v = u^{1-n}]$$

$$v' + (1-n)P v = (1-n)Q \quad (2)$$

b)  $v' + 2xv = v^2 \rightarrow P(x) = 2x, Q(x) = 1, n = 2, v = u^{-1}$

(2):  $v' + (-1) 2x v = (-1)$

$$v' - 2xv = 0$$

how:  $v' = 2xv \quad f(x) = 2x \quad g(v) = v$

$$\int f(x) dx = \int \frac{1}{v} dv \Rightarrow x^2 + C = \ln v$$

$$v_H = C e^{x^2}$$

We finish this section with a general formula for the computation of particular solutions for systems

$$u'(x) = a(x)u(x) + g(x),$$

The general idea behind the variation of constant formula is that a particular solution should have a similar structure, i.e., look like

$$u_p(x) = U(x) \cdot c(x),$$

$$v' = \underbrace{2x}_{a(x)} \underbrace{v}_{v(x)} - \underbrace{1}_{g(x)}$$

$$V_p = e^{x^2} c(x) \rightarrow U = e^{x^2}, U^{-1} = e^{-x^2}$$

$$v' = 2x e^{x^2} c(x) + c'(x) e^{x^2}$$

-> insert:

$$\cancel{2x e^{x^2} c(x)} + c'(x) e^{x^2} = \cancel{2x e^{x^2} c(x)} - 1$$

$$c'(x) = -e^{-x^2} \quad | \int$$

$$\rightarrow c(x) = -\int e^{-x^2} dx$$

For first order linear equations (i.e.  $u, a, g$  are scalar functions), this formula reduces to

$$u_p(x) = e^{A(x)} \int e^{-A(y)} g(y) dy,$$

where  $A(x) = \int a(x) dx$ .

$$A(x) = \int a(x) dx = \int 2x dx = x^2$$

$$V_p = e^{x^2} \int e^{-x^2} (-1) dx$$

$$\rightarrow V = V_H + V_p$$

$$v = c e^{x^2} - e^{x^2} \int e^{-x^2} dx$$

$$v = e^{x^2} \left( c - \int e^{-x^2} dx \right)$$

$$[v = v^{-1}] \quad u = v^{-1} = \left[ e^{x^2} \left( c - \int e^{-x^2} dx \right) \right]^{-1}$$

2. Use the variation of constants formula to solve the ODE

$$u'' - \frac{3}{x}u' - \frac{5}{x^2}u = \log(x).$$

In order to derive the general solution for the homogeneous equation, make the ansatz  $u(x) = x^\alpha$  and compute  $\alpha \in \mathbb{N}$ .

hom:  $u(x) = x^\alpha$

$$u'(x) = \alpha x^{\alpha-1}$$

$$u''(x) = \alpha(\alpha-1)x^{\alpha-2}$$

$$\rightarrow \alpha(\alpha-1)x^{\alpha-2} - \frac{3}{x}\alpha x^{\alpha-1} - \frac{5}{x^2}x^\alpha = 0$$

$$(\alpha^2 - \alpha)x^{\alpha-2} - 3\alpha x^{\alpha-2} - 5x^{\alpha-2} = 0$$

$$(\alpha^2 - \alpha - 3\alpha - 5)x^{\alpha-2} = 0$$

$$\alpha^2 - 4\alpha - 5 = 0$$

$$2 \pm \sqrt{4+5} = 2 \pm 3$$

$$\alpha_1 = 5, \quad \alpha_2 = -1 \notin \mathbb{N}$$

$$u_H = c_1 x^5 + c_2 x^{-1}$$

$$V_1 = u, \quad V_2 = u'$$

$$\rightarrow V_1' = V_2$$

$$V_2' = \frac{3}{x}V_2 + \frac{5}{x^2}V_1 + \log(x)$$

**Example.** We want to solve the inhomogeneous ODE

$$u''(x) + u'(x) = e^{-x}$$

with the variation of constants formula. Using the characteristic equation  $\lambda^2 + \lambda = 0$ , we obtain the general solution  $c_1 + c_2 e^{-x}$  of the homogeneous equation. Now, we rewrite the ODE into a first order system by introducing  $v_1 := u$  and  $v_2 := u'$ . Then,

$$v_1' = v_2$$

$$v_2' = -v_2 + e^{-x}$$

or in matrix notation

$$\begin{pmatrix} v_1 \\ v_2 \end{pmatrix}' = \begin{pmatrix} 0 & 1 \\ 0 & -1 \end{pmatrix} \cdot \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} + \begin{pmatrix} 0 \\ e^{-x} \end{pmatrix}.$$

$$\begin{aligned} \underbrace{\begin{pmatrix} v_1 \\ v_2 \end{pmatrix}'} &= \underbrace{\begin{pmatrix} 0 & 1 \\ \frac{3}{x^2} & \frac{1}{x} \end{pmatrix}}_{\text{matrix}} \underbrace{\begin{pmatrix} v_1 \\ v_2 \end{pmatrix}}_{\text{vec}(x)} + \underbrace{\begin{pmatrix} 0 \\ \log(x) \end{pmatrix}}_{g(x)} \\ u &= \begin{pmatrix} u_H \\ u_H' \end{pmatrix} = \begin{pmatrix} x^5 & x^{-1} \\ 5x^4 & -x^{-2} \end{pmatrix} \end{aligned}$$

$$U^{-1}: \left( \begin{array}{cc|cc} x^5 & x^{-1} & 1 & 0 \\ 5x^4 & -x^{-2} & 0 & 1 \end{array} \right) \rightarrow \left( \begin{array}{cc|cc} x^4 & x^{-2} & x^{-1} & 0 \\ 5x^4 & -x^{-2} & 0 & 1 \end{array} \right)$$

$$\rightarrow \left( \begin{array}{cc|cc} x^4 & x^{-2} & x^{-1} & 0 \\ 0 & -6x^{-2} & -5x^{-1} & 1 \end{array} \right) \rightarrow \left( \begin{array}{cc|cc} x^4 & x^{-2} & x^{-1} & 0 \\ 0 & x^{-2} & \frac{5}{6}x^{-1} & -\frac{1}{6} \end{array} \right)$$

$$\rightarrow \left( \begin{array}{cc|cc} x^4 & 0 & x^{-1} - \frac{5}{6}x^{-1} & \frac{1}{6} \\ 0 & x^{-2} & \frac{5}{6}x^{-1} & -\frac{1}{6} \end{array} \right) \rightarrow \left( \begin{array}{cc|cc} 1 & 0 & x^{-5}(1 - \frac{5}{6}) & \frac{x^{-6}}{6} \\ 0 & 1 & \frac{5}{6}x & -\frac{x^2}{6} \end{array} \right)$$

$$\rightarrow U^{-1} = \frac{1}{6} \begin{pmatrix} x^{-5} & x^{-4} \\ 5x & -x^2 \end{pmatrix}$$

$$v_p(x) = U(x) \int U^{-1}(y)g(y)dy$$

$$v_p = \begin{pmatrix} x^5 & x^{-1} \\ 5x^4 & -x^{-2} \end{pmatrix} \int \frac{1}{6} \begin{pmatrix} y^{-5} & y^{-4} \\ 5y & -y^2 \end{pmatrix} \begin{pmatrix} 0 \\ \log(y) \end{pmatrix} dy$$

$$= \begin{pmatrix} x^5 & x^{-1} \\ 5x^4 & -x^{-2} \end{pmatrix} \frac{1}{6} \int \begin{pmatrix} y^{-4} \log(y) \\ -y^2 \log(y) \end{pmatrix} dy \quad [ \int f'g = fg - \int fg' ]$$

$$\int y^{-4} \log y \, dy = -\frac{y^{-3}}{3} \log y - \int -\frac{y^{-3}}{3} \frac{1}{y} dy = -\frac{y^{-3}}{3} \log y + \int \frac{y^{-4}}{3} dy$$

$$= -\frac{y^{-3}}{3} \log y - \frac{y^{-3}}{9}$$

$$\int -y^2 \log y \, dy = -\frac{y^3}{3} \log y - \int -\frac{y^3}{3} y^{-1} dy = -\frac{1}{3} y^3 \log y + \frac{1}{6} \frac{y^3}{3}$$

$$= \begin{pmatrix} x^5 & x^{-1} \\ 5x^4 & -x^{-2} \end{pmatrix} \frac{1}{6} \begin{pmatrix} -\frac{1}{3}x^{-3}(\log x + \frac{1}{3}) \\ -\frac{x^3}{3}(\log x - \frac{1}{3}) \end{pmatrix}$$

$$= -\frac{1}{18} \begin{pmatrix} x^5 x^{-3}(\log x + \frac{1}{3}) + x^{-1} x^3(\log x - \frac{1}{3}) \\ 5x^4 x^{-3}(\log x + \frac{1}{3}) - x^{-2} x^3(\log x - \frac{1}{3}) \end{pmatrix}$$

$$= -\frac{1}{18} \begin{pmatrix} x^2(\log x + \frac{1}{3}) + x^2(\log x - \frac{1}{3}) \\ 5x(\log x + \frac{1}{3}) - x(\log x - \frac{1}{3}) \end{pmatrix}$$

$$= -\frac{1}{18} \begin{pmatrix} 2x^2 \log x \\ 4x \log x + \frac{6}{3}x \end{pmatrix} = \begin{pmatrix} -\frac{1}{9}x^2 \log x \\ -\frac{2}{9}x \log x - \frac{1}{9}x \end{pmatrix}$$

$$\begin{pmatrix} v_1 \\ v_2 \end{pmatrix} = U \cdot \begin{pmatrix} c_1 \\ c_2 \end{pmatrix} + v_p$$

with  $v_n = u$

$$u = c_1 x^5 + c_2 x^{-1} - \frac{1}{9}x^2 \log x$$

3. (Picard iteration) The Picard-Lindelöf/Cauchy-Lipschitz theorem in fact gives more than what we discussed in the lecture: it also provides a way to approximate solutions of the initial value problem

$$\begin{aligned} u' &= f(t, u) \\ u(0) &= u_0. \end{aligned}$$

This is the so-called *Picard iteration*, defined as

$$\begin{aligned} \varphi_0(t) &= u_0 \\ \varphi_n(t) &= u_0 + \int_0^t f(s, \varphi_{n-1}(s)) ds \quad n \geq 1. \end{aligned}$$

The produced sequence  $(\varphi_n)_{n \in \mathbb{N}}$  converges (provided the conditions in the Picard-Lindelöf theorem hold) to the solution  $u$  of the ODE.

- a) Compute the first 3 elements in the Picard iteration for

$$u'(t) = tu(t), \quad u(0) = 1.$$

To which function does the sequence converge?

- b) Compute the first 4 elements in the Picard iteration for

$$u'(t) = u^2(t), \quad u(0) = 1.$$

To which function does the sequence seem to converge?

a)  $u' = tu$ ,  $u := u(t)$ ,  $f(t, u) = tu$

$$u(0) = u_0 = 1 \rightarrow \varphi_0(t) = 1$$

$$\varphi_1(t) = 1 + \int_0^t (s \cdot 1) ds = 1 + \frac{s^2}{2} \Big|_0^t = 1 + \frac{t^2}{2} = \varphi_1(t)$$

$$\varphi_2(t) = 1 + \int_0^t \left( s \cdot \left( 1 + \frac{s^2}{2} \right) \right) ds$$

$$= 1 + \int_0^t s ds + \int_0^t \frac{s^3}{2} ds$$

$$= 1 + \frac{t^2}{2} + \frac{t^4}{8} = \varphi_2(t)$$

$$\left[ e^x = \sum_{i=0}^{\infty} \frac{x^i}{i!} \right]$$

$$\left[ e^{\frac{t^2}{2}} = 1 + \frac{t^2}{2} + \frac{\frac{t^4}{4}}{2} + \dots \right]$$

converges to  $e^{\frac{t^2}{2}}$

$$b) \quad u' = u^2(L) \quad u(0) = u_0 = 1$$

$$\varphi_0 = 1$$

$$f(t, u) = u^2(L)$$

3. (Picard iteration) The Picard-Lindelöf/Cauchy-Lipschitz theorem in fact gives more than what we discussed in the lecture: it also provides a way to approximate solutions of the initial value problem

$$\begin{aligned} u' &= f(t, u) \\ u(0) &= u_0. \end{aligned}$$

This is the so-called *Picard iteration*, defined as

$$\begin{aligned} \varphi_0(t) &= u_0 \\ \varphi_n(t) &= u_0 + \int_0^t f(s, \varphi_{n-1}(s)) \, ds \quad n \geq 1. \end{aligned}$$

The produced sequence  $(\varphi_n)_{n \in \mathbb{N}}$  converges (provided the conditions in the Picard-Lindelöf theorem hold) to the solution  $u$  of the ODE.

$$\varphi_1 = 1 + \int_0^t 1 \, ds = 1 + t = \varphi_1(L)$$

$$\varphi_2 = 1 + \int_0^t (1+s)^2 \, ds = 1 + \int_0^t 1 + 2s + s^2 \, ds$$

$$\varphi_2(t) = 1 + t + t^2 + \frac{t^3}{3}$$

$$\varphi_3 = 1 + \int_0^t \left(1 + s + s^2 + \frac{s^3}{3}\right)^2 \, ds$$

$$\left(1 + s + s^2 + \frac{s^3}{3}\right) \left(1 + s + s^2 + \frac{s^3}{3}\right) =$$

$$\underbrace{1} + \underbrace{s} + \underbrace{s^2} + \underbrace{\frac{s^3}{3}} + \underbrace{s} + \underbrace{s^2} + \underbrace{\frac{s^3}{3}} + \underbrace{\frac{s^4}{3}} + \underbrace{s^2} + \underbrace{s^3} + \underbrace{\frac{s^4}{3}} + \underbrace{\frac{s^5}{3}}$$

$$+ \underbrace{\frac{s^3}{3}} + \underbrace{\frac{s^4}{2}} + \underbrace{\frac{s^5}{2}} + \underbrace{\frac{s^6}{9}} = 1 + 2s + 3s^2 + \frac{8}{3}s^3 + \frac{5}{2}s^4 + \frac{2}{3}s^5 + \frac{s^6}{9}$$

$$\varphi_3 = 1 + \int_0^s 1 + 2t + 3t^2 + \frac{8}{3}t^3 + \frac{5}{2}t^4 + \frac{2}{3}t^5 + \frac{t^6}{9} \, dt \leftarrow s \text{ and } L \text{ should be switched here but too much writing}$$

$$\varphi_3 = 1 + t + t^2 + t^3 + \frac{8}{12}t^4 + \frac{5}{15}t^5 + \frac{2}{18}t^6 + \frac{t^7}{63}$$

Seems to converge to

$$\sum_{i=0}^{\infty} t^i = \frac{1}{1-t}$$

4. Consider the function

$$f(u) = \begin{cases} 1 & \text{if } u \leq 0 \\ -1, & \text{if } u > 0. \end{cases}$$

a) Show that the initial value problem

$$\begin{aligned} u' &= f(u) \\ u(0) &= 1 \end{aligned}$$

does not have a solution.

b) Check why Peano's theorem doesn't apply.

$$a) \quad u' = \begin{cases} 1 & , u \leq 0 \\ -1 & , u > 0 \end{cases} \quad , u(0) = 1$$



$u \leq 0$ :

$$\begin{aligned} \text{Ansatz: } u &= x + C \rightarrow u(0) = 1 \rightarrow C = 1 \\ u' &= 1 \end{aligned} \quad \rightarrow \underline{u = x + 1}$$

$u > 0$ :

$$\begin{aligned} \text{Ansatz: } u &= -x + C \\ u' &= -1 \end{aligned}$$

$$\rightarrow u = \begin{cases} \underline{x + 1, u \leq 0} \\ \underline{-x + C, u > 0} \end{cases} \rightarrow$$



$f(u)$  is not continuous,  
hence Peano's theorem  
does not apply.



Continuing with the equation from the previous exercise, let us test some approximation methods.

- a) Compute the first 4 elements in the Picard iteration, and see if it looks convergent.
- b) Another approximation method for ODEs is the *Euler method*, defined as follows. For each  $n$ , we define not a continuous function  $u$ , but rather a function  $\varphi_n$  on gridpoints  $0, \frac{1}{n}, \frac{2}{n}, \dots$ , inductively as follows:

$$\varphi_n(0) = u_0, \quad \varphi_n\left(\frac{k+1}{n}\right) = \varphi_n\left(\frac{k}{n}\right) + \frac{f\left(\frac{k}{n}, u\left(\frac{k}{n}\right)\right)}{n}, \quad k \geq 0.$$

Compute the Euler approximation of the ODE from the previous exercise with  $n = 2, 4, 6$ . Does this look convergent?

$$f(u) = \begin{cases} 1 & \text{if } u \leq 0 \\ -1, & \text{if } u > 0. \end{cases}$$

3. (Picard iteration) The Picard-Lindelöf/Cauchy-Lipschitz theorem in fact gives more than what we discussed in the lecture: it also provides a way to approximate solutions of the initial value problem

$$\begin{aligned} u' &= f(t, u) \\ u(0) &= u_0. \end{aligned}$$

This is the so-called *Picard iteration*, defined as

$$\begin{aligned} \varphi_0(t) &= u_0 \\ \varphi_n(t) &= u_0 + \int_0^t f(s, \varphi_{n-1}(s)) \, ds \quad n \geq 1. \end{aligned}$$

The produced sequence  $(\varphi_n)_{n \in \mathbb{N}}$  converges (provided the conditions in the Picard-Lindelöf theorem hold) to the solution  $u$  of the ODE.

$$u_0 = 1 = \varphi_0 \rightarrow \varphi_0 = 1$$

$$\varphi_1 = 1 + \int_0^t f(1) \, ds = 1 + \int_0^t -1 \, ds = 1 - t$$

$$\varphi_2 = 1 + \int_0^t f(1-s) \, ds = ?$$

if  $t < 1$ :

$$\varphi_2 = 1 + t(-1) = 1 - t$$

$$1 - s > 0 \Rightarrow f(1-s) = -1$$

if  $t \geq 1$ :

$$\varphi_2 = 1 + \int_0^1 f(1-s) \, ds + \int_1^t f(1-s) \, ds$$

$$\varphi_2 = 1 + (-1) + (t-1) = t-1$$

$$\varphi_2 = \begin{cases} 1-t, & t < 1 \\ t-1, & t \geq 1 \end{cases}$$

$$\varphi_3 = \begin{cases} 1 + \int_0^t f(1-s) \, ds, & t < 1 \\ 1 + \int_0^1 f(1-s) \, ds + \int_1^t f(s-1) \, ds, & t \geq 1 \end{cases} = \begin{cases} 1-t, & t < 1 \\ ? & t \geq 1 \end{cases}$$

if  $t \geq 1$ :

$$\begin{aligned} 1 + \int_0^1 f(1-s) \, ds + \int_1^t f(s-1) \, ds &= 1 + 1 + \left( \int_1^t -1 \, ds \right) \\ &= 2 + (-(t-1)) = 3-t \end{aligned}$$

$$\varphi_3 = \begin{cases} 1-t, & t \leq 1 \\ 3-t, & t \geq 1 \end{cases}$$

Does not look convergent (?)

- b) Another approximation method for ODEs is the *Euler method*, defined as follows. For each  $n$ , we define not a continuous function  $u$ , but rather a function  $\varphi_n$  on gridpoints  $0, \frac{1}{n}, \frac{2}{n}, \dots$ , inductively as follows:

$$\varphi_n(0) = u_0, \quad \varphi_n\left(\frac{k+1}{n}\right) = \varphi_n\left(\frac{k}{n}\right) + \frac{f\left(\frac{k}{n}, \varphi_n\left(\frac{k}{n}\right)\right)}{h}, \quad k \geq 0.$$

Compute the Euler approximation of the ODE from the previous exercise with  $n = 2, 4, 6$ . Does this look convergent?

$$\varphi_2 = \varphi_4 = \varphi_6 = u_0 = 1 \quad f(u) = \begin{cases} 1, & \text{if } u \leq 0 \\ -1, & \text{if } u > 0. \end{cases}$$

$h=2$

$$\varphi_2\left(\frac{1}{2}\right) = \varphi_2(0) + \frac{f(0, \varphi_2(0))}{2} = 1 - \frac{1}{2} = \frac{1}{2}$$

$$\varphi_2(1) = \varphi_2\left(\frac{1}{2}\right) + \frac{f\left(\frac{1}{2}, \varphi_2\left(\frac{1}{2}\right)\right)}{2} = \frac{1}{2} - \frac{1}{2} = 0$$

$$\varphi_2\left(\frac{3}{2}\right) = \varphi_2(1) + \frac{f(1, \varphi_2(1))}{2} = 0 + \frac{1}{2} = \frac{1}{2}$$

$$\varphi_2(2) = \varphi_2\left(\frac{3}{2}\right) + \frac{f\left(\frac{3}{2}, \varphi_2\left(\frac{3}{2}\right)\right)}{2} = \frac{1}{2} - \frac{1}{2} = 0$$

...

alternating between 0 and  $\frac{1}{2}$

$$\varphi_n\left(\frac{1}{n}\right) = \varphi_n(0) + \frac{f(0, \varphi_n(0))}{n} = 1 - \frac{1}{n} = \frac{n-1}{n}$$

$$\varphi_n\left(\frac{1}{2}\right) = \varphi_n\left(\frac{1}{n}\right) + \frac{f\left(\frac{1}{2}, \varphi_n\left(\frac{1}{2}\right)\right)}{n} = \frac{n-1}{n} - \frac{1}{n} = \frac{n-2}{n}$$

$$\varphi_n\left(\frac{3}{n}\right) = \dots = \frac{n-2}{n} - \frac{1}{n} = \frac{n-3}{n}$$

$$\varphi_n(1) = \frac{n-3}{n} - \frac{1}{n} = 0$$

$$\varphi_n\left(\frac{5}{n}\right) = 0 + \frac{f(0, \varphi_n(0))}{n} = 0 + \frac{1}{n} = \frac{1}{n}$$

alternating between 0 and  $\frac{1}{n}$

$$\varphi_6\left(\frac{1}{6}\right) = 1 - \frac{1}{6} = \frac{5}{6}$$

$$\varphi_6\left(\frac{2}{6}\right) = \frac{5}{6} - \frac{1}{6} = \frac{4}{6}$$


$$\varphi_6\left(\frac{3}{6}\right) = \frac{3}{6}$$

$$\varphi_6\left(\frac{4}{6}\right) = \frac{2}{6}$$

$$\varphi_6\left(\frac{5}{6}\right) = \frac{1}{6}$$

$$\varphi_6(1) = 0$$

$$\varphi_6\left(\frac{7}{6}\right) = \frac{1}{6}$$



alternating between 0 and  $\frac{1}{6}$

$\rightarrow 1 - t$  for  $t < 1$  and 0 for  $t \geq 1$