1

NCERT Discrete 11.5.9 Q20

EE23BTECH11061 - SWATHI DEEPIKA*

Question: If a,b,c are in A.P.;b,c,d are in G.P and $\frac{1}{c}$, $\frac{1}{d}$, $\frac{1}{e}$ are in A.P. prove that a,c,e are in G.P. **Solution:**

Symbol	Value	Description	Z-Transform
$x_1(n)$	{a, b, c}	A.P Sequence	$X_1(z)$
$x_2(n)$	{b, c, d}	G.P Sequence	$X_2(z)$
$x_3(n)$	$\left\{\frac{1}{c}, \frac{1}{d}, \frac{1}{e}\right\}$	A.P Sequence	$X_3(z)$
<i>y</i> (<i>n</i>)	{a, c, e}	Sequence	Y(z)

TABLE I Parameters

2b = a + c

 $b - a = c - b \tag{1}$

 $c^2 = b \times d \tag{3}$

$$d = \frac{c^2}{h} \tag{4}$$

$$\frac{1}{d} - \frac{1}{c} = \frac{1}{e} - \frac{1}{d}$$
$$\frac{2}{d} = \frac{1}{c} + \frac{1}{e}$$

From (4),

$$\frac{2b}{c^2} = \frac{1}{c} + \frac{1}{e}$$

From (2),

$$\frac{a+c}{c^2} = \frac{1}{c} + \frac{1}{e}$$
 (8)

$$\frac{a}{c^2} + \frac{1}{c} = \frac{1}{c} + \frac{1}{e} \tag{9}$$

$$a \times e = c^2 \tag{10}$$

$$y(1)^2 = y(0) \times y(2) \tag{11}$$

So, y(0),y(1),y(2) are in G.P

1) For y(n):

$$y(n) = y(0) \left(\frac{y(1)}{y(0)}\right)^n u(n)$$
 (12)

$$y(n) \longleftrightarrow Y(z)$$

$$Y(z) = \frac{y(1)}{1 - \frac{y(1)}{y(0)}z^{-1}}, \quad |z| > \left| \frac{y(1)}{y(0)} \right|$$
 (13)

2) For $x_1(n)$:

$$x_1(n) = (x_1(0) + n(x_1(1) - x_1(0)))u(n)$$
 (14)

$$x_1(n) \longleftrightarrow X_1(z)$$

$$X_1(z) = \frac{x_1(0)}{1 - z^{-1}} + \frac{(x_1(1) - x_1(0))z^{-1}}{(1 - z^{-1})^2}, \quad |z| > 1$$
(15)

3) For $x_2(n)$:

(2)

(6)

(7)

$$x_2(n) = x_2(0) \left(\frac{x_2(1)}{x_2(0)}\right)^n u(n)$$
 (16)

$$x_2(n) \longleftrightarrow X_2(z)$$

$$X_2(z) = \frac{x_2(1)}{1 - \frac{x_2(1)}{x_2(0)}z^{-1}}, \quad |z| > \left|\frac{x_2(1)}{x_2(0)}\right|$$
 (17)

(5) 4) For $x_3(n)$:

$$x_3(n) = (x_3(0) + n(x_3(0) - x_3(1))) u(n)$$
 (18)

$$x_3(n) \longleftrightarrow X_3(z)$$

$$X_3(z) = \frac{x_3(0)}{1 - z^{-1}} + (x_3(1) - x_3(0)) \frac{z^{-1}}{(1 - z^{-1})^2}, \quad |z| > 1$$
(19)