

# Assignment 5

## Probability and Random Variables

Swati Mohanty (EE20RESCH11007)

### I. PROBLEM

In a game, a man wins a rupee for a six and loses a rupee for any other number when a fair die is thrown. The man decided to throw a die thrice but to quit as and when he gets a six. Find the expected value of the amount he wins / loses.

### II. SOLUTION

Let  $X$  denote the random variable of winning/losing in the game.

$$P(X = 1) = P(\text{win}) = p = \frac{1}{6} = 0.167$$

$$P(X = 0) = P(\text{loss}) = 1 - p = \frac{5}{6} = 0.833.$$

Let  $Y$  denote the random variable of winning the game in  $N$ th trial,  $A(Y)$  denote the amount in each case. The possible cases are: (i) Wins in first throw:

$$P(Y = 1) = p = 0.167 \quad (1)$$

$$A(Y = 1) = +1 \quad (2)$$

(ii) Wins in the second throw :

$$P(Y = 2) = (1 - p) \times p = 0.139 \quad (3)$$

$$A(Y = 2) = -1 + 1 = 0 \quad (4)$$

(iii) Wins in the third throw :

$$P(Y = 3) = (1 - p) \times (1 - p) \times p = 0.107 \quad (5)$$

$$A(Y = 3) = -1 - 1 + 1 = -1 \quad (6)$$

(iv) Does not win in any throw :

$$P(Y = 3) = (1 - p) \times (1 - p) \times (1 - p) = 0.596 \quad (7)$$

$$A(Y = 3) = -1 - 1 - 1 = -3 \quad (8)$$

Net amount = Expectation =  $E[Y]$

$$E[Y] = \sum_{n=1}^3 P(Y = n)A(Y = n) \quad (9)$$

$$0.167 \times 1 + 0 + 0.107 \times -1 + 0.596 \times -3 \quad (10)$$

$$= -1.73 \quad (11)$$

```

Simulated results
0.167073
0.13918
0.13918
0.833175
-2.471632
Theoretical results
0.165
0.132
0.107
0.596
-1.7299999999999998

```

Figure 1: Simulation for tossing a fair coin

The probabilities were simulated using the python code.

**Download python code from here**

[https://github.com/Swati-Mohanty/AI5002/blob/main/Assignment\\_5/codes/die.py](https://github.com/Swati-Mohanty/AI5002/blob/main/Assignment_5/codes/die.py)

**Download latex code from here-**

[https://github.com/Swati-Mohanty/AI5002/blob/main/Assignment\\_5/codes/assignment5.tex](https://github.com/Swati-Mohanty/AI5002/blob/main/Assignment_5/codes/assignment5.tex)