On the properties of projectile motion and quadratic air resistance

Simon Halvdansson

April 18, 2016

The equations of motion for normal projectile motion are derived and some properties of such motion are determined. One dimensional quadratic air resistance is studied and modeled using numerical and analytical methods and compared to experimental results, producing negligible discrepancies. Analytical approximations to two dimensional quadratic air resistance are made using a model which separates low, high and split angle trajectories. The equations of motion for these are derived, plotted and compared to numerical solutions to the original differential equation.

1	Intro	oduction	1
2	Traj 2.1 2.2 2.3 2.4	Acceleration	1 2 2 2 2
3	Prop 3.1 3.2 3.3 3.4 3.5 3.6	Time to highest point	2 3 3 3 3 3
4	Cho 4.1 4.2 4.3	osing initial values to match result values Delimitations	4 4 4 4 5
5	Long 5.1 5.2	gest parabola starting at height Expression for range	5 5
6	Con 6.1 6.2	Change at every time interval	6 6 7
7	Ana 7.1 7.2	lytical solution to one-dimensional air resistanceDerivation7.1.1 Position as a function of time7.1.2 Determining the duration of the fall7.1.3 Terminal velocityPlots7.2.1 Examples7.2.2 Analysis	7 7 8 8 8 8 9
8	Ana 8.1 8.2	Introduction	9 10 10 10

		8.2.3 Position	10
		8.2.4 Equations of motion	11
	8.3	High Angle Trajectory approximation	11
		8.3.1 Differential equations of motion	11
		8.3.2 Ascending velocity	11
		8.3.3 Ascending position	11
		8.3.4 Descending velocity	12
		8.3.5 Descending position	12
		8.3.6 Equations of motion	13
	8.4	Split Angle Trajectory approximation	13
		8.4.1 Differential equations of motion	13
		8.4.2 Equations of motion	13
	8.5	Plots	13
		8.5.1 Low Angle Trajectory	13
		8.5.2 High Angle Trajectory	14
		8.5.3 Split Angle Trajectory	14
9	Ехр	eriment	14
	9.1	Setup	14
	9.2	Results	15
	9.3	Analysis	15
10	Con	culsions	15
	Furt	ther reading	15