On the properties of projectile motion and quadratic air resistance

Simon Halvdansson

March 22, 2016

The equations of motion for normal projectile motion are derived. Some properties of such motion are defined. Numerical approximations and analytical solutions to the effects of one dimensional quadratic air resistance are made and compared to experimental results with small error. Analytical approximations to two dimensional quadratic air resistance are made using a model which separates low, high and split angle trajectories. The equations of motion for these are derived, plotted and compared to numerical solutions to the original differential equation.

1	Intr	Introduction				
2	Traj 2.1 2.2 2.3 2.4	Acceleration	4 5 5 5 5			
3	Prop 3.1 3.2 3.3 3.4 3.5 3.6	Highest point				
4	Cho 4.1 4.2 4.3	Delimitations Specific projectile range Hitting a specific coordinate 4.3.1 Predetermined speed 4.3.2 Predetermined angle	6 7 7 7 7			
5	Lon ₅ 5.1 5.2	gest parabola starting at height Expression for range				
6	Con 6.1 6.2	nputational approach to air resistance Change at every time interval				
7	Ana 7.1 7.2	Ilytical solution to one-dimensional air resistance Derivation	10 10 10 11 11 11 11 12			
8	Ana 8.1 8.2	Ilytical approximations to two-dimensional air resistance Introduction	12 12 13 13 13			

		8.2.3	Position	. 13		
		8.2.4	Equations of motion	. 14		
	8.3	High A	Angle Trajectory approximation	. 14		
		8.3.1	Differential equations of motion	. 14		
		8.3.2	Ascending velocity	. 14		
		8.3.3	Ascending position	. 14		
		8.3.4	Descending velocity	. 15		
		8.3.5	Descending position			
		8.3.6	Equations of motion			
	8.4	Split A	Angle Trajectory approximation	. 16		
		8.4.1	Differential equations of motion			
		8.4.2	Equations of motion	. 16		
	8.5	Plots		. 16		
		8.5.1	Low Angle Trajectory	. 16		
		8.5.2	High Angle Trajectory	. 17		
		8.5.3	Split Angle Trajectory			
9	Experiment					
	9.1			. 17		
	9.2		S			
	9.3		sis			
10	Conculsions					
	Further reading					