On the properties of projectile motion and quadratic air resistance

Simon Halvdansson

March 21, 2016

The equations of motion for normal projectile motion are derived. Some properties of such motion are defined. Numerical approximations and analytical solutions to the effects of one dimensional quadratic air resistance are made and compared to experimental results with small error. Analytical approximations to two dimensional quadratic air resistance are made using a model which separates low, high and split angle trajectories. The equations of motion for these are derived, plotted and compared to numerical solutions to the original differential equation.

1	Intr	oduction	1
2	Traj 2.1 2.2 2.3 2.4	Acceleration	1 2 2 2 2
3	Prop 3.1 3.2 3.3 3.4 3.5 3.6	Time to highest point	2 3 3 3 3 3
4	Cho 4.1 4.2 4.3	osing initial values to match result values Delimitations Specific projectile range Hitting a specific coordinate 4.3.1 Predetermined speed 4.3.2 Predetermined angle	3 4 4 4 4
5	Lon s 5.1 5.2	gest parabola starting at height Expression for range	5 5
6	Con 6.1 6.2	Change at every time interval	6 6 7
7	Ana 7.1 7.2	lytical solution to one-dimensional air resistanceDerivation7.1.1 Position as a function of time7.1.2 Determining the duration of the fall7.1.3 Terminal velocityPlots7.2.1 Examples7.2.2 Analysis	7 7 8 8 8 8 9
8	Ana 8.1 8.2	8.2.1 Differential equations of motion	9 10 10

		8.2.3 Position	10
		8.2.4 Equations of motion	11
	8.3	High Angle Trajectory approximation	11
		8.3.1 Differential equations of motion	11
		8.3.2 Ascending velocity	11
		8.3.3 Ascending position	11
		8.3.4 Descending velocity	12
		8.3.5 Descending position	12
		8.3.6 Equations of motion	13
	8.4	Split Angle Trajectory approximation	13
		8.4.1 Differential equations of motion	13
		8.4.2 Equations of motion	13
	8.5	Plots	13
		8.5.1 Low Angle Trajectory	13
		8.5.2 High Angle Trajectory	14
		8.5.3 Split Angle Trajectory	14
9	Ехр	eriment	14
	9.1	Setup	14
	9.2	Results	15
	9.3	Analysis	15
10	Con	culsions	15
	Furt	ther reading	15