On the properties of projectile motion and quadratic air resistance

Simon Halvdansson

March 7, 2016

The equations of motion for normal projectile motion are derived. Some properties of such motion are defined. Numerical approximations and analytical solutions to the effects of one dimensional quadratic air resistance are made and compared to experimental results with small error. Analytical approximations to two dimensional quadratic air resistance are made using a model which separates low, high and split angle trajectories. The equations of motion for these are derived, plotted and compared to numerical solutions to the original differential equation.

1	Introduction	4
2	Trajectory 2.1 Acceleration	
3	Properties of a projectile parabola 3.1 Time to highest point	5 6 6 6 6
4	Choosing initial values to match result values 4.1 Delimitations	6 7 7 7
5	Longest parabola starting at height 5.1 Expression for range	8 8
6	Computational approach to air resistance 6.1 Change at every time interval	9 10
7	Analytical solution to one-dimensional air resistance 7.1 Derivation 7.1.1 Position as a function of time 7.1.2 Determining the duration of the fall 7.1.3 Terminal velocity 7.2 Plots 7.2.1 Examples 7.2.2 Analysis	10 10 11 11 11 11 12
8	. y 	12 13 13 13

		8.2.3 Position	13
		8.2.4 Equations of motion	14
	8.3	High Angle Trajectory approximation	14
		8.3.1 Differential equations of motion	14
		8.3.2 Ascending velocity	14
		8.3.3 Ascending position	14
		8.3.4 Descending velocity	15
		8.3.5 Descending position	15
		8.3.6 Equations of motion	16
	8.4	Split Angle Trajectory approximation	16
		8.4.1 Differential equations of motion	16
		8.4.2 Equations of motion	16
	8.5	Plots	16
		8.5.1 Low Angle Trajectory	16
		8.5.2 High Angle Trajectory	17
		8.5.3 Split Angle Trajectory	17
9	Ехр	eriment	17
	9.1	Setup	17
	9.2	Results	18
	9.3	Analysis	18
10	Con	culsions	18
	Furt	ther reading	18