

Transgenerational Developmental and Behavioral Plasticity of Threespine Stickleback *Gasterosteus aculeatus* of Lake Myvatn, Iceland

Spencer Edwards, Roll No.

April 2021

Introduction

Much research has been conducted exploring phenotypic plasticity, the ability for an organism to alter its phenotype in response to its environment within its lifetime (Denver and Middlemis-Maher, 2010; Kishida et al., 2010; Klemetsen, 2010). However, there is increasing interest in so-called transgenerational plasticity (TGP), wherein plastic responses to environmental conditions are passed down to offspring (Hellmann et al., 2020; Richter-Boix et al., 2014; Bell and Hellmann, 2019; Shama et al., 2014). This concept holds particular importance in the face of rapid environmental change, because it allows organisms to evolve without relying on actual genetic changes. One of the most well-known model systems for studying plasticity is *Gasterosteus aculeatus*, commonly named the threespine stickleback. Stickleback populations often form species pairs, with two or more morphs coexisting in a single body of water via niche partitioning. In lake Myvatn, two morphs exist, the “mud” and “lava” morphs, defined mainly by habitat type, as well as morphology and behavior (Kristjánsson et al., 2002; Millet et al., 2013). Different sections of lake Myvatn act as distinct habitats (referred to as the North, South, and (sometimes) East basins), and so the sticklebacks can also be distinguished by habitat of origin (Millet et al., 2013; Einarsson et al., 2004). Millet et al. (2013) in particular found that there was a large difference in size between sticklebacks from different habitats, with those of the

North basin being larger ($\geq 55\text{mm}$) and of different age classes than those of the South. They posit that these differences could be due to differences in life history strategies between the populations, or plasticity in response to resources. Interestingly, they also note that North basin stickleback have larger spines than those of the South basin, possibly owing to differences in abundance of gape-limited predators, which occur at rates of up ten times higher in the North basin than the south (Millet et al., 2013). While we know that plasticity in traits does occur in lake Myvatn stickleback, there is little data on whether this plasticity is transgenerational. Indeed, much of the research on TGP in stickleback has been performed on oceanic populations, and thus TGP in freshwater stickleback represents a current gap in our knowledge of the ecology and evolution of these organisms (Shama et al., 2014). Evidence from marine stickleback suggests that TGP does occur, although it varies between populations (Shama et al., 2014; Heckwolf et al., 2018; Kozak and Boughman, 2012)

Proposed Guides: Dr. Bjarni Kristjánsson, Department of Aquaculture and Fish Biology, Hólar University College.

References

- Bell, A. M. and Hellmann, J. K. (2019). An Integrative Framework for Understanding the Mechanisms and Multigenerational Consequences of Transgenerational Plasticity. *Annual Review of Ecology, Evolution, and Systematics*, 50:97–118.
- Denver, R. J. and Middlemis-Maher, J. (2010). Lessons from evolution: Developmental plasticity in vertebrates with complex life cycles. *Journal of Developmental Origins of Health and Disease*, 1(5):282–291.
- Einarsson, Á., Stefánsdóttir, G., Jóhannesson, H., Ólafsson, J. S., Már Gíslason, G., Wakana, I., Gudbergsson, G., and Gardarsson, A. (2004). The ecology of Lake Myvatn and the River Laxá: Variation in space and time. *Aquatic Ecology*, 38(2):317–348.
- Heckwolf, M. J., Meyer, B. S., Döring, T., Eizaguirre, C., and Reusch, T. B. (2018). Transgenerational plasticity and selection shape the adaptive potential of sticklebacks to salinity change. *Evolutionary Applications*, 11(10):1873–1885.

- Hellmann, J. K., Carlson, E. R., and Bell, A. M. (2020). Sex-specific plasticity across generations II: Grandpaternal effects are lineage specific and sex specific. *Journal of Animal Ecology*, 89(12):2800–2812.
- Kishida, O., Trussell, G. C., Mougi, A., and Nishimura, K. (2010). Evolutionary ecology of inducible morphological plasticity in predator-prey interaction: Toward the practical links with population ecology. *Population Ecology*, 52(1):37–46.
- Klemetsen, A. (2010). The Charr Problem Revisited: Exceptional Phenotypic Plasticity Promotes Ecological Speciation in Postglacial Lakes. *Freshwater Reviews*, 3(1):49–74.
- Kozak, G. M. and Boughman, J. W. (2012). Plastic responses to parents and predators lead to divergent shoaling behaviour in sticklebacks. *Journal of Evolutionary Biology*, 25(4):759–769.
- Kristjánsson, B. K., Skúlason, S., and Noakes, D. L. (2002). Morphological segregation of Icelandic threespine stickleback (*Gasterosteus aculeatus* L). *Biological Journal of the Linnean Society*, 76(2):247–257.
- Millet, A., Kristjánsson, B. K., Einarsson, Á., and Räsänen, K. (2013). Spatial phenotypic and genetic structure of threespine stickleback (*Gasterosteus aculeatus*) in a heterogeneous natural system, Lake Myvatn, Iceland. *Ecology and Evolution*, 3(10):3219–3232.
- Richter-Boix, A., Orizaola, G., and Laurila, A. (2014). Transgenerational phenotypic plasticity links breeding phenology with offspring life-history. *Ecology*, 95(10):2815–2825.
- Shama, L. N., Strobel, A., Mark, F. C., and Wegner, K. M. (2014). Transgenerational plasticity in marine sticklebacks: Maternal effects mediate impacts of a warming ocean. *Functional Ecology*, 28(6):1482–1493.