Hydrodynamic Topology as an Effective Model for Inertial Mass: A Self-Contained Derivation of a Topology-Modulated Mass Functional

Omar Iskandarani[*]

February 14, 2026

Abstract

We present a self-contained, hydrodynamically motivated effective model in which the invariant rest mass of a localized, topologically stable excitation is identified with the integral of a localized kinetic-energy density over its effective support volume. The model assumes an incompressible, inviscid background medium (Euler–Kelvin regime) supporting persistent filamentary defects whose stability is protected by circulation conservation. The resulting rest-mass functional factorizes into: (i) a universal energy-density scale $u = \frac{1}{2}\rho_{\mathrm{core}}\|\mathbf{v}_{\circlearrowleft}\|^2$, (ii) a geometric volume proxy $V = \pi r_c^3 L_{\mathrm{tot}}$, and (iii) a dimensionless kernel $\mathcal{K}(T)$ depending only on a small set of topological indices. We motivate a minimal closed kernel of the form

$$M(T) = \mathcal{K}(T)\,\frac{u\,V(T)}{c^2}, \qquad \mathcal{K}(T) = \left(\frac{4}{\alpha}\right)^{\mathsf{G}(T)} k(T)^{-3/2}\,\varphi^{-g(T)}\,n(T)^{-1/\varphi}, \tag{1}$$

where k is a complexity index (e.g. braid index/crossing proxy), g is Seifert genus, n is the number of components, α is the fine-structure constant, and φ is the golden ratio. The paper is written to be "orthodox-friendly": all strong statements are framed as model assumptions, the status of each scaling law is clearly labeled, and falsifiable preferred-frame and spectrum-selection consequences are listed. We also separate background density ρ_f (governing wave transport) from a core-localized density ρ_{core} (governing inertial mass) and explain why this separation is required for dimensional consistency and empirical scale matching.

[*] **Independent Researcher, Groningen, The Netherlands**
Email: info@omariskandarani.com
ORCID: 0009-0006-1686-3961
DOI: 10.5281/zenodo.xxx

1 Introduction and scope

The microscopic origin of inertial mass remains an open question spanning particle physics, condensed matter analogies, and quantum field theory. In the Standard Model (SM), fermion masses arise from Yukawa couplings to the Higgs condensate, with parameters not predicted by the theory [1, 2, 3]. This motivates exploration of alternative *effective* mechanisms in which inertia emerges from localized energy storage and topological stability.

This article develops a hydrodynamic-topology effective model where: (i) persistent localized excitations are protected by circulation conservation (Kelvin theorem), (ii) their rest mass equals a localized kinetic-energy content divided by c^2, (iii) topology modulates the efficiency of energy localization through a compact kernel $\mathcal{K}(T)$.

Status note. Equation (1) is presented as a *closed effective ansatz* motivated by Euler–Kelvin hydrodynamics, slender-filament energetics, and discrete scale invariance considerations. Certain exponent choices (notably $-3/2$) admit supporting scaling arguments; others (golden-ratio exponents) are principled but not unique. The goal is a journal-ready, checkable formulation with explicit assumptions and falsifiers.

2 Hydrodynamic substrate and topological stability

2.1 Euler–Kelvin regime

Assume a background medium described (on relevant scales) as incompressible and inviscid:

$$\nabla \cdot \mathbf{v} = 0, \qquad \partial_t \mathbf{v} + (\mathbf{v} \cdot \nabla)\mathbf{v} = -\frac{1}{\rho}\nabla p, \tag{2}$$

with barotropic closure $p = p(\rho)$ locally. In this regime, Kelvin's circulation theorem implies that the circulation along a material loop is conserved:

$$\Gamma = \oint_{\mathcal{C}(t)} \mathbf{v} \cdot d\boldsymbol{\ell} = \text{const.} \tag{3}$$

Consequently, filamentary circulation structures cannot smoothly terminate; closed loops are dynamically preferred and topologically robust [4, 5, 6].

2.2 Filamentary defects as persistent excitations

We model a localized excitation T as a closed filament (or link of n components) with a microscopic core radius r_c and a characteristic tangential swirl speed $\|\mathbf{v}_{\circlearrowleft}\| = \|\mathbf{v}_{\circlearrowleft}\|$. The physical picture is a slender tube of radius r_c carrying a quantized circulation scale Γ_0, which we parameterize as

$$\Gamma_0 \equiv 2\pi r_c \|\mathbf{v}_{\circlearrowleft}\|. \tag{4}$$

Equation (4) is a definition of the characteristic scale; quantization of Γ is treated as an additional postulate compatible with quantized vortices in superfluids [7, 8].

3 Mass as localized kinetic energy

3.1 Covariant definition (orthodox framing)

In relativistic field theory, invariant rest mass is the rest-frame energy divided by c^2. We adopt the operational definition: in the excitation rest frame,

$$M \equiv \frac{1}{c^2} \int_V \rho_E(\mathbf{x})\, dV, \tag{5}$$

where ρ_E is the localized energy density attributed to the excitation and V encloses its support. This is a standard bookkeeping identity once ρ_E is defined.

3.2 Core-localized energy density scale

Assume the excitation has a core region characterized by an effective core density ρ_{core} (distinct from the background density ρ_f that governs long-wavelength wave transport). The simplest kinetic-energy density scale is then

$$u \equiv \frac{1}{2}\rho_{\text{core}}\|\mathbf{v}_{\circlearrowleft}\|^2, \tag{6}$$

with units $\mathrm{J\,m^{-3}}$. We emphasize: (6) is the *local* kinetic-energy density of the organized core motion, not the background vacuum energy.

Why separate ρ_f and ρ_{core}? If one attempted to build particle-scale masses by integrating background density ρ_f over microscopic volumes, the resulting scale is generically far too small. The effective model requires a localized "core phase" with much larger inertial density to achieve observed masses while keeping the background "light." This is analogous in spirit (not identity) to the separation between condensate order-parameter stiffness and quasiparticle effective masses in superfluid/superconducting systems.

4 Geometry: volume proxy from ropelength

4.1 Slender-filament volume approximation

For a slender tube of radius r_c and centerline length ℓ, the volume is approximately

$$V \simeq \pi r_c^2\,\ell. \tag{7}$$

Write $\ell = r_c\,L_{\text{tot}}$, where L_{tot} is a dimensionless ropelength-like quantity (e.g. length in units of r_c). Then

$$V(T) = \pi r_c^3\,L_{\text{tot}}(T). \tag{8}$$

The use of a minimal (tight) ropelength as a ground-state proxy is consistent with the idea that line tension drives contraction until self-avoidance/self-interaction prevents further shrinkage [6].

5 Topology: a minimal closed kernel

5.1 Factorized master formula

Combining (5), (6), and (8), we obtain the factorized form

$$M(T) = \mathcal{K}(T)\,\frac{u\,V(T)}{c^2} = \mathcal{K}(T)\,\frac{\left(\frac{1}{2}\rho_{\text{core}}\|\mathbf{v}_{\circlearrowleft}\|^2\right)\left(\pi r_c^3 L_{\text{tot}}(T)\right)}{c^2}. \tag{9}$$

All nontrivial dependence on topology and "energy-localization efficiency" is now placed into $\mathcal{K}(T)$.

5.2 Minimal kernel choice and interpretation

We consider the dimensionless kernel

$$\mathcal{K}(T) = \left(\frac{4}{\alpha}\right)^{\mathsf{G}(T)} k(T)^{-3/2}\,\varphi^{-g(T)}\,n(T)^{-1/\varphi}. \tag{10}$$

Here:

- $k(T)$ is a complexity index (e.g. braid index, or a crossing-number proxy) representing coarse geometric/topological complexity relevant for self-induction and screening.
- $g(T)$ is Seifert genus, capturing surface complexity of a spanning surface.
- $n(T)$ is number of components (for links).
- $\varphi = \frac{1+\sqrt{5}}{2}$ is the golden ratio, used as a discrete-scale modulation factor (see below).
- α is the electromagnetic fine-structure constant, used here as a universal dimensionless normalization connecting energy localization to a measured coupling strength; we treat this as a model postulate rather than a theorem.

Status of each factor.

1. The factor $(4/\alpha)^{\mathsf{G}(T)}$ is a *gated normalization postulate.* When $\mathsf{G}(T) = 0$, the excitation is in a "shielded" sector and the vacuum-coupling amplification is inactive; when $\mathsf{G}(T) = 1$, the amplification is active.
2. The exponent $-3/2$ for $k(T)$ admits scaling support from diffusion/heat-kernel analogies and from dimensional considerations in filament energetics; nevertheless, it remains an effective choice.
3. The φ^{-g} and $n^{-1/\varphi}$ factors encode *discrete scale invariance (DSI)*-motivated suppression with increasing topological obstruction and multiplicity. These are principled but not unique; alternative monotone suppressions can be tested.

5.3 Formal shielding gate via rational sliceness (hard constraint)

The narrative "shielding vs. exposure" is promoted here to a *checkable gate* $\mathsf{G}(T) \in \{0, 1\}$ defined using rational sliceness (rational homology 4-ball bounding). This provides an explicit on/off rule for when the factor $4/\alpha$ is active.

Definition 5.1 (Rational homology 4-ball)**.** A smooth compact 4-manifold W is a *rational homology 4-ball* if $\partial W \cong S^3$ and

$$H_*(W; \mathbb{Q}) \cong H_*(B^4; \mathbb{Q}).$$

Definition 5.2 (Rational sliceness)**.** A knot $K \subset S^3$ is *rationally slice* if there exists a rational homology 4-ball W with $\partial W \cong S^3$ in which K bounds a smoothly embedded disk $D^2 \subset W$. Equivalently, K is slice in some W that is "B^4 up to $\mathbb{Q}$-homology." [12]

Definition 5.3 (Shielding gate)**.** Define the *shielding-gate exponent* $\mathsf{G}(T) \in \{0, 1\}$ by

$$\mathsf{G}(T) = \begin{cases} 0, & \text{if } T \text{ is rationally slice (shielded sector)}, \\ 1, & \text{otherwise (exposed sector)}. \end{cases}$$

In particular, $(4/\alpha)^{\mathsf{G}(T)} = 1$ in the shielded sector and equals $4/\alpha$ in the exposed sector.

Double-twist family as an explicit, computable gate. For the double twist knots $K_{m,n}$ (two twist regions with integer parameters m, n), rational sliceness is completely characterized. We use this as a *hard constraint* (not merely inspiration) because it yields an exact, finite test for $\mathsf{G}(K_{m,n})$.

Theorem 5.1 (Rational sliceness of double twist knots [13]). *A double twist knot $K_{m,n}$ is rationally slice if and only if*

$$mn = 0 \quad or \quad n = -m \quad or \quad n = -m \pm 1.$$

Remark 5.1 (Gate specialization for $K_{m,n}$). The gate becomes the explicit indicator of *not* satisfying the above condition:

$$\mathsf{G}(K_{m,n}) = \begin{cases} 0, & mn = 0 \text{ or } n = -m \text{ or } n = -m \pm 1, \\ 1, & \text{otherwise.} \end{cases}$$

The proof in [13] uses definite 4-manifold obstructions (Donaldson-type input) to rule out all other cases. [14]

5.4 DSI motivation (orthodox phrasing)

Discrete scale invariance appears in multiple contexts (fracture, turbulence cascades, renormalization-group limit cycles) and can generate preferred ratios in spectra [11]. In the present model, we adopt DSI as an organizing principle: increased topological obstruction (g) and component count (n) reduces the fraction of volume that can maintain coherent core motion, implemented via the golden-ratio suppression in (10). This is a hypothesis, testable by spectrum-fitting stability and by predictive out-of-sample checks.

6 Representative selection vs. mass kernel

Equation (9) depends on $L_{\text{tot}}(T)$, which in practice requires selecting a representative configuration within a topological sector. To keep the mass kernel (10) minimal, we separate:

1. **Selection functional:** choose a representative $K^\star$ within a sector by minimizing an effective topological energy

$$E_{\text{eff}}[K] = \alpha_C\, C(K) + \beta_L\, L(K) + \gamma_H\, H(K), \qquad K^\star = \arg\min E_{\text{eff}}[K]. \tag{11}$$

   Here C, L, H are monotone complexity/stability proxies; $\alpha_C, \beta_L, \gamma_H$ are weights. This stage defines which "tight" representative is used to compute L_{tot} and indices.

2. **Mass evaluation:** compute $(k, g, n, L_{\text{tot}})$ on $K^\star$ and evaluate (1).

This separation prevents double-counting complexity inside $\mathcal{K}$ and keeps the mass map compact.

7 Calibration, prediction program, and limitations

7.1 Calibration boundary

Any model with a universal scale must be anchored. A minimal approach is to calibrate one reference mass (e.g. the electron) by choosing L_{tot} of its representative so that (1) matches M_e. All other masses become predictions conditional on the selected topological assignments and computed indices/ropelengths.

7.2 Composite systems and binding energy

For nuclei and atoms, a first-pass assembly model is

$$M_{\text{atom}}(Z, N) \approx Z\, M_p + N\, M_n + Z\, M_e - \frac{E_{\text{bind}}(Z, N)}{c^2}, \tag{12}$$

where each constituent mass (e.g. M_p, M_n, M_e) is supplied by the same master kernel $M(T)$ evaluated on the chosen topological representatives, while E_{bind} is provided by a *bridge model* until a fully hydrodynamic binding theory is derived.

7.2.1 SEMF bridge model (drop-in compatible with the master kernel)

As an orthodox-accurate bridge, take the Bethe–Weizsäcker semi-empirical mass formula (SEMF) binding energy:

$$E_{\text{bind}}^{\text{SEMF}}(Z,N) = a_V A - a_S A^{2/3} - a_C \frac{Z(Z-1)}{A^{1/3}} - a_A \frac{(A-2Z)^2}{A} + \delta(A,Z), \qquad A = Z+N, \tag{13}$$

with the pairing term

$$\delta(A,Z) = \begin{cases} +a_P A^{-1/2}, & Z,N \text{ even}, \\ 0, & A \text{ odd}, \\ -a_P A^{-1/2}, & Z,N \text{ odd}. \end{cases} \tag{14}$$

This keeps the atomic/nuclear *bookkeeping* orthodox while delegating the *constituent rest masses* to the topology-modulated kernel (1). [9, 10]

7.2.2 Optional SST-compatible modulation (bridge extension)

If one wishes to encode "topological packing efficiency" at the nuclear scale without changing the SEMF structure, promote (a_V, a_S) to mild A-dependent functions through a monotone coherence factor $\Xi(A)$ (fit/validated on data):

$$a_V(A) = a_V^{(0)}\,\Xi(A), \qquad a_S(A) = a_S^{(0)}\,\Xi(A)^{-1/2}, \tag{15}$$

leaving a_C, a_A, a_P as orthodox baselines at first pass. This yields an SST-SEMF bridge that remains algebraically identical to (13) but allows controlled deviations for large A in out-of-sample tests.

7.3 Preferred-frame (foliation) extension and falsifiers

If the background medium defines a preferred rest frame, motion relative to it can generate small preferred-frame corrections. Parametrize the effective mass for an excitation moving at velocity v relative to the medium by

$$M_{\text{eff}}(v) = M_0 \left[1 + \eta_1 \frac{\mathbf{v}\cdot\hat{\mathbf{u}}}{c} + \eta_2 \frac{(\mathbf{v}\cdot\hat{\mathbf{u}})^2}{c^2} + \cdots \right], \tag{16}$$

where $\hat{\mathbf{u}}$ is the medium rest-frame direction field and η_i map to preferred-frame phenomenology constrained by precision tests (PPN/SME literature) [16, 17]. The model is falsified if required η_i violate existing bounds or if predicted anisotropies are not observed.

8 Numerical sanity check (dimension and scale)

Define the universal prefactor

$$M_0 \equiv \frac{u\,\pi r_c^3}{c^2} = \frac{\left(\frac{1}{2}\rho_{\text{core}}\|\mathbf{v}_{\circlearrowleft}\|^2\right)\pi r_c^3}{c^2}, \qquad M_1 \equiv \left(\frac{4}{\alpha}\right) M_0. \tag{17}$$

Then

$$M(T) = M_0 \left(\frac{4}{\alpha}\right)^{\mathsf{G}(T)} \left[k(T)^{-3/2}\,\varphi^{-g(T)}\,n(T)^{-1/\varphi} \right] L_{\text{tot}}(T). \tag{18}$$

Equation (18) makes explicit that the model requires only a universal base mass scale M_0, a *gated* amplification $(4/\alpha)^{\mathsf{G}(T)}$, and dimensionless topology/geometry factors.

10-year-old analogy (one sentence). A particle is modeled like a tiny knotted loop of spinning fluid: its mass is how much "spinning energy" is packed into the loop, and complicated knots pack that energy less efficiently.

9 Conclusions

We have presented a self-contained hydrodynamic-topology effective model for inertial mass in which rest mass equals localized kinetic energy divided by c^2, and topology enters through a minimal dimensionless kernel multiplying a filament volume proxy. The framework is explicitly modular: (i) covariant mass functional, (ii) core energy density scale, (iii) slender-filament geometry, (iv) topology-modulated localization efficiency, and (v) optional representative selection by an energy functional. The approach yields clear falsifiers (preferred-frame constraints, out-of-sample mass predictions, stability of selection under kernel variations) and provides a compact, checkable formula suitable for systematic benchmark testing.

Acknowledgments

None.

References

[1] P. W. Higgs (1964), *Broken symmetries and the masses of gauge bosons*, Phys. Rev. Lett. **13**, 508–509. doi:10.1103/PhysRevLett.13.508

[2] F. Englert and R. Brout (1964), *Broken symmetry and the mass of gauge vector mesons*, Phys. Rev. Lett. **13**, 321–323. doi:10.1103/PhysRevLett.13.321

[3] S. Weinberg (1967), *A model of leptons*, Phys. Rev. Lett. **19**, 1264–1266. doi:10.1103/PhysRevLett.19.1264

[4] W. Thomson (Lord Kelvin) (1869), *On vortex motion*, Trans. R. Soc. Edinburgh **25**, 217–260. (permalink: historical reprints widely available)

[5] G. K. Batchelor (1967), *An Introduction to Fluid Dynamics*, Cambridge University Press.

[6] P. G. Saffman (1992), *Vortex Dynamics*, Cambridge University Press.

[7] L. Onsager (1949), *Statistical hydrodynamics*, Nuovo Cimento Suppl. **6**, 279–287. (permalink: archival)

[8] R. P. Feynman (1955), *Chapter II: Application of quantum mechanics to liquid helium*, in *Progress in Low Temperature Physics*, Vol. 1 (ed. C. J. Gorter), Elsevier, pp. 17–53.

[9] C. F. von Weizsäcker (1935), *Zur Theorie der Kernmassen*, Z. Phys. **96**, 431–458. doi:10.1007/BF01337700

[10] H. A. Bethe and R. F. Bacher (1936), *Nuclear Physics A: Stationary States of Nuclei*, Rev. Mod. Phys. **8**, 82–229. doi:10.1103/RevModPhys.8.82

[11] D. Sornette (1998), *Discrete-scale invariance and complex dimensions*, Phys. Rep. **297**, 239–270. doi:10.1016/S0370-1573(97)00076-8

[12] J. C. Cha (2007), *The structure of the rational concordance group of knots*, Mem. Amer. Math. Soc. **189**, no. 885. doi:10.1090/memo/0885 (arXiv:math/0609408)

[13] W. Lee (2025), *Rational Concordance of Double Twist Knots*, arXiv:2504.07636. doi:10.48550/arXiv.2504.07636

[14] S. K. Donaldson (1983), *An application of gauge theory to four-dimensional topology*, J. Differential Geom. **18**(2), 279–315. doi:10.4310/jdg/1214437665

[15] J. Cantarella, R. B. Kusner, and J. M. Sullivan (2002), *On the minimum ropelength of knots and links*, Invent. Math. **150**, 257–286. doi:10.1007/s00222-002-0234-y

[16] C. M. Will (2014), *The Confrontation between General Relativity and Experiment*, Living Rev. Relativ. **17**, 4. doi:10.12942/lrr-2014-4

[17] V. A. Kostelecký (2004), *Gravity, Lorentz violation, and the Standard Model*, Phys. Rev. D **69**, 105009. doi:10.1103/PhysRevD.69.105009