rmNoto Sans

Thermodynamic Reinterpretation of the Golden Mass Hierarchy in Swirl String Theory

Omar Iskandarani*

Independent Researcher, Groningen, The Netherlands

March 5, 2026

Abstract

Definitions: $\varphi \equiv (1+\sqrt{5})/2$ and $\alpha \equiv e^2/(4\pi\varepsilon_0\hbar c)$.

Abstract

We formalize a thermodynamic reinterpretation of the particle mass spectrum within Swirl String Theory (SST). Building upon the quantum-thermodynamic isomorphism, we redefine the Golden-layer mass hierarchy (φ^n) not as a particle-defining topological property, but as a discrete thermodynamic state variable representing SST Heat (δQ_{SST}). The baseline structural identity of a particle is governed by its topological knot invariant (evaluated as adiabatically stored work), while its mass generation is governed by the excitation of Kelvin-modes and R/T-phase reconfigurations. By incorporating the topological Kelvin excitation gap (Δ_K) and the zero-free-parameter derivation of the vacuum equation of state, we demonstrate why low-energy states are highly stable. Applying this to ab initio computational braid data reveals that disparate Standard Model particles (e.g., the Electron and the Higgs Boson) share identical baseline topologies but exist in drastically different, quantized thermodynamic heat states within the swirl condensate.

Keywords: PAPER$_K EYWORDS$

1 The Thermodynamic Equation of State

In the SST Multi-Scale Thermodynamics framework, the physical vacuum is modeled as a frictionless, incompressible swirl condensate. Drawing upon the zero-free-parameter derivation of quantization [2], the total energy E (rest mass) of a vortex filament decomposes into mechanical work and heat transfer, governed by the Abe-Okuyama quantum-thermodynamic isomorphism:

$$dE = \delta W_{SST} + \delta Q_{SST}. \tag{1}$$

1.1 SST Work (Topological Baseline)

The mechanical work $\delta W_{SST} = \sum_k p_k \, dE_k$ corresponds to adiabatic geometric deformation of the core radius r_c against the surrounding vacuum swirl pressure. The baseline mass of an unexcited knot is entirely adiabatic work, constrained by the core momentum-flux invariant

$$A_C^{(SST)} = 4\pi\rho_{\text{core}}\|\mathbf{v}_{\circlearrowleft}\|^2 r_c^4. \tag{2}$$

From the ab initio Braid Search Engine, the baseline geometric mass kernel for a topology K is:

$$M_{\text{base}}(K) = \rho_0 \, \chi(K) \, IM(K) \, L_{\text{tot}}(K), \tag{3}$$

where $\rho_0 \approx 0.022706\,\text{MeV}/L_{\text{tot}}$ is an energy-per-L_{tot} calibration constant fixed once by calibrating the trefoil ground state (3_1), and L_{tot} is the dimensionless ropelength returned by the relaxation engine. The factor $\chi(K) \in \{1, 2\}$ encodes the boson/fermion parity (even components vs. odd components). The invariants (b, g, n) denote bridge/strand density, genus (screening layers), and component number, respectively and

$$IM(K) \equiv b(K)^{-3/2} \, \varphi^{-g(K)} \, n(K)^{-1/\varphi} \tag{4}$$

is the canonical SST topological screening kernel used by the computational engine. To satisfy the

* Email: info@omariskandarani.com
ORCID: 0009-0006-1686-3961

DOI: 10.5281/zenodo.xxx

Crucially, the excitation of these Kelvin modes is not continuous. Reconnection constraints and torsion impose a severe **Topological Excitation Gap** $\Delta_K \sim \mathcal{O}(10^2 - 10^3)\,\mathrm{eV}$ [3]. Below this gap, Kelvin excitations are exponentially suppressed,

$$p_{\mathrm{exc}} \propto e^{-\Delta_K/(k_B T)}, \qquad C_V \propto e^{-\Delta_K/(k_B T)}, \tag{6}$$

guaranteeing extreme stability of fundamental ground states (e.g. the electron) in atomic regimes where thermal coupling is effectively frozen.

1.3 Isotropic Stress Normalization ($V_{eff} = 3V_{core}$)

The derivation of the core momentum-flux invariant $A_C^{(SST)}$ relies strictly on the effective excluded volume of the vortex core. In continuum mechanics for an inviscid medium, the Cauchy stress tensor is given by $\sigma_{ij} = -p\delta_{ij}$. The scalar pressure is therefore related to the trace of the tensor by $p = -\frac{1}{3}\mathrm{tr}(\sigma) = -\frac{1}{3}\sigma_{ii}$.

When a compact spherical core of geometric volume $V_{core} = \frac{4\pi}{3} r_c^3$ is inserted into the background stress field, we adopt an effective isotropic insertion channel model. The scalar work is expressed as $\Delta E \equiv -\delta p V_{eff}$, which absorbs the conversion between the scalar pressure δp and the trace of the normal stresses $\delta\sigma_{ii}$. Because $\delta\sigma_{ii} = -3\delta p$, this normalization convention effectively counts three equal normal-stress contributions (one for each spatial axis).

Thus, the effective thermodynamic volume is exactly three times the geometric volume:

$$V_{eff} = 3V_{core} = 3\left(\frac{4\pi}{3} r_c^3\right) = 4\pi r_c^3. \tag{7}$$

Multiplying this effective volume by the canonical core stress scale $p_* \equiv \rho_{\mathrm{core}} \|\mathbf{v}_{\circlearrowleft}\|^2$ yields the exact invariant $A_C^{(SST)} = 4\pi\rho_{\mathrm{core}}\|\mathbf{v}_{\circlearrowleft}\|^2 r_c^4$. This mechanical derivation eliminates any free parameter in the coupling magnitude [8].

Analogy for a Smart Teenager: Imagine blowing up a balloon underwater. You have to push the water away in all 3 directions of space (up/down, left/right, forward/back). The total thermodynamic effort takes exactly 3 times the work of pushing the water in just a single dimension.

2 Reinterpreting the Braid Sweep Generations

Heating a knot does not change its topology K; rather, it pumps energy across the Kelvin gap into discrete higher-order resonances. The computational parameter $G \in \{0, 1, 2\}$ identified in the SSTcore Braid Sweep is thus formally recognized as the **Thermodynamic Heat Index**.

The total thermodynamic mass functional of a particle is the product of its adiabatic topological kernel and its isothermal heat multiplier:

$$m(K, G) = M_{\mathrm{base}}(K)\,\Lambda_Q(G), \qquad \Lambda_Q(G) \equiv \left(\frac{4}{\alpha}\right)^G. \tag{8}$$

Using discrete scale invariance, the heat multiplier admits a compact golden-ladder approximation

$$\left(\frac{4}{\alpha}\right)^G \approx \varphi^{\nu G}, \qquad \nu \equiv \log_\varphi\left(\frac{4}{\alpha}\right) \approx 13.1 \simeq 13, \tag{9}$$

so that one may write $\Lambda_Q(G) \approx \varphi^{13G}$ as a canonical scaling approximation (not a separate particle-defining topology).

2.1 Kinematic and Geometric Derivation of the Heat Multiplier ($4/\alpha$)

The exact scalar base of the heat multiplier, $4/\alpha$, is not an arbitrary phenomenological fit; it derives directly from the fundamental kinematic and geometric invariants of the swirl string.

From the SST parameter primitives, the fine-structure constant is intrinsically linked to the characteristic swirl speed (SST Identity Group 9):

$$\|\mathbf{v}_{\circlearrowleft}\| = c\left(\frac{\alpha}{2}\right) \quad \Longrightarrow \quad \alpha = \frac{2\|\mathbf{v}_{\circlearrowleft}\|}{c}. \tag{10}$$

Substituting this into the base of the heat multiplier gives a pure velocity ratio:

$$\frac{4}{\alpha} = \frac{4}{2\|\mathbf{v}_{\circlearrowleft}\|/c} = 2\left(\frac{c}{\|\mathbf{v}_{\circlearrowleft}\|}\right). \tag{11}$$

This reveals that the generation scalar represents exactly twice the Mach ratio of the longitudinal wave speed (c) to the transverse core swirl speed ($\|\mathbf{v}_{\circlearrowleft}\|$).

Furthermore, this velocity ratio is strictly bound to the spatial geometry of the vortex loop. From the recomputed primitives, the Compton wavelength of the fundamental reference state (the electron) is defined by the core's rotational parameters (SST Identity Group 8):

$$\lambda_c = \frac{2\pi c r_c}{\|\mathbf{v}_{\circlearrowleft}\|}. \tag{12}$$

Rearranging this geometric relation isolates the velocity ratio:

$$\frac{c}{\|\mathbf{v}_{\circlearrowleft}\|} = \frac{\lambda_c}{2\pi r_c}. \tag{13}$$

Substituting this back into the heat multiplier base yields its definitive geometric form:

$$\frac{4}{\alpha} = 2\left(\frac{\lambda_c}{2\pi r_c}\right) = \frac{\lambda_c}{\pi r_c}. \tag{14}$$

This constitutes a rigorous closure identity within the SST primitive network: the thermodynamic generation jump ($\Lambda_Q(G)$) is precisely governed by the scale-invariant geometric ratio between the longitudinal interaction wavelength of the string (λ_c) and the transverse half-circumference of its core (πr_c). Heating the knot by one generation ($G \to G+1$) corresponds to amplifying its available thermodynamic phase space by this exact scale-invariant spatial ratio.

2.2 Thermodynamic Equivalences (Isotopes of Heat)

Particles that share the same knot topology K but differ in generation G are interpreted as the same geometric entity, existing at different quantized thermodynamic swirl temperatures (T_{swirl}).

Trefoil family (3_1). For the trefoil topology we use the canonical invariants

$$b = 2, \quad g = 1, \quad n = 1, \quad \chi = 2, \quad L_{\text{tot}} = 46.450,$$

which yields (Braid Engine output)

$$m(3_1, G=0) = 0.4609 \text{ MeV} \quad \text{(electron-scale)}, \qquad m(3_1, G=2) = 138.488 \text{ GeV} \quad \text{(Higgs-scale)}.$$

SST identification: the Higgs-scale state is modeled as a high-heat excitation of the same baseline trefoil topology (not an orthodox Standard Model identity).

Cinque-foil torus family (5_1). For 5_1 we use

$$b = 2, \quad g = 2, \quad n = 1, \quad \chi = 2, \quad L_{\text{tot}} = 57.410,$$

giving

$$m(5_1, G = 1) = 192.989 \text{ MeV} \quad (\pi^0\text{-scale}), \qquad m(5_1, G = 2) = 105.786 \text{ GeV} \quad (Z\text{-scale}).$$

Unknot family (1_1). A one-component unknot ring with fermionic parity choice $\chi = 2$ and invariants

$$b = 1, \quad g = 0, \quad n = 1, \quad \chi = 2, \quad L_{\text{tot}} = 25.932,$$

gives

$$m(1_1, G = 1) = 645.508 \text{ MeV} \quad (\text{strange-quark constituent scale}).$$

3 The Geometric Limit of the SST Mass Functional

3.1 Definitions (to avoid hidden assumptions)

We use the electron Compton wavelength

$$\lambda_c := \frac{h}{M_e c}, \tag{15}$$

and the SST geometric identity (Group 8 closure)

$$\lambda_c = \frac{2\pi c\, r_c}{\|\mathbf{v}_{\circlearrowleft}\|}. \tag{16}$$

We define a *core energy density* ρ_E ($\mathrm{J\,m^{-3}}$) and its mass-equivalent

$$\rho_m := \frac{\rho_E}{c^2} \qquad (\mathrm{kg\,m^{-3}}). \tag{17}$$

In the canonical ground-state normalization used throughout this manuscript, we identify $\rho_m \equiv \rho_{\text{core}}$. This avoids dimensional ambiguity if one uses "energy density" language in the mass functional.

3.2 Elimination of kinematic variables via the Compton–core identity

From (16),

$$\frac{\|\mathbf{v}_{\circlearrowleft}\|}{c} = \frac{2\pi r_c}{\lambda_c}, \qquad \frac{c}{\|\mathbf{v}_{\circlearrowleft}\|} = \frac{\lambda_c}{2\pi r_c}. \tag{18}$$

3.3 Geometric baseline mass scale

Assume a localized kinetic-energy density of the form

$$u \;=\; \frac{1}{2}\rho_m \,\|\mathbf{v}_{\circlearrowleft}\|^2 \;=\; \frac{1}{2}\frac{\rho_E}{c^2}\,\|\mathbf{v}_{\circlearrowleft}\|^2, \tag{19}$$

and a geometric core volume

$$V := \pi r_c^3\, L_{\text{tot}}(T), \tag{20}$$

where $L_{\text{tot}}(T)$ is a dimensionless ropelength-like factor (or a dimensionless length ratio in your discretization).

Then the inertial mass scale defined by energy equivalence is

$$M_0(T) \;:=\; \frac{uV}{c^2} = \frac{1}{2}\rho_m \left(\frac{\|\mathbf{v}_{\circlearrowleft}\|}{c}\right)^2 \pi r_c^3\, L_{\text{tot}}(T). \tag{21}$$

Substituting (18) into (21) gives the *pure geometric* form

$$\boxed{M_0(T) = 2\pi^3\,\rho_m\,\frac{r_c^5}{\lambda_c^2}\,L_{\mathrm{tot}}(T).} \tag{22}$$

Dimensional check: ρ_m ($\mathrm{kg\,m^{-3}}$) times r_c^5/λ_c^2 ($\mathrm{m^3}$) yields kg. *No free kinematic variable remains.*

3.4 Substitution of the shielding gate

From the previously established geometric identity for the sector gate,

$$\boxed{\frac{4}{\alpha} = \frac{2c}{\|\mathbf{v}_{\circlearrowleft}\|} = \frac{\lambda_c}{\pi r_c},} \tag{23}$$

define the exposure gate $G(T) \in \{0, 1, 2\}$ and a dimensionless topology kernel $\mathcal{K}(T)$. The canonical mass functional becomes

$$\boxed{M(T) = \left(\frac{\lambda_c}{\pi r_c}\right)^{G(T)} \mathcal{K}(T)\,\left(2\pi^3\,\rho_m\,\frac{r_c^5}{\lambda_c^2}\,L_{\mathrm{tot}}(T)\right).} \tag{24}$$

This form contains no explicit α and no explicit $\|\mathbf{v}_{\circlearrowleft}\|$.

3.5 Geometric reduction of the GR-like time dilation term

Consider the standard Schwarzschild-like factor appearing in GR time dilation,

$$\Phi(T; r) \;:=\; \frac{2GM(T)}{rc^2}. \tag{25}$$

Substituting (24) into (25) yields

$$\boxed{\Phi(T; r) = \frac{4\pi^3\,G\,\rho_m}{c^2}\,\left(\frac{\lambda_c}{\pi r_c}\right)^{G(T)} \mathcal{K}(T)\,\frac{r_c^5}{r\,\lambda_c^2}\,L_{\mathrm{tot}}(T).} \tag{26}$$

Key point. The explicit symbols G and M no longer appear separately: the potential is now controlled by *geometric ratios* multiplied by the composite prefactor $(G\rho_m/c^2)$. Thus the "cancellation" is a *reparameterization* once you posit $M(T)$ geometrically; it is not a GR identity.

3.6 Numerical gate validation (dimensionless)

Using canonical lengths r_c and λ_c,

$$\frac{\lambda_c}{\pi r_c} \approx 5.48144 \times 10^2, \tag{27}$$

matching $4/\alpha$ at rounding precision.

3.7 Status, assumptions, falsifiers

Assumptions: (i) energy density model (19); (ii) geometric core volume (20); (iii) closure identity (16); (iv) binary/integer exposure gate $G(T)$.

Falsifiers: any empirical inconsistency in the identity network (e.g. λ_c vs r_c vs $\|\mathbf{v}_{\circlearrowleft}\|$) breaks (23); any failure of (24) to predict sector ratios when L_{tot} and $\mathcal{K}$ are fixed breaks the physical interpretation of $G(T)$ as "exposure."

4 Gravity as Pure Geometric Impedance in SST

4.1 Geometric reduction of the GR coupling term

The GR (Schwarzschild) proper-time factor may be written in terms of the dimensionless potential

$$\Phi(T;r) \;:=\; \frac{2GM(T)}{rc^2}. \tag{28}$$

Using the standard definitions $L_p^2 = \hbar G/c^3$ and $\lambda_c = 2\pi\hbar/(M_e c)$, the electron gravitational radius

$$r_s(e) \;:=\; \frac{2GM_e}{c^2} \tag{29}$$

reduces identically to the purely geometric form

$$\boxed{r_s(e) = 4\pi \frac{L_p^2}{\lambda_c}.} \tag{30}$$

Thus, at the electron scale, the GR "source strength" GM_e/c^2 is equivalently a ratio of the Planck area to the Compton boundary.

4.2 SST topological mass ratio

Define the dimensionless SST mass ratio relative to the electron,

$$\boxed{\tilde{M}(T) := \frac{M(T)}{M_e} = \left(\frac{\lambda_c}{\pi r_c}\right)^{G(T)} \mathcal{K}(T)\, \tilde{L}_{\text{tot}}(T),} \tag{31}$$

where $G(T)$ is the exposure (shielding) gate, $\mathcal{K}(T)$ is the topology kernel, and $\tilde{L}_{\text{tot}}(T)$ is the relative ropelength factor.

4.3 GR time dilation with SST substitution

For a knot state T, the GR potential term becomes

$$\Phi(T;r) = \frac{r_s(e)}{r}\,\tilde{M}(T) = \boxed{\frac{4\pi L_p^2}{r\lambda_c}\left(\frac{\lambda_c}{\pi r_c}\right)^{G(T)} \mathcal{K}(T)\, \tilde{L}_{\text{tot}}(T).} \tag{32}$$

Substituting this into the Schwarzschild-like proper-time factor and appending SST-specific local swirl terms yields

$$t_{\text{adj}} = \Delta t \sqrt{1 - \Phi(T;r) - \frac{\|\mathbf{v}_{\circlearrowleft}\|^2}{c^2} e^{-r/r_c} - \frac{(\omega r_c)^2}{c^2} e^{-r/r_c}}. \tag{33}$$

Interpretation (precise claim). The GR structure is retained; SST enters only through $\tilde{M}(T)$. In this combined model, the gravitational time-dilation term can be written as a product of (i) a universal length-ratio prefactor $4\pi L_p^2/(r\lambda_c)$ and (ii) a purely dimensionless topological impedance factor $(\lambda_c/(\pi r_c))^{G(T)}\,\mathcal{K}(T)\tilde{L}_{\text{tot}}(T)$. No additional mass parameters are introduced beyond the electron normalization.

5 Hydrodynamic Isothermal Equilibrium of the Bohr Radius

In standard quantum mechanics, the Bohr radius a_0 is derived as a probabilistic expectation value. In the SST Multi-Scale Thermodynamics framework, it emerges strictly as a thermodynamic equilibrium surface.

An electron represented by a swirl string possesses two relevant geometric scales: the internal core radius r_c and the macroscopic orbital envelope a_0. During the formation of a hydrogen atom, the vortex string undergoes a geometric expansion from r_c to a_0.

Applying Euler's radial force balance for a steady circular flow:

$$\frac{dp}{dr} = \rho_f \frac{v_\theta^2}{r} \qquad \text{where} \qquad v_\theta(r) = \frac{\Gamma_0}{2\pi r}. \tag{34}$$

Integrating from the exterior vacuum reservoir inward (p_∞), the internal pressure drops according to:

$$p(r) = p_\infty - \frac{1}{2}\rho_f v_\theta^2(r). \tag{35}$$

As r decreases, the local swirl velocity v_θ increases, lowering the internal pressure at the core boundary and generating a net inward compressive hydrodynamic force.

Thermodynamically, atomic binding corresponds to mechanical work performed on the swirl string. However, the system must remain in equilibrium with the reservoir's background swirl fluctuations. Thus, the binding process is strictly an **isothermal expansion**. The stable equilibrium point occurs where the derivative of the free energy vanishes ($\frac{dF}{dr} = 0$), mapping precisely to the radius $r = a_0$ where the internal centrifugal swirl pressure exactly balances the external vacuum tension:

$$p_{swirl}(a_0) = p_{vac}. \tag{36}$$

Consequently, atomic structure in SST is fundamentally a hydrodynamic pressure-balance, effectively bypassing probability amplitudes.

Analogy for a Smart Teenager: Imagine spinning a heavy weight on a highly elastic rubber band. The faster you spin it, the more the rubber band stretches outward. The Bohr radius (a_0) is the exact, stable distance where the outward pull of the spinning fluid perfectly cancels out the inward snap of the vacuum's rubber band.

6 The Unruh Echo: Two-Stage Thermodynamic Transduction

The thermodynamic framework naturally extends to highly accelerated reference frames. Standard Unruh radiation arises from coupling to the electromagnetic vacuum at the speed of light c [9]. SST introduces a dual-vacuum ontology: the swirl condensate governed by the characteristic velocity $\mathbf{v}_{\circlearrowleft}$, and the electromagnetic transduction sector governed by c.

Accelerated motion of a vortex filament mechanically stretches local vorticity, directly raising the local swirl temperature:

$$T_{swirl}(a) \propto a. \tag{37}$$

This mechanical stretching forces an excitation of internal Kelvin modes (SST Heat). However, due to the severe impedance mismatch between the two vacuum sectors ($\|\mathbf{v}_{\circlearrowleft}\| \ll c$), the resulting energy radiation cannot occur instantaneously. Instead, we hypothesize that it manifests as a Two-Stage Thermodynamic Pulse with characteristic scaling delays:

1. **Primary Burst ($\sim$ 0.1 ns scale):** The initial acceleration causes a rapid thermal excitation within the pure swirl sector, registering as an immediate fluctuation of local vorticity (δQ_{sw}).

2. **Secondary Echo ($\sim$ 30 ns scale):** The excited Kelvin modes interact with boundaries or charges in the condensate. A portion of this thermal energy is re-radiated as electromagnetic work (δW_{EM}), creating a distinctly delayed transduction pulse.

The Unruh effect in Swirl String Theory is therefore reconceptualized not as abstract particle creation from an event horizon, but as a macroscopic heat-work transduction problem, where:

$$\delta Q_{sw}(t = 0.1 \text{ ns}) \longrightarrow \delta W_{EM}(t = 30 \text{ ns}). \tag{38}$$

Analogy for a Smart Teenager: Imagine throwing a heavy rock into a muddy pond. First, you see the immediate, violent splash of the mud (the 0.1 ns swirl burst). But several seconds later, the ripples finally travel across the pond and hit a glass bottle floating at the edge, making a "clink" sound (the 30 ns electromagnetic echo). It is the same event, but the energy takes time to switch from mud-waves into sound-waves.

7 SST Effective Ansatz for Suppression of an Inverse-Square Interaction under Tangential Swirl Excitation

7.1 Scope and Status

This section introduces a *phenomenological* SST ansatz for suppression of an effective inverse-square interaction amplitude under tangential swirl excitation. It is not derived from the SST field equations. The purpose is a dimensionally consistent model suitable for calibration and falsification.

7.2 Baseline Inverse-Square Interaction (Effective Coupling Form)

Let Γ_1 and Γ_2 denote circulation-like SST state variables (units $\mathrm{m}^2\,\mathrm{s}^{-1}$), and let r be the separation (m). We postulate an effective inverse-square baseline term

$$F_0(r) = \mathcal{K}_\Gamma \frac{\Gamma_1 \Gamma_2}{r^2}, \tag{39}$$

with

$$[\mathcal{K}_\Gamma] = \mathrm{N}\,\frac{\mathrm{s}^2}{\mathrm{m}^2} = \mathrm{kg}. \tag{40}$$

7.3 Bounded SST Suppression Factor (Clock-Aligned Form)

Define a bounded suppression factor

$$S(v_t; v_\star) \equiv 1 - \frac{v_t^2}{v_\star^2}, \qquad 0 \le v_t \le v_\star, \tag{41}$$

so that $0 \le S \le 1$. For the special choice $v_\star = c$, one has $S = 1 - v_t^2/c^2 = S_t^2$ where S_t is the SST speed-based time-slow factor.

The effective interaction is then

$$F_{\text{eff}}(r, v_t) = F_0(r)\, S(v_t; v_\star) = \mathcal{K}_\Gamma \frac{\Gamma_1 \Gamma_2}{r^2} \left(1 - \frac{v_t^2}{v_\star^2}\right). \tag{42}$$

7.4 Resonance-Driven Response (Bounded Power-to-Speed Map)

Let $\mathcal{P}_{\text{res}}$ be the resonance drive power and Q the resonance quality factor. Introduce a dimensionless response function η:

$$v_t(\mathcal{P}_{\text{res}}) = v_\star\, \eta(\mathcal{P}_{\text{res}}), \qquad 0 \le \eta < 1. \tag{43}$$

A convenient bounded choice is

$$\eta(\mathcal{P}_{\text{res}}) = \tanh\!\big(\alpha Q \mathcal{P}_{\text{res}}\big), \qquad [\alpha] = \mathrm{W}^{-1}, \tag{44}$$

so that $\alpha Q \mathcal{P}_{\text{res}}$ is dimensionless.

Thus

$$F_{\text{eff}}(r, \mathcal{P}_{\text{res}}) = \mathcal{K}_\Gamma \frac{\Gamma_1 \Gamma_2}{r^2} \left[1 - \eta(\mathcal{P}_{\text{res}})^2\right]. \tag{45}$$

7.5 Load-Compression Dependence

If loading modifies the separation, write $r = r(P_{\text{load}}) > 0$. Then

$$\mathcal{F}_{\text{int}}(P_{\text{load}}, \mathcal{P}_{\text{res}}) = \mathcal{K}_\Gamma \frac{\Gamma_1\Gamma_2}{r(P_{\text{load}})^2} \left[1 - \eta(\mathcal{P}_{\text{res}})^2\right]. \tag{46}$$

7.6 Small-Signal Limit (Fit Form)

For $\alpha Q \mathcal{P}_{\text{res}} \ll 1$, $\tanh x \approx x$ gives

$$\mathcal{F}_{\text{int}} \approx \mathcal{K}_\Gamma \frac{\Gamma_1\Gamma_2}{r(P_{\text{load}})^2} \left[1 - (\alpha Q \mathcal{P}_{\text{res}})^2\right]. \tag{47}$$

7.7 SST Falsifiable Split: Limiting Speed Choice

Two discriminating hypotheses:

$$v_\star = c \qquad \text{(clock-aligned relativistic limiter: } S = S_t^2\text{)}, \tag{48}$$

$$v_\star = \|\mathbf{v}_{\circlearrowleft}\| \qquad \text{(SST characteristic swirl-speed limiter)}. \tag{49}$$

7.8 Numerical SST Sanity Check (Swirl-Speed Limiter)

Using the SST Canon reference constant

$$\|\mathbf{v}_{\circlearrowleft}\| = 1.09384563 \times 10^6 \ \mathrm{m\,s^{-1}}, \tag{50}$$

if $v_t = 0.9\,\|\mathbf{v}_{\circlearrowleft}\|$, then $S = 1 - 0.9^2 = 0.19$ (i.e. 81% suppression relative to F_0).

8 Thermodynamic Consistency and the Heat Capacity

Because the Golden Layers dictate thermodynamic states rather than distinct particles, the heat capacity of the vacuum must reflect both the Δ_K suppression at low energies and the log-periodic structure at high energies.

In the frozen regime ($k_B T \ll \Delta_K$), the heat capacity is exponentially suppressed, ensuring that $C_V \to 0$ without destabilizing atomic orbitals [3]. At extreme collision energies where Δ_K is overcome, the discrete Golden Ladder (truncated at N modes) activates:

$$C_V = k_B \beta^2 \left[\frac{1}{Z} \sum_{n=0}^{N} E_n^2 e^{-\beta E_n} - \left(\frac{1}{Z} \sum_{n=0}^{N} E_n e^{-\beta E_n} \right)^2 \right], \tag{51}$$

where $E_n = E_0 \varphi^n$.

The discrete steps of $\sim 13n$ layers (i.e. φ^{13}) identified by the Braid Engine align with the expected logarithmic spacing of activated Kelvin thresholds. As an external interaction (such as a high-energy particle collision) raises the local T_{swirl}, the collision energy is absorbed as δQ_{SST}, pumping the vortex string up the Golden Ladder ($G = 0 \to 1 \to 2$) until it reaches the fundamental breakdown limit of the continuum.

References

[1] Iskandarani, O. (2026). "Multi-Scale Thermodynamics of the Swirl Condensate: A Unified Hydrodynamic-Topological Framework for Swirl-String Theory". *Zenodo*, DOI: 10.5281/zenodo.17813283.

[2] Iskandarani, O. (2026). "Thermodynamic Origin of Quantization in Swirl-String Theory: From Clausius Work-Heat Structure to Parameter-Free Constants". *Independent Researcher, Groningen.*

[3] Iskandarani, O. (2026). "Kelvin Mode Suppression in Atomic Orbitals: A Vortex-Filament Gap". *Independent Researcher, Groningen.*

[4] Iskandarani, O. (2026). "Atomic Masses from Topological Invariants of Knotted Field Configurations", *Independent Research (manuscript).*

[5] Iskandarani, O. (2026). "Velocity-Dependent Mass Functional, Preferred-Frame Effects, and Intrinsic Temporal Stochasticity", *Independent Research.*

[6] Abe, S., & Okuyama, S. (2011). "Similarity between quantum mechanics and thermodynamics: Entropy, temperature, and Carnot cycle". *Physical Review E*, 83, 021121. DOI: 10.1103/PhysRevE.83.021121.

[7] Sornette, D. (1998). "Discrete scale invariance and complex dimensions". *Physics Reports*, 297, 239-270. DOI: 10.1016/S0370-1573(97)00076-8.

[8] Landau, L. D., & Lifshitz, E. M. (1987). *Fluid Mechanics*, 2nd ed. (Course of Theoretical Physics, Vol. 6). Pergamon Press. ISBN: 978-0750627672.

[9] Unruh, W. G. (1976). "Notes on black-hole evaporation". *Physical Review D*, 14, 870-892. DOI: 10.1103/PhysRevD.14.870.

[10] Tiesinga, E., Mohr, P. J., Newell, D. B., & Taylor, B. N. (2023). "The 2022 CODATA Recommended Values of the Fundamental Physical Constants". *Rev. Mod. Phys.* **95**, 025003. DOI: 10.1103/RevModPhys.95.025003.

[11] Misner, C. W., Thorne, K. S., & Wheeler, J. A. (1973). *Gravitation.* W. H. Freeman. ISBN: 978-0716703440.

[12] Abbott, B. P. et al. (LIGO Scientific Collaboration and Virgo Collaboration) (2017). "GW170817: Observation of Gravitational Waves from a Binary Neutron Star Inspiral". *Phys. Rev. Lett.* **119**, 161101. DOI: 10.1103/PhysRevLett.119.161101.