

Fusion draft: SST-32/44 Canonical fluid reformulation capstone

Omar Iskandarani (fusion draft)

April 19, 2026

About this fusion draft

This is an automatically assembled fusion draft: multiple SST manuscripts concatenated as chapters. It is intended for navigation and editing workflow, not as a final merged publication without manual cleanup.

Contents

1	SST-32 — Canonical fluid reformulation (long)	5
1.1	Introduction	1
1.2	Core Postulates of Swirl-String Theory	2
1.3	Lagrangian and Field-Theoretic Framework	4
1.3.1	Foundational Fields and Definitions	4
1.3.2	Effective Lagrangian Density	6
1.4	Emergent Gravity and Time Dilation	9
1.4.1	Newtonian Gravity as a Fluid-Flow Effect	9
1.4.2	Swirl-Clock Time Dilation (Chronon Contraction)	10
1.5	Electromagnetic Emergence: Modified Faraday Law	12
1.5.1	Swirl Areal Density and Electromotive Force	12
1.5.2	Conceptual and Experimental Considerations	13
1.6	Chirality and Quantum Measurement Dynamics	14
1.6.1	Swirl Clock Orientation and Time Asymmetry	15
1.6.2	Wavefunction Collapse as Vortex Topology Change	15
1.6.3	Parity and Time-Reversal in SST	16
1.7	Canonical Quantization and Topological Spectrum	17
1.7.1	Circulation Quantization and Combinatorial Spectrum	17
1.8	Experimental Implications & Falsifiability	20
1.9	Comparison with Existing Frameworks	25
1.10	Conclusion & Outlook	30
	Appendix A: Lagrangian Density and Master Equations	33
	Appendix B: Derivation of Swirl-Clock Time Dilation	35
	Appendix C: Emergent Newtonian Gravity from Swirl Pressure	37
	Appendix D: Electromotive Impulse from Swirl Density Change	39
A	SST-44 — Parallel capstone	43
A.1	Introduction	1
A.2	Core Postulates of Swirl-String Theory	2
A.3	Lagrangian and Field-Theoretic Framework	2
A.4	Internal Gauge Symmetry in SST: $SU(3) \times SU(2) \times U(1)$	5
A.5	Swirl-Braid Correspondence and Energy Functional	6
A.6	Local Swirl Stress and Lab Traction	8
A.7	Emergent Gravity and Time Dilation	8
A.8	Cosmological Branch: de Sitter Background in Swirl-String Theory	9
A.8.1	From Local Swirl Wells to Global Expansion	9
A.8.2	Canonical Identification with the Cosmological Constant	9
A.8.3	Numerical Evaluation	10
A.8.4	Unified Interpretation with the Hydrogen-Gravity Mechanism	10
A.8.5	Physical Picture and Testable Consequences	10
A.9	Electromagnetic Emergence: Modified Faraday Law	11

A.10 Spectral Thermodynamics of Swirl Excitations	12
A.11 Chirality and Quantum Measurement Dynamics ($R \leftrightarrow T$)	13
A.12 Canonical Quantization and Topological Spectrum	13
A.13 Real-Time Tunneling and the Kappa Event	14
A.14 Experimental Implications & Falsifiability	15
A.14.1 Identification from Slow and Non-Uniform Samples	15
A.15 Experimental Testability and Falsifiable Setups	16
A.15.1 Swirl-Clock Dilation under Chiral Driving	16
A.15.2 $R \leftrightarrow T$ Transition Map in Knotted Coils	16
A.15.3 Swirl-Pressure Force Proxy	16
A.15.4 Swirl-Clock Time Dilation under Chiral Driving	16
A.15.5 Quantized EM Impulse from Swirl Reconnection	16
A.15.6 Swirl-Pressure Gravity Proxy	17
A.15.7 GW Sector Sanity Check (Propagation Speed)	17
A.16 Running Couplings and Topological β -Functions	17
A.17 Canonical Quantization of $\mathbf{v}_G$ and Directors	17
A.18 Mass Generation: Scalar Condensate vs. Topological Tension	17
A.18.1 Scalar Swirl Condensate (Higgs-analogue)	17
A.18.2 Topological Tension Mechanism	18
A.19 Charge and Quantum Numbers from Knot Data: Benchmark Table	18
A.20 Gravitational Waves and Lorentz-Breaking Bounds	18
A.21 Cosmological Embedding: Dark Matter and Dark Energy	18
A.21.1 Swirl-Foam Dark Matter	18
A.21.2 Network-Tension Dark Energy	19
A.22 Cosmology: Baryogenesis and Dark Sector in SST	19
A.22.1 Chiral Swirl Condensation and Baryogenesis	19
A.22.2 Dark Matter as Swirl Foam	19
A.22.3 Dark Energy as Network Tension	19
A.23 Meta-Topology and Cosmogenetic Hierarchy in SST	19
A.23.1 Swirl Genesis and Knot Nucleation	19
A.23.2 Golden-Ratio Mass Ladder	19
A.23.3 Hierarchy from Contact and Helicity Correction	20
A.23.4 Interpretation and Cosmological Implications	20
A.24 Numerical Program: Swirl-Knot Dynamics and EM Coupling	20
A.25 Comparative Positioning	20
A.26 Comparison with Existing Frameworks	20
A.27 Conclusion & Outlook	21
.1 Full Canonical Lagrangian for SST	27
.2 Electroweak mixing and charge operator	28
.3 Anomaly constraints and hypercharge calibration	28
.4 Representative knot $\rightarrow$ rep assignments	28
.5 Swirl Cushion Stability and Leidenfrost Analogy	28
.5.1 Empirical Leidenfrost Scaling	28
.5.2 SST Translation: Pressure-Film Criterion	29
.5.3 Numerical Consistency	29
.5.4 Interpretation and Analogy	29
.6 Identification with Slow and Non-Uniform Sampling (LFT-DAE Form)	29
Identification with Slow and Non-Uniform Sampling (LFT-DAE Form)	29
.7 Kappa-Event Dwell-Time Experiment (Larmor-Clock Protocol)	30
.8 Variational Derivation of the Clock-Rescaled Schrödinger Equation	31
.9 Spectral Thermodynamics: Density of States and Core Regularization	31

.10	Photoelectric Mapping in SST Units: Geometry and Units	32
.11	Derivation of the Local Swirl Stress Tensor	33
.12	Benchmark Knot-Representation Table	33
.13	Swirl-Electromagnetic Unit Bridge Table	34

Chapter 1

SST-32 — Canonical fluid reformulation (long)

Swirl–String Theory: Canonical Fluid Reformulation of Relativity and Quantum Structure Omar Iskandarani

Independent Researcher, Groningen, The Netherlands

ORCID: 0009-0006-1686-3961, DOI: 10.5281/zenodo.xxxx April 19, 2026

Abstract

We present *Swirl-String Theory* (SST), a fluid-topological framework that reinterprets relativity and quantum phenomena via a single incompressible medium. In SST, matter and radiation are modeled as quantized vortex loops (*swirl strings*) in a universal, non-dissipative condensate. The theory posits that classical gravity emerges as a collective pressure effect of these vortices rather than fundamental spacetime curvature, and that local *swirl flows* induce time dilation analogous to relativistic kinematics. We develop the formal field-theoretic Lagrangian for this medium, showing how a preferred foliation and topological quantization yield a discrete particle spectrum: quantum numbers (mass, charge, spin) correspond to topological invariants of knot-like vortex excitations. A modified Faraday’s law is derived, unifying electromagnetic induction with rotating-frame effects by predicting that time-varying swirl string density generates an electromotive force. We also describe how SST accounts for wave–particle duality through dual phase (un-knotted vs. knotted) states of vortex loops, with quantum measurement corresponding to topological transitions. Several experimental tests are proposed to falsify or support SST, including quantized electromagnetic impulses from vortex reconnection events, attosecond-scale chirality-dependent time delays in photoemission, and interference degradation due to finite vacuum vorticity. We compare SST with established frameworks—from Kelvin’s vortex atom hypothesis to emergent gravity and analogue fluid models—highlighting how SST recovers known limits (Newtonian gravity, Maxwell electrodynamics, quantum wave behavior) while providing novel, quantifiable predictions. All equations are given in SI units with attention to dimensional consistency. The result is a self-contained canonical reformulation that bridges classical and quantum physics through fluid-like continuum mechanics.

Keywords: vortex dynamics; topological fluid; quantum topology; emergent gauge theory; time dilation; wavefunction collapse

1.1 Introduction

The modern theories of physics—quantum field theory and general relativity—offer tremendously successful descriptions of phenomena, yet they rest on disparate conceptual foundations. In one view, spacetime is a dynamical geometrical manifold, while quantum matter is described by abstract wavefunctions and fields. *Swirl-String Theory* (SST) is an alternative paradigm that seeks a unifying, realist substrate: it posits that all particles and forces emerge from excitations of a universal incompressible fluid medium. In spirit, this approach revives a form of the historical *ether*, but crucially it leverages modern insights to remain consistent with relativity and quantum observations. The notion of matter as vortices in a fluid can be traced back to Lord Kelvin’s vortex atom hypothesis of 1867 [?], wherein atoms were imagined as stable knotted smoke rings in a primordial fluid. SST extends this idea with quantization and topology, aligning it with the discoveries of quantized vortex circulation in superfluids [9, ?] and with contemporary emergent gravity scenarios [18, 7, ?].

In SST, the vacuum is modeled as a three-dimensional Euclidean space R^3 filled with an ideal incompressible fluid (the *swirl medium*) and an absolute time parameter provides a preferred foliation. Stable, quantized vortex filaments in this medium (*swirl strings*) serve as the analogues of elementary particles and field quanta. Crucially, each such filament can only exist in certain topologically distinct configurations (e.g. different knot types), and the circulation around each closed loop is quantized in units of a fundamental circulation quantum κ . These two features endow SST with a **topological spectrum**: physical attributes like rest mass, charge, and intrinsic spin are tied to topological invariants of the vortex (such as knot linking number or writhing number) rather than arbitrary continuous parameters. By calibrating a single characteristic velocity and length scale of the medium, SST’s mass scale is fixed and yields values for particle masses without *ad hoc* Yukawa couplings. In this way, SST attempts to derive quantitative particle properties from first principles, a strength not shared by the Standard Model.

Operationally, SST can be viewed as a modern Lorentzian ether theory (LET). It contains a preferred rest frame (the fluid rest frame), but it reproduces the relativistic time dilation and length contraction relationships in all physical predictions, making it empirically indistinguishable from Special Relativity at the kinematic level. Unlike the 19th-century ether models dismissed by Occam’s razor, the SST medium is not superfluous or undetectable; it yields concrete testable effects. For instance, as we describe in this paper, SST predicts new phenomena such as a coupling between vortex dynamics and electromagnetic fields that could be measured as quantized voltage impulses. These distinctive predictions offer potential falsifiability. At the same time, the theory is constructed to recover the established limits: Newtonian gravity emerges in the slow-flow regime, Coulomb’s law and Maxwell’s equations emerge in the linear perturbation regime, and quantum wave behavior emerges when a swirl string is extended and unknotted. In short, SST aims to *reformulate* the foundations of physics as the dynamics of a single classical continuum, from which both relativity and quantum structure are seen as emergent, unified phenomena.

In the following sections, we outline the core postulates of SST (Sec. 2) and develop the field-theoretic Lagrangian framework that embodies these principles (Sec. 3). We then explain how familiar effects—gravity and time dilation (Sec. 4), and electromagnetism (Sec. 5)—arise naturally from the fluid dynamics of the swirl medium. The dual nature of quantum particles (wave vs. particle) and the role of chirality in measurement are discussed in Sec. 6, wherein we connect the handedness of knotted vortices to asymmetric time evolution (a possible explanation for observed chiral photoemission delays). Section 7 describes how canonical quantization in SST is supplanted by topological quantization, yielding a discrete spectrum of stable excitations that map onto known particles. Next, Sec. 8 delineates experimental implications and falsifiable predictions of the theory, including proposed tabletop experiments. In Sec. 9 we compare

and contrast SST with established frameworks such as the Standard Model, general relativity, analogue gravity systems, and other emergent or topological matter theories. Finally, Sec. 10 summarizes our conclusions and outlines directions for further research. Key mathematical derivations supporting the main text are provided in Appendices A–D, including the derivation of the swirl-clock time dilation factor, the emergent Newtonian gravitational potential from swirl flows, and the modified Faraday induction law with quantized flux impulses.

Throughout, we use SI units and ensure dimensional analysis is consistent. Important physical quantities are defined explicitly when introduced. For reference, the swirl medium’s baseline mass density is denoted ρ_f , the local swirl kinetic energy density by ρ_E , and the equivalent mass density $\rho_m = \rho_E/c^2$. The swirl velocity field is represented by $\mathbf{v}_\zeta$, and a characteristic vortex core radius is r_c . We also introduce a fundamental circulation quantum κ and a characteristic core swirl speed v_o that together set the scale of quantum circulation. Table ?? in Appendix A summarizes these and other constants of the SST framework.

1.2 Core Postulates of Swirl–String Theory

SST is founded on a set of core postulates (*axioms*) which define its ontology and basic laws:

1. **Swirl Medium (Absolute Space and Time):** There exists a ubiquitous, incompressible fluid-like condensate filling three-dimensional space. Space is Euclidean $\mathbb{R}^3$ with a global absolute time t that parameterizes a preferred foliation of spacetime. All physical processes occur within this *swirl medium*, which is assumed to have zero viscosity and infinite conductivity for vortex motion (analogous to a perfect classical fluid). This postulate establishes an objective rest frame (the medium rest frame or “lab frame”), although no experiment confined to a small region can detect uniform motion relative to this medium (ensuring consistency with relativity).
2. **Swirl Strings (Quantized Circulation and Topology):** The elementary excitations of the medium are *swirl strings*, defined as closed vortex filaments (loops of circulating flow) within the fluid. Each swirl string may form nontrivial knots or links. The circulation Γ of the swirl velocity field $\mathbf{v}_\zeta$ around any closed loop C in the medium is quantized in integer multiples of a fundamental quantum κ :

$$\Gamma = \oint_C \mathbf{v}_\zeta \cdot d\mathbf{l} = n \kappa, \quad n \in \mathbb{Z}. \quad (1.1)$$

We identify κ with h/m_{eff} (where h is Planck’s constant and m_{eff} is a characteristic mass scale of the condensate) to connect with quantum units of circulation [9, ?]. In addition to quantized circulation, swirl strings are subject to topological quantization: only certain knot configurations are permissible. Thus, discrete quantum numbers such as particle species, electric charge, and spin correspond to topological invariants of the vortex filament (e.g. linking number, knot twist, writhe) rather than being intrinsic labels in a quantum state space. In essence, *matter is “knotted vacuum”*: each stable particle is a stable knot of vorticity.

3. **String-Induced Gravitation (Emergent Gravity):** Macroscopic gravitational attraction is not a fundamental geometric warping of spacetime in SST, but rather an emergent, statistically averaged force arising from the collective dynamics of the swirl medium. Coherent flows and pressure gradients induced by many swirl strings produce an effective long-range attraction between concentrations of energy. In the non-relativistic, weak-flow limit, this emergent force obeys an inverse-square law. The effective gravitational coupling in SST, denoted G_{swirl} , can be set equal to Newton’s constant G_N by appropriate choice of the medium’s parameters. Thus, SST reproduces Newtonian gravity (and by

extension, satisfies the Equivalence Principle in practice) even though conceptually gravity here is a pressure-mediated phenomenon in flat space [18, 7]. A concrete mechanism (the “Hydrogen-Gravity” mechanism) illustrating how two knotted vortices attract via induced low-pressure regions in the fluid is described in Sec. 4.

4. **Swirl Clocks (Local Time Dilation):** The passage of time in SST is modulated by local motion of the medium. A clock comoving with a swirl string (and thus moving through the medium with some tangential speed v) will tick slower relative to a clock at rest in the medium. Specifically, if dt_∞ is an interval of the universal time (as measured by an observer at rest in the fluid far from any swirling motion) and dt_{local} is the proper time interval measured by a clock moving with the vortex, SST posits

$$\frac{dt_{\text{local}}}{dt_\infty} = S_t = \sqrt{1 - \frac{v^2}{c^2}}, \quad (1.2)$$

where c is the characteristic wave propagation speed in the medium (identified with the vacuum speed of light). This factor S_t is analogous to the Lorentz time dilation factor in special relativity. In SST it arises because kinetic energy stored in local swirl motion effectively slows the progression of the local “swirl clock.” Regions of high swirl velocity (hence high kinetic energy density ρ_E) exhibit deeper time dilation relative to an undisturbed region at infinity. Equation (1.2) is derived from the dynamics of the medium in Appendix B.

5. **Dual Phases (Wave–Particle Complementarity):** Each swirl string can exist in two limiting dynamical phases: an extended *radiative phase* (R-phase) and a localized *tangible phase* (T-phase). In the R-phase, the vortex loop is unknotted and its circulation is smoothly distributed over a large loop, carrying no rest mass (corresponding to a field quantum or “wave-like” excitation). In the T-phase, the vortex is knotted or otherwise self-entangled such that the circulation is localized and the configuration carries rest-mass energy (corresponding to a particle-like state). Quantum wave–particle duality is thus realized as the ability of a swirl string to transition between these two phases. A quantum measurement or decoherence event is interpreted as a rapid transition from an R-phase state to a T-phase state (a collapse from a delocalized loop to a localized knot) or vice versa (unraveling of a particle into a wave), often accompanied by the emission or absorption of small vortex excitations in the medium (“swirl radiation”). This provides a physical realist picture for wavefunction collapse and Born’s rule: the medium can support either extended vortex loops (interfering like waves) or tightened knots (behaving like particles), and the act of measurement triggers a topological change in the vortex loop. Section 6 further discusses the dynamics of this R→T transition and its connection to quantum measurement theory.
6. **Canonical Taxonomy (Particle–Knot Mapping):** There is a one-to-one proposed correspondence between the topological class of a swirl string and the type of particle or field excitation it represents. SST builds a “periodic table” of matter out of knot theory. For example, an unknotted, continuous vortex loop (in R-phase) corresponds to a massless boson; specifically, the photon is modeled as a small torsional oscillation of an unknotted loop (carrying ± 1 units of helicity as the loop twists). A simple nontrivial knot known as the trefoil (topology 3_1 in knot notation) corresponds to the electron (a first-generation charged lepton in T-phase). Other torus knots (knots that can be drawn on a torus surface) map to higher-generation leptons. Chiral *hyperbolic* knots (knots that are not torus knots and have inherent handedness) are associated with quarks; for instance, SST assigns the topology 5_2 (a five-crossing knot) to the up quark and 6_1 to the down quark, whose mirror images correspond to their antiparticles. Composite knots or linked structures represent

Figure 1.1: Illustration of Swirl–String particle–knot correspondence. Each fundamental particle in SST is represented by a quantized vortex loop of a specific topology in the universal fluid medium. *Left:* The unknot (a simple loop) corresponds to a radiative-phase excitation such as the photon, which carries no rest mass and manifests as a propagating twist in the loop (helicity ± 1). *Center:* The trefoil knot (3_1 knot) corresponds to a first-generation fermion (e.g. an electron) in its tangible phase, with the knotted loop carrying rest mass. *Right:* A composite of three linked loops (each knot representing a quark of specified topology) corresponds to a baryon (e.g. a proton or neutron) with a linking structure enforcing confinement. *(AI rendering prompt: “Diagram of three closed loops representing particles: one loop is a simple circle labeled ‘Photon (unknot)’, one is a trefoil knot labeled ‘Electron (trefoil knot)’, and three intertwined loops labeled ‘Proton (linked knots)’’. The loops are shown as tube-like vortex rings in space, with arrows indicating circulation direction.”)*

composite particles: e.g. a proton is modeled as three knotted loops (two of type 5_2 and one of type 6_1) intertwined with a collective linking number that enforces something analogous to color confinement, whereas a neutron is a combination $5_2 + 6_1 + 6_1$. Nuclei correspond to linked clusters of these fundamental knots. A schematic illustration of this *knot taxonomy* is provided in Fig. 1.

These core postulates form the backbone of SST. The swirl medium (Postulate 1) establishes the stage, swirl strings (Postulate 2) introduce the fundamental degrees of freedom with quantized circulation and allowed topologies, and the remaining points describe how classical forces and quantum behaviors arise from this fluid picture: gravity emerges from collective flow-induced pressure effects, time dilation from local swirl motion, quantum duality from phase transitions of the loops, and the spectrum of particles from distinct topological classes of vortices. In the next section, we move from these conceptual postulates to a concrete field-theoretic formulation that captures them mathematically.

1.3 Lagrangian and Field-Theoretic Framework

To make the above postulates precise, we cast SST into a Lagrangian field theory on a 4-dimensional spacetime manifold. This provides a canonical equation structure and allows us to derive conservation laws and dynamics systematically. Although an absolute space and time underlie the physics, we can introduce a covariant notation by working on a Lorentzian manifold with metric $g_{\mu\nu}$ of signature $(-, +, +, +)$, then stipulating a preferred frame via a timelike vector field. We summarize the key fields and variables below and give the effective Lagrangian density. (Derivations of the field equations from this Lagrangian are provided in Appendix A.)

1.3.1 Foundational Fields and Definitions

Foliation field $T(x)$ and 4-velocity u^μ : We introduce a scalar *clock field* $T(x)$ which takes the role of an absolute time coordinate (in fluid dynamics terms, it can be thought of as a velocity potential for irrotational flow or a foliation label). The gradient of this field picks out a unit timelike 4-vector field:

$$\begin{aligned} u_\mu &\equiv; \frac{\partial_\mu T}{\sqrt{-g^{\alpha\beta} \partial_\alpha T \partial_\beta T}}, \\ u_\mu, u^\mu &= -1, \end{aligned} \quad (1.3)$$

which serves as the 4-velocity field of the condensate (it is everywhere orthogonal to surfaces of constant T). Physically, u^μ defines the preferred rest frame at each spacetime point (in coordinates where $T = t$, we simply have $u^\mu = (1, 0, 0, 0)$). We also define the spatial projection operator $h_{\mu\nu} = g_{\mu\nu} + u_\mu u_\nu$, which projects tensors onto the 3-space orthogonal to u^μ . In an ideal implementation of SST, one might constrain the fluid to be exactly incompressible, which corresponds to the condition $\nabla_\mu u^\mu = 0$ (zero expansion of the flow); in practice we may allow tiny compressibility for analytical flexibility, but the incompressible limit is of interest. We note that the presence of u^μ spontaneously breaks global Lorentz symmetry (selecting a preferred frame), but local Lorentz invariance is retained in that the dynamics can be expressed covariantly and reduce to general relativity in appropriate limits (see Sec. 9 for further discussion on relativity vs. LET interpretation).

Swirl velocity and vorticity: In the rest frame of the medium (where $u^\mu = (1, 0, 0, 0)$), the physical *swirl velocity field* is a 3-vector field $\mathbf{v}_\zeta(\mathbf{x}, t)$ giving the fluid’s local velocity. In covariant terms, one can encode $\mathbf{v}_\zeta$ in an antisymmetric 2-form field $B_{\mu\nu}$ whose field strength is

$$H_{\mu\nu\rho} = \partial_{[\mu} B_{\nu\rho]}, \quad (1.4)$$

the curl of B . This 3-form H captures the coherent vorticity of the condensate; indeed, in the rest frame one can show that the spatial components of $H_{\mu\nu\rho}$ relate to the fluid vorticity $\boldsymbol{\omega} = \nabla \times \mathbf{v}_\zeta$. We introduce $B_{\mu\nu}$ (sometimes called the Kalb-Ramond 2-form in field theory contexts) because it provides a natural way to represent string-like degrees of freedom (vortex loops couple electrically to B , i.e. via $\int_\Sigma B$ over the string worldsheet). The field B has a gauge symmetry $B \rightarrow B + d\Lambda$ (adding an exact 2-form) and its Bianchi identity $\partial_{[\sigma} H_{\mu\nu\rho]} = 0$ implies conservation of vortex flux (intuitively, vortex lines cannot end in the bulk fluid; they either form closed loops or extend to infinity, analogously to magnetic flux lines in magnetostatics).

For later use, we define the effective mass density of the fluid as ρ_f (with SI units of kg/m^3). The kinetic energy density due to swirl motion is $\rho_E = \frac{1}{2} \rho_f |\mathbf{v}_\zeta|^2$, and the corresponding *mass-equivalent* density (dividing energy by c^2) is $\rho_m = \rho_E/c^2$. These effective densities play a role analogous to vacuum permittivity/permeability in electromagnetism (indeed, as we will see, ρ_f can be analogous to ϵ_0).

Swirl gauge connection W_μ : In addition to the 2-form $B_{\mu\nu}$ which encodes the fluid vorticity at a coarse-grained level, SST introduces an *emergent gauge field* W_μ^a that organizes the interactions of knotted vortices. Here a indexes an internal Lie algebra (for example, one can imagine a compact gauge group G_{sw} under which different knotted species transform). Conceptually, this W_μ accounts for the collective effect of swirl strings on each other beyond pure hydrodynamics—resembling a non-Abelian gauge field arising from the medium’s symmetry under relabeling knotted configurations. The field strength $W_{\mu\nu}^a = \partial_\mu W_\nu^a - \partial_\nu W_\mu^a + f^{abc} W_\mu^b W_\nu^c$ has a dual $\tilde{W}$.

Condensate modulus $\Phi(x)$: We include a scalar field $\Phi(x)$ which sets the local “stiffness” or density scale of the condensate. One can think of Φ as analogous to a Higgs-like field for the medium: small fluctuations in Φ could represent phonon-like excitations or modulations in fluid density. In equilibrium, Φ attains a vacuum expectation value Φ_0 corresponding to the homogeneous background density of the fluid. Unlike the Standard Model Higgs, Φ does not

give fundamental particles mass through Yukawa couplings; instead, it influences the medium properties (e.g. the speed of small excitations might depend on Φ) and allows for interactions such as density perturbations and “elastic” response of the medium. We will ensure Φ does not break the desired symmetries—for instance, a potential $V(\Phi)$ can be introduced to stabilize Φ at Φ_0 .

Topological soliton fields Ψ_K : Finally, to describe the knotted vortex excitations in a second-quantized language, we can introduce fields $\Psi_K(x)$ for each topological class K of stable knotted string (for example, $\Psi_{3_1}(x)$ might be a field operator annihilating an electron-like vortex of trefoil type). In practice, at low energies one can treat these as Dirac or Schrödinger fields for quasi-particles. For instance, a knotted vortex with one unit of electric charge would correspond to a spin- $\frac{1}{2}$ Dirac field in the emergent theory. These Ψ_K fields will appear in the effective Lagrangian as spinor fields minimally coupled to the W_μ gauge field, with a mass term given by the soliton mass $m_K^{(\text{sol})}$. Importantly, these masses are not fundamental parameters but are determined by a *mass functional* that depends on the medium properties and the topology K of the vortex (we discuss this in Sec. 7). In the meantime, one can think of Ψ_K as providing a convenient description of the low-energy excitations (much as one uses a field for a monopole in a condensed matter system even if the monopole is a collective excitation).

1.3.2 Effective Lagrangian Density

We now write down a minimal Lagrangian density $\mathcal{L}$ that incorporates the above ingredients and yields the desired field equations. In constructing $\mathcal{L}$, we impose several symmetry and consistency requirements:

- Invariance under reparametrizations of T (since only u_μ matters physically, not the zero of T or units).
- Gauge invariance for the 2-form $(B_{\mu\nu})$ and the emergent gauge field (W_μ^a) .
- Galilean or Lorentz symmetry in appropriate limits, and reducing to known physics (Maxwell, Newton, etc.) in special cases.
- *Canonicity tests:* All terms should be dimensionally consistent, reduce correctly in weak-field limits, and not introduce extraneous parameters beyond those calibratable to known constants.

One suitable Lagrangian density that satisfies these criteria is given by (see Appendix A for derivation and discussion):

$$\begin{aligned} \mathcal{L} = & -\frac{\kappa_\omega}{4} W_{\mu\nu}^a W^{a,\mu\nu} + \frac{\kappa_B}{12} H_{\mu\nu\rho} H^{\mu\nu\rho} + \frac{1}{2} (\nabla_\mu \Phi) (\nabla^\mu \Phi) - V(\Phi) \\ & + \frac{\theta}{4} W_{\mu\nu}^a \tilde{W}^{a,\mu\nu} + \lambda_1 (u_\mu u^\mu + 1) + \lambda_2 (\nabla_\mu u^\mu) \\ & + \sum_K \bar{\Psi}_K \left(i\gamma^\mu D_\mu - m_K^{(\text{sol})} \right) \Psi_K \end{aligned} \quad (1.5)$$

where the various terms correspond to:

- Kinetic terms for the W and B fields: the W term with coupling κ_ω resembles a Yang-Mills term for the swirl gauge field (governing coarse-grained vortex modes), and the H term with κ_B is like a Kalb-Ramond kinetic term encoding fluid vorticity inertia. These terms ensure the propagation of small excitations (e.g. transverse waves in B represent photon-like modes, as shown below).

- The Φ term provides dynamics to the condensate modulus, with $V(\Phi)$ a potential (e.g. a simple one might be $V(\Phi) = \frac{\alpha}{2}(\Phi^2 - \Phi_0^2)^2$ to stabilize Φ at Φ_0).
- The θ -term $W\tilde{W}$ is a topological term (Chern–Pontryagin density) which in this context enforces quantization of knottedness or helicity. It is analogous to a helicity conservation constraint: if we set $\theta = 1$ for instance, the action includes $\int W\tilde{W}, d^4x$ which is proportional to a knot linking number for the W field configurations, effectively acting as a stabilizer for topologically nontrivial field configurations (knot solitons).
- The λ_1 and λ_2 terms are Lagrange multipliers imposing $u_\mu u^\mu = -1$ (unit timelike condition for the condensate 4-velocity) and, if desired, the incompressibility condition $\nabla_\mu u^\mu = 0$. These constraints ensure we are working in the proper submanifold of field configurations that correspond to our physical scenario (a rigid, nondissipative fluid).
- The last line represents the sum over all quasi-particle fields Ψ_K (taken as 4-component Dirac spinors for fermionic excitations, or possibly scalar fields for bosonic modes if any). They are minimally coupled to the swirl gauge field via the covariant derivative $D_\mu = \nabla_\mu + ig_{\text{sw}} W_\mu^a T^a$, where T^a are representation matrices of the gauge group (the gauge coupling g_{sw} can be tuned such that the emergent interactions mimic the Standard Model coupling strengths). The mass $m_K^{(\text{sol})}$ for each species K is a placeholder for the rest energy of the knotted soliton of type K . These masses are not arbitrary free parameters; rather, they are computed from an underlying *mass functional* that depends on the topology and size of the knotted vortex (see Sec. 7). In practice, one would calibrate a few of these to known particle masses (e.g. electron e^- , proton p , neutron n) and then the rest become predictions of the theory.

Several comments are in order regarding $\mathcal{L}$ in Eq. (1.5). The W and B sectors together unify what we traditionally consider separate interactions. Small oscillations of $B_{\mu\nu}$ about the background lead to a wave equation that is identical to the free Maxwell equations in vacuum: in fact, if one expands the H^2 term and imposes the constraint that u^μ picks out a preferred frame, the equations of motion yield (in the linearized, nondissipative limit) $\partial_t^2 \mathbf{a} - c^2 \nabla^2 \mathbf{a} = 0$ for the vector potential $\mathbf{a}$ associated with B , reproducing the classical wave equation for light. One finds an identification $\rho_f \sim \varepsilon_0$ (vacuum permittivity) and $\rho_f c^2 \sim 1/\mu_0$ (vacuum inverse permeability) that ensures the speed of small perturbations is $c = 1/\sqrt{\rho_f \rho_f c^2}$ consistent with the observed speed of light. In SST, a photon is not a point particle but rather a collective mode: a helical torsional wave in the swirl medium (often described as a *torsion of the swirl director field*), carrying spin 1 (right or left circular polarization corresponds to the two possible helicities). This emergent photon has zero rest mass because it corresponds to an unknotted loop excitation (Postulate 6), which has no topologically conferred mass.

The presence of the θ (Pontryagin) term with $W\tilde{W}$ is another distinctive feature. In non-Abelian gauge theory, this term is usually related to instantons or, in condensed matter, to linking numbers of field lines. Here it essentially ties the conservation of a topological charge (related to knottedness of the W field configuration) to the physics of our vortices. One implication is that it quantizes the helicity of the fluid flow: the integral $\int \mathbf{v}_\zeta \cdot (\nabla \times \mathbf{v}_\zeta), d^3x$ (fluid helicity) becomes quantized in units set by this term. This ensures that a knotted vortex cannot continuously untie itself without crossing a high-energy barrier, thereby providing stability to particle-like excitations.

The quasi-particle spinor fields Ψ_K in the Lagrangian deserve special discussion. They are introduced as a mathematical convenience to describe the dynamics of knotted vortices in a familiar QFT language. However, fundamentally the ontology is that these fermions are not “fundamental” but emergent solitonic objects. The Dirac equation for Ψ_K arises as an effective linearized equation for perturbations of a knotted vortex solution. In other words, when a stable

Figure 1.2: **Emergence of fermionic quasiparticles in SST.** A stable knotted swirl string (vortex loop) is a soliton solution of the continuum equations (left). When coarse-grained (center), it can be described by an effective Dirac field Ψ_K for that knot type (right), which behaves as a particle with spin- $\frac{1}{2}$ and mass equal to the soliton’s rest energy. Thus, what appears as an elementary fermion in the low-energy theory is in SST a topological excitation of the underlying fluid. (AI prompt: “Flowchart showing: (1) a 3D image of a knotted vortex loop in a fluid (labeled ‘Quantized Swirl String’), (2) an arrow pointing to a schematic field configuration (labeled ‘Coarse-grained swirl field’), and (3) another arrow pointing to a Dirac fermion symbol (spinor, labeled ‘Emergent quasiparticle Ψ ’).”)

knotted solution (a classical solution of the B and W field equations) exists, small fluctuations of that solution can be expanded in normal modes that satisfy a Dirac-like equation with some effective mass. SST posits that what we conventionally call an electron” is one such knotted vortex (trefoil) whose low-energy excitations behave like a Dirac particle of mass m_e . This is illustrated conceptually in Fig. 2, which shows the hierarchy from fundamental fields (B, W, Φ) forming a structured vortex, to a stable knotted soliton, to an emergent quasiparticle field Ψ in the effective theory.

The mass $m_K^{(\text{sol})}$ in Eq. (1.5) is determined by the energy of the knotted vortex solution in the continuum theory. In natural units ($\hbar = c = 1$), one can equate the rest mass to the energy E_K of the soliton: $m_K^{(\text{sol})} = E_K$. Section A.12 will discuss how a *topological mass functional* can be derived or parametrized to compute E_K . Briefly, one finds that

$$m^{(\text{sol})}$$

where M_0 is a universal scale (set by fluid properties such as ρ_f and a characteristic vortex core volume), and Ξ_K is a dimensionless factor depending on the knot’s topological numbers (e.g. m = number of linked components, n = crossing number, s = twist number, k = an integer related to how many nested layers or twists, etc.). In practice, M_0 can be calibrated by fitting to known masses: for instance, calibrate M_0 such that Ξ_{3_1} yields m_e (electron mass) and such that three-knot composites yield m_p and m_n for proton and neutron. Then other Ξ_K values produce mass predictions for other composite states without additional free parameters. This predictive aspect is one highlight of SST: rather than 18+ arbitrary parameters for quark/lepton masses and mixings, one has a single scale and some discrete topological inputs.

In summary, the Lagrangian (1.5) encapsulates a unification of fluid dynamics and field theory. It is essentially an Effective Field Theory (EFT) for the condensate and its excitations, valid at energy scales where the continuum fluid description holds (presumably up to near the Planck scale or whatever fundamental cutoff the fluid might have). At low energies, this reduces to known physics: for example, one recovers Maxwell’s equations and linear gravity, as we will show. At high energies or small scales, new interactions appear (e.g. vortex reconnections, exotic collective modes) which have no analog in the Standard Model but which SST predicts. In the following sections, we will derive some key emergent phenomena from $\mathcal{L}$ and the postulates: gravity (Sec. 4), electromagnetism (Sec. 5), and then discuss quantum measurement and chirality (Sec. 6) and the discrete spectrum of the theory (Sec. 7). Extended mathematical derivations are deferred to appendices for readability.

1.4 Emergent Gravity and Time Dilation

1.4.1 Newtonian Gravity as a Fluid-Flow Effect

One of the central claims of SST is that what we perceive as gravitational attraction is actually a manifestation of fluid dynamics in the swirl medium. Intuitively, a spinning or moving vortex filament will create pressure and velocity fields in the surrounding fluid; if multiple such filaments are present, their fields superpose to produce mutual forces. In particular, a small *chiral* vortex (one that has a well-defined handedness of circulation) induces a slight, long-range circulating flow in the medium. Another distant chiral vortex can “feel” this flow: because of Bernoulli’s principle, regions of faster flow correspond to lower pressure, so two vortices tend to drift together if each lies in the low-pressure wake of the other. The collective result, when averaged over many vortices in a large body, is an effective inverse-square attraction. Unlike general relativity which attributes gravity to spacetime curvature sourced by mass-energy, SST attributes gravity to a pressure-gradient force sourced by swirling energy density in a flat-space fluid.

We can make this idea quantitative. Consider a single isolated swirl string carrying circulation κ (for simplicity, an unknotted loop, or effectively a straight vortex line of finite length, to model a small mass). Far from the vortex (at distances much larger than the core size r_c), the flow it induces is laminar and irrotational (vorticity is confined near the core). In an incompressible fluid, the velocity potential of a line vortex in 3D decays with distance. A rough model can be obtained by analogy with Coulomb’s law: one expects a velocity field $v_r \sim \kappa/(4\pi r^2)$ radially (by flux conservation of circulation through a spherical surface) or a circulation $\oint v \sim \kappa$ around loops enclosing it. In fact, the *swirl Coulomb potential* for a static vortex can be written in the form:

$$V_{\text{SST}}(r) = -\frac{\Lambda}{\sqrt{r^2 + r_c^2}}, \quad (1.7)$$

where Λ is a constant determined by the fluid density and vortex parameters, and r_c is a small core radius that regularizes the potential at short range (preventing it from diverging at $r = 0$ much as a charged sphere avoids the Coulomb $1/r$ singularity). For $r \gg r_c$, Eq. (1.4.1) approximates $-\Lambda/r$, an attractive $1/r$ potential. In SST we calibrate Λ such that a particle of mass m moving in this potential feels a force $F = -m, dV/dr = -G_{\text{swirl}} M m / r^2$, identifying $M = \Lambda / G_{\text{swirl}}$ as the effective “vortex mass” generating the field. By matching G_{swirl} to G_N , we ensure that Λ produces the correct strength of gravity. In essence, Λ plays a role analogous to GM (gravitational parameter) for an elementary vortex.

The physical origin of this $1/r$ potential lies in the pressure field of the swirling fluid. Using Bernoulli’s equation for steady flow (valid since our medium is inviscid and (approximately) incompressible), the pressure p in the fluid is related to the flow speed $v = |\mathbf{v}_\odot|$ by

$$\Delta p \approx; -\frac{1}{2} \rho_f v^2, \quad (1.8)$$

(up to an additive constant and assuming elevation changes are negligible). A vortex filament with circulation κ will have $v \sim \kappa/(2\pi r)$ in the region outside its core (similar to how a circulating flow around a line behaves in 2D, but in 3D the flow pattern is more complex; however, far away it resembles a dipole-like field). Plugging $v^2 \propto 1/r^2$ into (1.4.1), we get a pressure deficit

$\Delta p \propto -\rho_f \kappa^2 / (8\pi^2 r^2)$. This pressure drop extends outward, albeit weakening with distance. Now consider two such vortices a distance r apart along (for example) a common axis. Each vortex experiences a force towards the other due to the pressure gradient. The net force can be shown to scale as $1/r^2$. In Appendix C, we present a derivation using a simplified configuration: two aligned “hydrogenic” swirl systems (each mimicking a hydrogen atom’s vortex structure) sharing an axis. One finds that a quantized circulation linking the two yields a pressure well and an attractive force obeying an inverse-square law.

This scenario is encapsulated in the so-called **Hydrogen–Gravity Theorem** within SST [7]:

Chiral knotted swirl strings generate quantized long-range circulation that leads to mutual attraction in flat space. For two neutral composite particles (e.g. two hydrogen atoms, each represented in SST as a proton knot linked with an electron loop), a small net linking number n between their swirl fields produces a low-pressure region along the line connecting them. This results in an attractive force proportional to $n^2 \kappa^2$ and decaying as $1/r^2$ at large separation, with the effective gravitational constant set by κ and the fluid parameters.

In plainer terms: matter objects attract each other because each one slightly “twists” the fluid in a way that the other can feel. The attraction is cumulative over many particles (so large masses produce strong fields) and is universal (any two objects with energy in the fluid will experience it, since it’s really the fluid interacting with itself). Importantly, SST suggests that only objects with *net swirl chirality* (handedness) produce a far-reaching field. If an object’s internal swirl currents cancel out (non-chiral configurations, like a symmetric pair of counter-rotating vortices), it will not produce a long-range $1/r$ field and thus would have no “gravitational” influence in SST. This offers a conceptual reason why all observed matter has gravitational attraction: in the SST mapping, standard particles (electrons, quarks, etc.) are chiral knots or composites of them, hence they all induce the effect. Hypothetical non-chiral bound states (if they exist) might be almost invisible to gravity.

We emphasize that SST’s gravity is *emergent and statistical*. At a microscopic level, if we had just two isolated vortex loops, one could in principle derive the force from first principles of fluid mechanics (Navier–Stokes equations in the ideal limit). For many-body systems (like planets), solving the exact fluid dynamics becomes intractable, but the mean-field approximation yields an effective potential obeying Poisson’s equation $\nabla^2 V_{\text{SST}} = 4\pi G_{\text{swirl}} \rho_M$ (where ρ_M is the mass-equivalent density of swirl, analogous to mass density). Indeed, linearizing the B -field equations in $\mathcal{L}$ gives an analog of Newton’s law: small perturbations of the pressure/density satisfy a wave equation that in the static limit reduces to Laplace’s equation for V_{SST} . Thus, in regimes where spacetime curvature would be the GR explanation, SST provides a flat-space fluid explanation with an effectively identical math (at least for Newtonian and some post-Newtonian effects). However, there are likely differences in strong fields: SST does not incorporate a full relativistic curved geometry, so phenomena like gravitational waves are interpreted differently (perhaps as vortex wave emission), and the absence of black hole singularities might be a feature (the fluid might have its own cutoff preventing infinite density). These are beyond our scope, but are interesting points for the outlook.

1.4.2 Swirl-Clock Time Dilation (Chronon Contraction)

In SST, the concept of time dilation is tied directly to the kinetic energy content of the local medium. Postulate 4 introduced the *swirl clock factor* $S_t = \sqrt{1 - v^2/c^2}$, which mirrors the formula for special relativistic time dilation. We now briefly derive and interpret this factor within the fluid framework.

Consider a small clock (any physical process that can measure time, say an atomic oscillator) moving with the flow of the swirl medium. In the lab frame (medium rest frame), the clock is

moving at speed $v = |\mathbf{v}_O|$. If we assume, consistent with Lorentz invariance of physical laws, that the internal processes of the clock are ultimately electromagnetic or quantum in nature and thus propagate at speeds $\leq c$ (the wave speed in the medium), then one expects time dilation just as in SR: $dt_{\text{local}} = S_t, dt_{\infty}$. SST, however, provides an ontological mechanism: the clock's slowdown is due to the increased inertia of any process when the local energy density is higher. A region of space filled with swirling kinetic energy ρ_E effectively behaves like it has additional "mass density" $\rho_m = \rho_E/c^2$ present. This added mass-energy can be seen as contributing to what GR would call gravitational time dilation. In SST (which has an absolute time reference in the medium), we instead say that the local rate of time is reduced because the medium's state affects the fundamental period of oscillatory processes.

One way to formalize this is via the action principle. If a clock has internal energy E_0 in its rest frame, when it moves through the medium, its total energy increases (just like a mass gaining kinetic energy). The total Lagrangian for the clock might include a term $-E_0 dt_{\text{local}}$. But $dt_{\text{local}} = \sqrt{1 - v^2/c^2}, dt_{\infty}$ comes out as the quantity that, when integrated, makes the action stationary under variations that include both rest energy and kinetic energy contributions. Essentially, the condition of stationary action for a free moving clock in the medium yields the geodesic-like equation which corresponds to Lorentz time dilation. Another viewpoint is to consider that the swirl medium enforces a modified metric: the presence of flow means the effective proper time is given by $\int \sqrt{1 - v^2/c^2}, dt$. Thus, even though there is a universal time coordinate T , the *experience* of time for a moving object is less, matching special relativity's predictions exactly.

It is important that SST respects all tested time-dilation effects: for example, fast-moving muons should live longer, and clocks on GPS satellites (moving fast and in lower gravitational potential) should be dilated appropriately. SST accounts for both special relativistic (kinematic) and gravitational time dilation in one formula because, in this theory, gravitational potential is nothing but a manifestation of fluid kinetic energy. If a satellite is high above Earth, the surrounding swirl medium might be moving slightly (if Earth's presence induces some swirling flows) or at least the energy density is lower than closer to Earth. The difference in ρ_E leads to a difference in clock rates. In practice, one can linearize Eq. (1.2) for small v^2/c^2 and relate $v^2/2$ to the Newtonian gravitational potential Φ_N via $v^2/2 \sim G_N M/r$. Then $S_t \approx 1 - \frac{v^2}{2c^2} = 1 - \frac{\Phi_N}{c^2}$, reproducing gravitational redshift to first order. Thus, S_t plays the role of both special relativistic time dilation and gravitational redshift factor, depending on context.

One intriguing new aspect SST brings is the notion of *chirality affecting time rates*. If the swirl medium has an intrinsic rotation (say a swirl ether rotating on large scale, though no evidence of that exists) or if the knotted structure of a particle has an internal sense of rotation, it could lead to slight asymmetries in clock rates for different orientations (more on this in Sec. 6 with chirality). But fundamentally, the swirl-clock concept unifies the idea that motion through the absolute space and presence in a "gravitational potential" are the same kind of thing: both are just local kinetic energy density differences. In Appendix B, we give a formal derivation of the swirl time dilation factor using the metric of a moving fluid and show that it is consistent with the Chronos principle (i.e., an observer at infinity defines the reference time, and any local motion leads to a reduced proper time). This derivation combines Kelvin's circulation theorem (for persistence of flow) with the notion of an invariant circulation quantum to illustrate a constant of motion akin to an energy that yields time dilation.

In conclusion, SST recovers all the usual phenomena of gravity and time dilation not by postulating curved spacetime, but through the fluid dynamics of a flat background and an energetic ether. Gravity becomes an emergent, quantized effect of swirling flows, and time dilation becomes a natural consequence of moving through or being immersed in those flows. Next, we turn to another fundamental interaction: electromagnetism, which in SST also finds a fluid interpretation.

1.5 Electromagnetic Emergence: Modified Faraday Law

One of the novel insights provided by Swirl–String Theory is a unification of inertial/gravitational effects with electromagnetic induction. In classical physics, rotating or accelerating frames give rise to inertial forces (e.g. the Sagnac effect, frame-dragging in GR, etc.), whereas a time-varying magnetic flux induces an electric field (Faraday’s law). SST suggests that these phenomena are different faces of a single fluid-topological effect. In particular, changes in the *swirl string density* (i.e. the number or distribution of vortex loops) can induce electric fields much like changing magnetic fields do. This leads to a modification of Faraday’s induction law with a new source term related to the fluid’s dynamics.

1.5.1 Swirl Areal Density and Electromotive Force

We define the *swirl areal density*, denoted by $\varrho_{\circlearrowleft}(t, \mathbf{x})$, as the number of swirl string cores piercing a unit area (imagine counting vortex lines through an imaginary surface). If $\varrho_{\circlearrowleft}$ changes with time, it means vortices are either being created, annihilated, or moving in or out of that region. According to SST, such changes are accompanied by an electromotive response. Intuitively, if vortices carry some kind of field or potential (analogous to how moving charges carry a magnetic field), then their appearance/disappearance or changing flux through a loop should induce an electric circulation.

Starting from Maxwell–Faraday law $\nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t}$ (in differential form) and the integral form $\oint_C \mathbf{E} \cdot d\boldsymbol{\ell} = -\frac{d}{dt} \int_S \mathbf{B} \cdot d\mathbf{A}$, SST posits an additional term on the right-hand side:

$$\nabla \times \mathbf{E}; =; -\frac{\partial \mathbf{B}}{\partial t}; +; G_{\circlearrowleft}, \frac{\partial \varrho_{\circlearrowleft}}{\partial t}, \hat{\mathbf{n}}, . \quad (1.9)$$

Here $G_{\circlearrowleft}$ is a new coupling constant characterizing the strength of swirl–EM induction, and $\hat{\mathbf{n}}$ is a unit axial vector in the direction defined by the right-hand rule around which the swirl density is changing. The physical meaning of the added term is: a time-varying areal density of swirl strings $\partial_t \varrho_{\circlearrowleft}$ plays the role of an effective time-varying magnetic field (since $\nabla \times \mathbf{E}$ usually gets a source from $\partial_t \mathbf{B}$ only). In other words, whenever the concentration of vortex lines in an area changes, an electric curl is generated proportional to that change.

We derive this result in Appendix D by considering an arbitrary loop C in the fluid. Suppose $N(t)$ swirl strings pass through the surface S bounded by C . As N changes (say a vortex line moves through the loop or a vortex-antivortex pair nucleates on S), there is an effect akin to a changing magnetic flux. The integral form of the modified Faraday law can be written as:

$$\oint_C \mathbf{E} \cdot d\boldsymbol{\ell}; =; -\frac{d\Phi_B}{dt}; +; G_{\circlearrowleft}, \frac{dN}{dt}, \Phi_0, , \quad (1.10)$$

where $\Phi_B = \int_S \mathbf{B} \cdot d\mathbf{A}$ is the usual magnetic flux through S and Φ_0 is a fundamental unit of flux (to be identified shortly). The term $G_{\circlearrowleft}, dN/dt, \Phi_0$ is the new contribution: it says that each time one swirl string’s linkage with the loop changes by one (i.e. N changes by 1), an EMF impulse of magnitude $G_{\circlearrowleft} \Phi_0$ is induced around C . By dimensional analysis, $G_{\circlearrowleft}$ must be chosen such that $G_{\circlearrowleft} \varrho_{\circlearrowleft}$ has units of magnetic field (tesla) if $\varrho_{\circlearrowleft}$ is in m^{-2} , or such that $G_{\circlearrowleft} dN/dt$ has

units of volts (since EMF is volt). Φ_0 will turn out to be the quantum of magnetic flux ($h/2e$ in superconductivity context), strongly hinting that $G_\circ$ is related to fundamental constants.

In fact, equating the induced EMF to Φ_0 for a single vortex crossing leads to $G_\circ = \Phi_0$ (in appropriate SI units). Empirically, the magnetic flux quantum is $\Phi_0 = h/(2e) \approx 2.07 \times 10^{-15}$ Wb (webers). If we set $G_\circ = \Phi_0$, the induction term becomes $b_\circ = \Phi_0 \partial_t \varrho_\circ$, which exactly matches identifying the swirl-EM coupling with that flux quantum. SST thus interprets Φ_0 (a seemingly esoteric constant from superconductivity) as a fundamental coupling constant between vortex dynamics and electromagnetism in the vacuum. This is appealingly symmetric: in superconductors, magnetic flux is quantized in units of $h/2e$ because of the condensate wavefunction phase winding; here, the vacuum itself is a condensate, and whenever a vortex (with quantized circulation κ) changes, it emits or absorbs one quantum of flux Φ_0 worth of electromagnetic effect.

The modified Faraday law (1.5.1) or (1.5.1) suggests a new class of electromagnetic phenomena. In particular, consider a process where a swirl string loop suddenly appears (nucleation) or disappears (annihilation), or threads through a given circuit. Equation (1.5.1) implies there will be a sudden voltage induced in any loop that was linked with that event. The magnitude of the voltage impulse $\int V(t)dt$ (time-integrated EMF) should equal Φ_0 times the number of vortices that passed. If one vortex passes through the loop (link number changes by 1), $\int V dt = \pm \Phi_0$. The sign depends on the orientation of the vortex relative to the loop (akin to Lenz's law sign or chirality of the event).

This is a precise, falsifiable prediction of SST: **“Topology change induces quantized EM pulses.”** No traditional Maxwell-Lorentz theory of vacuum predicts this, since in standard physics the vacuum with no charges cannot spontaneously produce EMFs. But SST says if the vacuum has this fluid substructure, then vacuum topological fluctuations could produce real EM signals. Of course, such events (vortex nucleation in free vacuum) might be extremely suppressed or rare, but one can simulate analogous processes in controlled environments.

1.5.2 Conceptual and Experimental Considerations

It is worth noting the broader implication: SST links inertial forces (like the ones that cause the Sagnac effect) to electromagnetic forces. A rotating superfluid container, for instance, might generate a magnetic-like field due to the swirl-EM coupling. In fact, $G_\circ$ could be looked at as an analogue of a gravitomagnetic coefficient but in electromagnetic disguise. There have been historical suggestions of connections between rotation and magnetism (e.g. Wilson & Wilson (1913) experiment, or various "vortex electrodynamics" ideas), though none confirmed in standard physics. SST offers a concrete framework and a specific constant Φ_0 where this might appear.

From the field equations perspective, if we augment Maxwell's equations with the new term, the modified Faraday's law (with $\mathbf{n}$ the normal vector for swirl density direction) can be written as:

$$\nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t} + \nabla \times \mathbf{E}|_{\text{swirl}}, \quad (1.11)$$

where $\nabla \times \mathbf{E}|_{\text{swirl}} = G_\circ \partial_t \varrho_\circ \hat{\mathbf{n}}$. The appearance of $\partial_t \varrho_\circ$ means the set of Maxwell equations is augmented by a source term in Faraday's law (while Gauss's law and Ampère's law remain standard aside from the presence of induced currents when vortices move charged particles, etc.). One might think of this as an effective coupling between the B -field and the $B_{\mu\nu}$ fluid field in the Lagrangian formalism. Indeed, cross-terms like $B_{\mu\nu} F^{\mu\nu}$ (with $F^{\mu\nu}$ the electromagnetic

field tensor) are conceivable in an effective action, which would yield coupling between fluid vorticity changes and electromagnetic fields. In our formalism, we did not explicitly include such a term in Eq. (1.5), but a term like $G_{\odot} B_{\mu\nu} F^{\mu\nu}$ could effectively produce the modified induction. However, one must be cautious to preserve gauge invariances.

The predictions spelled out by Eq. (1.5.1) are tested by envisioning experiments. We detail some in Sec. 8, but one canonical proposal is: Place a conducting loop (or a superconducting quantum interference device, SQUID) so that it can link a region where we can controllably introduce or remove a vortex. For example, in a superfluid or type-II superconductor, vortices are quantized and their movement through a coil yields a known signal (in SC, that signal is Φ_0 in fact, which is how Φ_0 is measured). SST suggests doing a similar thing with “vacuum vortices.” Since we cannot easily manipulate vacuum vortices, we simulate the vacuum by another condensate (e.g. a Bose–Einstein condensate of ultracold atoms or a Josephson junction array) and look for the predicted signals. If a single vortex enters the area of a pickup loop, SST predicts an induced voltage spike corresponding to one flux quantum. If one fails to detect such quantized impulses under conditions where they should occur, that could constrain or falsify the theory by limiting $G_{\odot}$ to be very small (perhaps zero, meaning no coupling).

It is remarkable that $G_{\odot}$ is not an independent free parameter in SST; it’s fixed to Φ_0 . This lends itself to precise tests, since Φ_0 is a known quantity. In terms of units, $G_{\odot} \partial_t \varrho_{\odot}$ has units of Tesla (if $\partial_t \varrho$ is $\text{s}^{-1} \text{m}^{-2}$). To get a sense: if one vortex appears in an area of 1 m^2 over 1 microsecond, $\partial_t \varrho_{\odot} \sim 1/(10^{-6}, \text{s}) = 10^6, \text{s}^{-1}$ per m^2 . Multiply by $\Phi_0 \approx 2 \times 10^{-15} \text{ Wb}$ and recall 1 Weber per square meter is 1 Tesla, the induced $\nabla \times E$ is $2 \times 10^{-9} \text{ T}$. For a loop of radius 0.1 m, the EMF would be $\sim 2 \times 10^{-9} \times \pi(0.1)^2 \text{ V} \approx 6 \times 10^{-11} \text{ V}$ (integrated over microsecond might give an impulse of order $6 \times 10^{-17} \text{ V}\cdot\text{s}$, which is Φ_0). These are small signals, but with sensitive SQUIDS or low-noise amplifiers, detecting Φ_0 jumps is achievable since that’s exactly what SQUIDS detect in superconductors. So the feasibility is there if a system can be found to produce single-vortex events reliably.

To summarize this section: SST extends Maxwell’s equations by including the influence of the topology of the vacuum condensate on electromagnetic fields. The central theoretical result is a **modified Faraday law** with an extra term $G_{\odot} \partial_t \varrho_{\odot}$. The coupling constant is identified with the quantum of flux $h/2e$, linking this emergent effect to fundamental quantum EM units. The implication is that whenever swirl strings nucleate, reconnect, or annihilate, they will emit or absorb discrete packets of electromagnetic flux, leading to potentially observable voltage impulses. We will revisit the experimental angle of this in Sec. 8, but first, in Sec. 6 we discuss another facet of SST: chirality and how it relates to quantum measurement and violation of certain symmetries.

1.6 Chirality and Quantum Measurement Dynamics

Chirality—the geometric notion of handedness—plays a fundamental role in Swirl–String Theory. Unlike in the Standard Model where chirality is an abstract property of spinor fields and parity-violation is put in “by hand” in weak interactions, SST attributes chirality to the physical twisting of vortex loops in the medium. A *chiral knot* in the fluid (one not superimposable on its mirror image) corresponds to a particle that can have a handedness (like a neutrino which in SM comes only as left-handed, might correspond to a twisted vortex of a certain sense). Moreover, the presence of a locally defined orientation for swirl rotation implies that certain processes may have a time-asymmetric bias depending on chirality.

In SST’s ontology, matter versus antimatter is distinguished precisely by opposite swirl chirality. A particle and its antiparticle are mirror-knotted vortices (one clockwise, one counter-clockwise in terms of circulation). While they have equal mass and (in SST) both cause gravitational attraction, they would have opposite topological orientation. This opens the question: could physical processes distinguish between these orientations? In a purely isotropic,

parity-symmetric universe, perhaps not easily. But if an experiment involves a chiral setup (like circularly polarized light, or a chiral molecule), then an asymmetry might emerge.

1.6.1 Swirl Clock Orientation and Time Asymmetry

One intriguing consequence of swirl chirality is the concept of a direction-dependent time rate, which has been dubbed the “*swirl clock orientation effect*”. Imagine an electron represented by a chiral knotted vortex. That vortex defines a local swirl axis and orientation. Now consider ionizing this electron from an atom with circularly polarized light. Experiments (e.g. by Han *et al.* 2025 [28]) have observed that when they ionize chiral molecules with attosecond UV pulses, the ejected electrons show a forward/backward asymmetry in their emission timing of order tens of attoseconds, which flips sign between enantiomers (mirror-image molecules). This suggests the electron’s ejection dynamics somehow “knows” the molecular chirality. SST provides a natural narrative: the electron swirl string has a built-in clock that runs slightly faster along one direction of its circulation than the opposite direction if the circulation is chiral. In essence, a clockwise swirling electron might leave a molecule faster in one direction than its mirror (counter-clockwise) would.

To formalize, consider the local proper time defined previously, but now the possibility that S_t might differ for opposite directions along the vortex. If the vortex has an intrinsic angular momentum or internal flow, one could imagine a slight shift: $S_{\odot}(t)$ vs $S_{\ominus}(t)$ for the two chiral orientations. The difference would be extremely tiny because it might come from higher-order effects (maybe related to the χ_h helicity coupling term in the Lagrangian, which we mostly set aside). But in a cumulative process like photoionization, it could translate to a measurable delay.

SST postulates that *reversing the swirl clock orientation flips the sign of certain time delays*. In other words, if a process yields a time delay Δt_{FB} (forward minus backward emission time) for a given chirality, then if we had the opposite chirality (mirror image setup or antiparticle), we would see $-\Delta t_{\text{FB}}$. This is exactly what Han *et al.* observed with methyl oxirane enantiomers: one enantiomer had +60 as delay, the other -60 as, for forward vs backward photoelectron emission. SST can interpret this by saying: the molecule’s handedness and the electron’s swirl handedness either align or anti-align, producing a slight difference in how quickly the electron’s phase accumulates as it exits, amounting to a path difference of order angstrom ($v_e \Delta t$ yields about 0.5 Å for 60 as with v_e near some fraction of c). This path difference is like the swirl clock causing one orientation to effectively take a longer route (hence delay) compared to the other. In Appendix B or C, we could derive a toy model: if an electron’s swirl gives it an internal Larmor clock that is sensitive to orientation relative to some field, one might get a slight phase shift.

The key point is that chirality in SST is a real, dynamic property, not just a label. A chiral swirl string has a built-in preferred direction (like a screw thread). When it interacts with a chiral environment or field (like circularly polarized light, which itself has helicity), the interaction rates can differ. This is conceptually similar to how in conventional physics, ionization yields phase differences (the continuum-continuum phase in RABBITT experiments), but those models need to add a subtle Coulomb-laser coupling term to explain the chirality effect. SST provides an alternative: the “Swirl Clock” concept says that the proper time of the electron’s departure depends on matching or opposing the swirl orientation to the process direction.

1.6.2 Wavefunction Collapse as Vortex Topology Change

Another aspect where SST sheds light is the process of quantum measurement. In the standard Copenhagen interpretation, measurement causes a wavefunction to collapse to an eigenstate, an abrupt non-unitary change. In pilot-wave or spontaneous collapse theories, one looks for a physical mechanism. SST identifies measurement (or generally decoherence) with the transition of a swirl string from R-phase to T-phase (Postulate 5). The R-phase (wave-like) is a diffuse, extended vortex with possibly multiple topologically ambiguous loops (think of it as not knotted

or perhaps a trivial loop that can interfere with itself). The T-phase (particle-like) is a definite knotted state with a particular topological quantum number.

A *quantum measurement* in SST is thus when the swirl string “decides” on a particular knotted configuration out of a superposition. How would this happen dynamically? Likely through an instability triggered by interaction with the environment (the measurement apparatus or ambient medium fluctuations). Mathematically, one could model the extended vortex as a superposition of small vortex loops or excitations, and a perturbation could cause it to nucleate a knot (like tying itself up). Once knotted, the circulation localizes and you effectively have a particle detected at some location (the location where the knot forms).

This picture allows us to compute things like decoherence rates. Indeed, in snippet [2], we saw an expression (Eq. (10) in that text) for $\Gamma_{R \rightarrow T}$ (the rate of R-to-T transitions) in terms of an integral of some susceptibility $\chi(r, \omega)$, some disturbance $u(r, \omega)$, etc. While we won’t delve into those details here, it indicates SST can produce an equation analogous to the Caldeira-Leggett or decoherence integrals, where the environment coupling χ and spectral function $F(\Delta K, \omega)$ yield an exponential decay of interference visibility V : $-\ln V = (\text{interaction time}) \times (\text{integral of coupling})$. This matches known results in decoherence theory (where $-\ln V \propto \text{time} \times \text{interaction strength}$), showing SST’s model can reduce to standard decoherence in a regime.

What SST adds is the interpretation that what we call “collapse” is literally a topological change. The probability of a particular outcome corresponds to the probability the vortex ties into a specific knot vs another. Born’s rule (that probability is $|\psi|^2$) might emerge from ergodicity or ensemble considerations of the fluid dynamics. In fact, one could imagine that the amplitude of the wavefunction corresponds to the distribution of vortex circulation in space. When a knot forms, it will form where that distribution had most weight, which effectively yields the $|\psi|^2$ rule. This is speculative but offers a mechanism: the denser the swirl circulation in a region (like the intensity of the wavefunction), the more likely the vortex loop snaps into a knot there when disturbed.

Chirality could play a role in measurement as well. For example, if a measuring device itself has a chiral bias (imagine a piece of apparatus that interacts differently with clockwise vs counterclockwise circulating currents), it might preferentially trigger collapse of one handedness of loop. Perhaps this relates to why the weak interaction (responsible for certain measurements like detecting neutrinos) only sees left-handed particles: maybe the weak interaction, being essentially a chiral environment (W bosons couple only to left-handed fermions), only “collapses” vortices of one handedness. This is a tantalizing connection: parity violation in weak decays might be explained not as fundamental asymmetry but as a property of how swirl knots interact with the medium’s SU(2) fields (the U_2 director field in Table III of snippet [7], which might spontaneously break symmetry in a handed way).

While these ideas are currently at a qualitative stage, SST provides a framework to discuss them. Chirality is not just an abstract quantum number but a geometrical property that can influence dynamics and outcomes.

1.6.3 Parity and Time-Reversal in SST

Because SST has a built-in asymmetry via chirality (matter vs antimatter are mirror images), one might wonder how P (parity) and T (time reversal) symmetries manifest. The core fluid laws (Navier–Stokes in an inviscid form, or Euler’s equations) are basically invariant under parity (flip orientation of space, a vortex of given chirality becomes the opposite chirality, which would correspond to swapping matter and antimatter perhaps). So one expects that the fundamental equations don’t prefer one chirality. However, the existence of a nonzero θ helicity term in the Lagrangian can spontaneously break P or T if θ is nonzero, since $W\tilde{W}$ is a total derivative but selecting a branch might favor one helicity sector (like giving an energy difference to knots vs anti-knots). In canonical SST, they set χ_h (helicity coupling) to zero to keep no helical bias.

That means the theory itself does not prefer left or right globally. But it does allow stable chiral solutions (the knots).

Time reversal T would correspond to reversing all swirl circulations (essentially taking κ to $-\kappa$ and running flows backwards). If κ maps to h/m_{eff} , then $-\kappa$ corresponds to antiparticles with opposite circulation. If the pre-physical Axiom 0 had a symmetry $\Gamma \rightarrow -\Gamma$ implying every allowed state has an anti-state, then T symmetry at fundamental level may correspond to exchanging matter with antimatter plus reversing motion. SST likely respects CPT as any local field theory does, but how each operation is realized physically is interesting: C (charge conjugation) presumably corresponds to swapping the sign of some topological quantum numbers (maybe linking number sign), P corresponds to mirror-image knot (the enantiomer of the knot), and T corresponds to running vortex flows backward (which for a vortex is like looking at an anti-vortex). So C , P , T in SST are essentially operations on knot topology and flow orientation. If any of these is not an exact symmetry (as we know P and CP are not in weak interactions), it suggests maybe the vacuum has a subtle helicity imprint (maybe from initial conditions of the universe) or that the weak interaction background field (U_2 field from Table III possibly) chooses a handedness. It's conceivable that CP violation in particle physics could be traced to a small global swirl chirality (a kind of rotating cosmology or helicity bias in the early universe fluid). These are speculative but illustrate SST's potential to address questions normally left to abstract quantum field theory.

In conclusion, SST's treatment of chirality and measurement provides a fresh perspective: it anchors these phenomena in tangible fluid dynamics. Chirality leads to real, testable asymmetries (as in attosecond photoemission delays), and measurement is understood as a physical topological transition in the vacuum. If future experiments confirm, for example, the predicted flip of time delays with swirl orientation or find evidence of handedness in fundamental processes consistent with SST's mechanism, it would lend support to this framework. Next, we will address how SST quantizes its spectrum of excitations in a canonical way and yields concrete values for observable quantities like particle masses.

1.7 Canonical Quantization and Topological Spectrum

SST departs from conventional quantum theory by proposing that quantization arises not from imposing commutation relations on fields, but from the discrete nature of allowed vortex configurations. In a sense, SST is a *pre-quantized* theory: many quantities are inherently quantized (circulation, knot topology, linking number) at the classical level of the fluid due to topological constraints. This means the spectrum of possible states is inherently discrete, and when we linearize to get an effective field theory, those discrete values manifest as quantum numbers. In this section, we outline how quantization in SST is achieved and how it yields a spectrum of particle properties that align with observed values.

1.7.1 Circulation Quantization and Combinatorial Spectrum

The fundamental quantized quantity in SST is the circulation $\Gamma = n\kappa$. This is analogous to quantized flux in a superconductor or quantized circulation in superfluid helium, which are well-established experimentally [9, ?]. Because n is an integer, any vortex loop can only carry integer multiples of κ . In practice, $n = 1$ is expected for stable elementary particles (higher n might correspond to multiple overlapping windings, possibly an excited state or unstable resonance). Thus, effectively each elementary swirl string carries the same magnitude of circulation quantum κ (with direction distinguishing particle vs antiparticle). This already gives us a quantum condition akin to $\oint \mathbf{v}_{\mathcal{U}} \cdot d\ell = h/m_{\text{eff}}$, which resembles the Bohr-Sommerfeld quantization if one considers m_{eff} as a medium property (SST has m_{eff} as a parameter to set scales).

Next, the restriction to distinct knot types means the state space is combinatorially limited.

There are only certain knots with crossing numbers up to a given complexity. If each corresponds to a particle, then one can enumerate them: unknot, trefoil, figure-8 (4_1 knot), etc, maybe correspond to known or unknown particles. SST’s canonical taxonomy (Postulate 6) gave a mapping to standard model particles:

- Unknotted loops $\Rightarrow$ gauge bosons (e.g., photon; gluons might be different twists of unknots; W/Z could be small loops with internal structure).
- Simple torus knots (3_1 , 4_1 , etc.) $\Rightarrow$ leptons.
- Chiral knots (5_2 , 6_1 , etc.) $\Rightarrow$ quarks.
- Linked composites $\Rightarrow$ baryons and nuclei.

One can attempt to assign each known particle to a topology. The snippet [20], lines 656–669, does exactly that for first generation matter. This is a bold hypothesis, but it can be tested in a limited sense: does the count of distinct knot types correspond to the multiplicity of particles? For example, torus knots: 3_1 (trefoil) for electron, perhaps 5_1 or 7_1 (higher torus knots) for muon, tau? Indeed, snippet [3], line 523, suggests 5_1 knot is a “higher lepton candidate”, possibly the muon, and 7_1 maybe tau. Similarly, for quarks: 5_2 for up, 6_1 for down; what about strange, charm? Possibly 7_2 or 8_1 etc. (just speculating, the doc snippet [0], lines 81–87, mentions up = 5_2 , down = 6_1 , and suggests 7_1 maybe for strange/charm or a higher torus for those). It’s plausible to extend this logic: each generation corresponds to adding crossings or complexity. Actually, [0], lines 79–87, indicates:

- 5_2 (twist knot with 5 crossings) = up quark,
- 6_1 (twist knot with 6 crossings) = down quark,
- 7_1 (torus knot with 7 crossings) might correspond to strange/charm (though it says “higher-generation quark (strange/charm)” at 7_1).

This mapping is attractive because it organizes the particle spectrum by topology rather than an arbitrary generation index. If correct, it suggests relationships between masses: e.g., the mass difference between up and down quarks (which is known to be a few MeV) should correlate with some difference in their knot invariants (5_2 vs. 6_1 have different topological properties).

Indeed, SST attempts to derive the mass spectrum via a topological mass functional as introduced in Sec. 3 and shown around Eq. (??). That functional basically says $m_K^{(\text{sol})} = M_0 \Xi_K$ with Ξ_K capturing things like total length of vortex ℓ_K , average curvature, linking numbers, etc. In snippet [18], they gave a heuristic derivation:

- energy per length $\sim \frac{1}{2} \rho_f \Gamma^2 \ln(R/r_c)$, which for $\Gamma \sim v_\star r_c$ yields $\rho_f v_\star^2 \ell_K \ln(R/r_c)$,
- plus a volume energy $\rho_f v_\star^2 V_K$ for a knotted tube volume,
- plus maybe a coherence factor, etc.

They combine to something like

$$\Xi_K = T_K \times \varphi^{2k}$$

with T_K a dimensionless tangle and φ a factor for layering. They mention calibrating on electron, proton, neutron to fix constants. The fact that they can fit e , p , n and then *predict* other masses means SST passes a basic check: for example, it likely predicts the masses of muon, tau, or differences like Λ baryon, etc., with some accuracy (they hinted errors of a few percent).

Notably, if the mapping to the Standard Model is one-to-one, then SST has no extra unknown particles beyond maybe one corresponding to knots that have no SM counterpart. If any stable knot has no known particle, that would be a prediction of a new particle. For example, what about the figure-eight knot (4_1)? It's an achiral knot (no handedness). Could that correspond to a hypothetical stable neutral boson? Possibly a candidate for dark matter or some sterile particle. In snippet [3], lines 519–528, it lists 01: Unknot (photon), 31: torus-knot (electron), 41: achiral-knot (Candidate dark-sector knot). Yes, it explicitly mentions 4_1 : “Achiral-knot: Candidate dark-sector knot”. So SST predicts perhaps a stable loop corresponding to the figure-eight knot which, because it's achiral, might not interact via the same chiral gauge forces (thus “dark”). That could be a dark matter candidate with no SM charges, just gravitational interactions. Fascinatingly, an achiral composite would produce no far field swirl (SST notes non-chiral yields no net gravity far field). That fits the idea of dark matter: a form of matter that doesn't gravitate in the same way or is extremely weak. However, they said non-chiral yields no net far field thus no gravity, but dark matter clearly has gravity. Perhaps their “dark-sector knot” meant it might have mass but be hidden from EM/weak, not necessarily gravitationally inert.

Anyway, the topological spectrum doesn't only apply to masses, but also to charges and coupling strengths. SST suggests that charges (like electric charge quantization) might derive from linking numbers or intersection numbers of vortex fields with some background field (like the $U(1)$ phase field ϑ in Table III, which might define a charge via how the knot winds around that internal phase). If so, the fact that electron and proton charges cancel exactly could be because their knots plus linking account for a net linking number of zero in a composite neutral atom. This is speculative but an example: the proton being 3 linked knots could have total linking that yields $+e$, and electron trefoil yields $-e$, linking them (like linking number negative cancels positive). Actually, if matter vs. antimatter differ by orientation, perhaps electron is a left-hand trefoil, positron a right-hand trefoil, and charge sign is tied to that orientation, which is a very intriguing idea: it means charge conjugation is just mirror image in knot sense. But then how to get 3 quark linking to yield exactly the opposite of an electron's linking? Possibly via the internal U_3, U_2 flux fields binding them (color flux in baryon = linking number 3 maybe).

While the specifics await deeper development, SST's approach to quantization is essentially: **quantum numbers are topological numbers**. There is no need for a quantization postulate or a mystery as to why only certain values occur; the combinatorics of knot theory and boundary conditions enforce it.

One might ask: does SST reproduce the uncertainty principle and other hallmarks of quantum mechanics? Yes, in that an extended vortex R-phase has an inherent trade-off: if it's very well-defined in momentum (circulation), it's likely extended in space. In T-phase (localized knot), it has a definite position but then internally the circulation vector might be kind of not well-defined (maybe analogous to having internal spin uncertainty). The formal derivation of uncertainty is not trivial here, but one could attempt to map the equations of small excitations to a Schrödinger equation. Possibly, small oscillations of an unknotted loop are like a delocalized wavefunction, and when you restrict allowed modes by a boundary (like tying a knot imposes boundary conditions), that corresponds to localizing.

Another way canonical quantization enters is in the linear wave regime: if you linearize about a trivial vacuum, you'll find normal modes (phonons or photons) that can be quantized by usual means, yielding creation/annihilation operators for excitations of frequency ω . This standard route would recover that, e.g., the photon mode is quantized in units of energy $\hbar\omega$, presumably because the fluid's excitations carry momentum and energy in quantized lumps. However, importantly, these “quanta” are not fundamental random quanta; they are discrete because the underlying continuum has a countable set of modes.

SST's canonical framework likely defines a Poisson bracket or Hamiltonian formulation for the fluid (which is well known for classical fluids; there's a Hamiltonian description with the

Clebsch potentials or vorticity). Then imposing canonical commutation relations might not even be needed if the states are already discrete. But if one insisted on a field quantization, one could treat $B_{\mu\nu}$ and W_μ fields as quantum fields. The presence of stable solitons then implies a quantum Hilbert space that includes those soliton states. In that quantized field theory viewpoint, one would find a spectrum of bound states (like knotted flux tubes) possibly analogous to how in lattice QCD one finds glueball states or flux tube excitations. The big difference: those excitations are stable here due to topology, whereas in QCD flux tubes break (though QCD strings between quarks can't break because quarks are permanently attached; in SST maybe knot can't untie because of helicity conservation unless some energy is provided).

One consequence of the topological spectrum is that there might be relationships or selection rules for reactions. For example, if two vortex loops collide, to satisfy topological charge conservation, they might reconnect into new loops only if overall linking and knot numbers match. This could correlate to conservation laws like baryon number or lepton number. Indeed, baryon number might correspond to an invariant like “3 times linking number mod something” (since baryon is 3 quark knots linked, and maybe that linking number is conserved mod 3). If so, proton decay would require unraveling that composite link, which might be topologically forbidden without extremely high energy (so baryon number is effectively conserved unless extreme conditions). That fits with observed proton stability.

Another selection rule could involve spin: In SST, spin is related to how a knotted vortex rotates or twists. A trefoil has perhaps one unit of intrinsic angular momentum ($\hbar/2$). Two trefoils linked might produce an integer spin state, etc. Possibly it's akin to how you can combine twist states. The details are open, but presumably it's arranged to match spin-statistics: being knotted might impose fermionic behavior (trefoils are essentially twisted once, giving half-integer spin?), whereas unknotted loops are bosonic (photon is boson). If that holds, SST might naturally incorporate spin-statistics via topology: an unknotted loop has an even (or zero) twist, making it effectively a boson, whereas a knot has an odd half-twist requiring a 2π rotation to return to same configuration, yielding fermionic properties. If proven, that would be a major success (spin-statistics theorem from topology of fluid flow).

In summary, SST's quantization is realized in a **canonical ensemble of discrete vortex states**. The allowed states are countable and correspond to quantum states we observe. The continuous symmetries (like phase rotations, isospin rotations of internal fields) yield conserved quantities (like charge, isospin) which are quantized by topological winding numbers. The discrete symmetries and exclusion (spin-statistics) come from whether a vortex configuration can be continuously deformed into another or not (for identical fermions, swapping two knotted loops might yield a sign change if one has to be rotated 360° around the other to swap due to linking).

While many of these statements remain qualitative or conjectural within SST, the theory does produce at least one concrete outcome: a table of particle masses and quantum numbers that can be checked against reality. The authors of SST have done some of that calibration. If the predictions (for instance, predicting the neutron-proton mass difference, or the muon mass, etc.) are accurate, that is evidence in favor of the approach. Conversely, any glaring mismatch (like wrong mass ratios or missing a known particle) would be a problem. So far, the glimpses show encouraging correspondence within a few percent for leptons and nucleons.

Having laid out how SST conceptualizes quantization and obtains a topological spectrum of states, we now proceed to discuss concrete experiments that could validate or falsify various aspects of the theory.

1.8 Experimental Implications & Falsifiability

A physical theory as radical as Swirl-String Theory must be subjected to experimental tests. SST is constructed to reproduce known phenomena in its domain of validity, but it also makes distinct

predictions that depart from the Standard Model and general relativity. Here we highlight several key experimental implications of SST and how one might test them. These include: (I) quantized electromotive impulses from vortex events, (II) decay of quantum interference visibility due to finite vacuum vorticity, (III) structured photon propagation effects tied to knot chirality, and (IV) chiral-dependent time delays in photoemission (already discussed in Sec. 6). We also mention tests of emergent gravity (though those might be harder to isolate given gravity’s weakness). Crucially, each of these is *falsifiable*: if careful experiments fail to observe the predicted effect where SST predicts it should occur, the theory would be tightly constrained or refuted. Conversely, observing one of these novel effects would provide strong evidence for SST’s underlying fluid medium.

I. Quantized Flux Impulses from Reconnection Events: Perhaps the most striking test is the prediction that whenever a swirl string is created, destroyed, or changes its linkage through a loop, an electromotive force impulse of magnitude $\Phi_0 = h/2e$ (one flux quantum) is induced (Sec. 5). To test this, one doesn’t need to literally create vortex loops in vacuum (which might be difficult), but one can simulate analogous conditions in a controlled setup. One proposed platform is a type-II superconductor or a superfluid: these systems are well described by quantum fluid dynamics and support quantized vortices (in superconductors, magnetic flux lines; in superfluids, flow vortices). We can arrange a superconducting pickup loop (part of a SQUID circuit, for example) and then trigger single vortex events in its vicinity. For instance, using a superconducting film adjacent to the loop: by applying a small magnetic field, one can coax exactly one vortex to enter the film through the loop (once the film’s critical field is just exceeded). When that vortex enters, the loop should register a sharp voltage spike corresponding to one flux quantum crossing it. Indeed, in superconductivity, such a signal is expected by standard physics because a vortex carries one flux quantum Φ_0 and the SQUID is essentially measuring that (this is how SQUID magnetometers detect flux). However, here’s the key: SST expects that even in the absence of a pre-existing magnetic flux, a swirl vortex event (which in a superconductor is basically the same as a magnetic flux quantum event) produces that impulse. The real test is in a neutral superfluid (like a Bose–Einstein condensate of atoms) where quantized vortices carry circulation but not necessarily magnetic flux. If one could engineer a tiny loop of wire in a BEC and create or annihilate a single quantized vortex threading that loop, the moving vortex’s rotating frame should induce an electric field via the SST coupling. The predicted impulse is again Φ_0 in magnitude. This is exotic because a neutral BEC vortex should not couple to electromagnetism in the Standard Model, yet SST says the analog coupling in the true vacuum would produce an EMF. Since we can’t directly test vacuum, using a BEC with maybe a synthetic gauge field might mimic it. Alternatively, one could attempt to observe in heavy-ion collisions or other violent processes if any sudden $\nabla \times E$ fields are generated without charges present (though that’s messy).

The superconducting version of the experiment is simpler and essentially amounts to measuring known flux quanta, which is more a proof-of-concept. The BEC or other condensed matter analogs are challenging but conceivable. If such experiments find no evidence of impulses (especially in conditions where SST would predict them), it could imply either $G_\odot$ is very small or zero in vacuum. However, one might argue that if we only test analog systems, we have to be cautious: a BEC might not exactly replicate vacuum SST coupling unless the coupling is truly universal. Possibly, it might require coupling BEC vortices to actual electromagnetism via some mechanism (maybe using an atom with large polarizability such that vortex rotation produces an effective magnetic moment).

Nevertheless, the general expectation is that if swirl strings exist, they will reveal themselves through this quantized EM coupling if we are clever enough to isolate such events. The impulse detection via SQUID is at least technologically feasible, as SQUIDs can easily detect single flux quanta. The main challenge is controlling the single-vortex events in a clean way and ensuring

any measured signal isn't just from known inductive coupling (in the superconductor case, it is known and expected; in a neutral fluid it would be new).

II. Interference-Visibility Decay from Finite-Amplitude Swirl: Another prediction mentioned in the Lagrangian paper's abstract is that finite amplitude swirl flows can cause a decay in interference visibility. The idea is that if the vacuum (or experimental environment) isn't perfectly still but has some residual swirl excitations, these can decohere quantum superpositions over time. In simpler terms, if a particle is in a double-slit experiment and the vacuum medium around it has turbulence or oscillations, those fluid perturbations can carry information about which path the particle took, thus reducing interference contrast.

This effect would manifest as a slight reduction in fringe visibility that worsens with time or with certain conditions (like if vacuum fluctuations of swirl are induced by a strong field). The user text suggests a concrete scenario: interference-visibility decay driven by finite-amplitude swirl. Possibly one could test this by doing electron double-slit experiments in environments of varying conditions. For example, if one could "stir" the vacuum by placing a spinning heavy mass or an electromagnetic field that might drag the vacuum a tiny bit, does it reduce coherence of electrons? Standard theory would say no if it doesn't interact significantly. But SST might say yes, the swirl medium motion causes time dilation differences along the electron paths leading to phase jitter.

One way to check could be to use ultra-stable interferometers (like for neutrons or atoms) and see if rotation or motion of apparatus (or environment changes like temperature affecting vacuum modes) influences coherence beyond known effects. The challenge is to separate this from ordinary noise and vibrations. The effect might be subtle. The Lagrangian mention suggests that this is a concrete, parameter-free prediction: presumably, once we calibrate ρ_f and v_o , we can estimate how a given swirl amplitude affects decoherence rate. The abstract enumerated it as II after flux impulses. It implies that if we had a known amplitude swirl (like a vortex of certain circulation in vicinity), it will cause a quantifiable exponential decay of interference fringe contrast.

No such effect is known in standard physics except if the vortex physically interacts (like air flow or fields scattering the particles). But here it would be a "frictionless" fluid's effect.

This is reminiscent of random ether drift causing decoherence – an idea that was historically associated with stochastic gravity or something. But SST can put a number: maybe an amplitude of swirl corresponding to velocity v yields a decoherence time τ inversely related to v^2 (just guessing, since higher $v \rightarrow$ more time dilation differences $\rightarrow$ faster decoherence). If an experiment sees no such effect when v is known (like rotating the apparatus at 1 m/s tangential speed), it sets an upper bound on ρ_f or coupling.

Thus, interference experiments, particularly with massive particles or long coherence times, can constrain SST. For instance, interference of C_{60} fullerene molecules has been done. If SST's swirl medium had a density that caused any noticeable decoherence at those scales, it might have been seen. Possibly SST authors tuned parameters so that in normal conditions (no swirling flows), decoherence is negligible (consistent with observed long coherence lengths of, say, photons from distant stars, etc.). But if we artificially excite swirl flows, maybe by bounding a region with moving boundaries or intense laser fields, we could test if coherence is lost more rapidly than expected.

III. Skymionic Photon Textures (Polarization Patterns Indexed by Knot Chirality):

The Lagrangian abstract mentions "skymionic photon textures indexed by knot chirality." This suggests that photons (or electromagnetic fields) might acquire complex polarization or phase patterns (like skyrmions in their field configuration) that depend on the chirality of the source or medium. Possibly, a knotted vortex could imprint a specific swirling polarization pattern on photons it emits. For example, maybe radiation from a knotted electron (like synchrotron

radiation from a swirling electron vortex) has a distinctive polarization state.

An example might be: If an electron in SST is a trefoil knot, when it oscillates (emits a photon), the photon might carry not just spin-1 polarization, but a subtle topological phase (like optical vortices, where light can have phase singularities or orbital angular momentum). So SST might predict certain optical or gamma-ray emissions have spiral wavefront or polarization singularities that correlate with particle chirality. If one could measure the fine structure of light from certain atomic transitions or in scattering, one might find deviations like a small circular dichroism due to vacuum structure.

One accessible test: produce entangled photons from matter-antimatter annihilation or other processes and check if their polarization correlations reveal any pattern that could hint at underlying swirl structures. Or measure whether photon propagation in vacuum exhibits any self-rotation of polarization (birefringence) tied to global matter distribution. Standard QED says vacuum is not birefringent unless in fields (except in some quantum gravity ideas). SST might allow a scenario: if lots of swirl strings align (like a net helicity in cosmic structures), could that rotate polarization of traversing light (i.e. cosmic polarization rotation)? Possibly. That could be tested with astrophysical observations (light from distant galaxies through cosmic matter distribution – though that’s messy because cosmic dust etc also affects polarization).

Skyrmionic textures might also refer to the solutions of the B field equation where a photon is represented as a twisted field configuration. In that case, measuring them would require mapping the field in space, which is not easy for real photons except via interferometry. But one could attempt to simulate in metamaterials or other analogs.

This prediction is less straightforward to test in a tabletop experiment. It sounds like something that might come out in specialized circumstances or under careful analysis of polarization patterns in high-intensity laser experiments (where vacuum polarization might reveal structure). At least, it’s a potential distinctive feature to look for in photonic experiments.

IV. Attosecond Chirality-Sensitive Dynamics: We already covered this in Sec. 6: the photoionization delay experiments with chiral molecules [28]. Those results exist: enantiomers show opposite sign delays. Conventional theory can attribute those to subtle continuum scattering differences, but it’s somewhat surprising and has generated interest in “time-dependent chirality in photoionization.” SST offers an explanation and more importantly a prediction: if one could flip the swirl orientation of the electron artificially, it would flip the delay sign. We cannot flip an electron’s knot handedness easily (that’s like converting it to positron or something), but we can compare to maybe an antiparticle if similar experiments were possible (like photoionizing positronium or something with chiral environment? Hard to do). At least, the existence of this effect in molecules is a hint consistent with SST’s swirl clock idea.

To further test this aspect, one might try different molecules, different energies to see if the delay correlates with predicted path length differences. Or try to isolate whether it’s the vacuum property or molecular potential causing it. For example, in a purely achiral potential, SST would predict no forward-backward asymmetry (since swirl orientation doesn’t matter if environment symmetric). If one artificially introduces a chiral field (like a second laser with circular polarization in a known phase relation), does it amplify or invert the asymmetry? That could sort out how much is due to external fields vs internal swirl. At present, this line of investigation is already happening in attosecond science; SST provides a framework to interpret any anomalies or patterns in terms of a vacuum chirality effect.

Emergent Gravity Tests: While not enumerated in I–III, another big domain is testing the emergent gravity aspect. Some possible tests or constraints:

- Does G_{swirl} exactly equal G_N for all conditions? Could there be deviations in strong fields? Perhaps look at precision Cavendish experiments or inverse square law tests at short ranges:

if a swirl fluid has any slight compressibility or additional Yukawa-like behavior (via r_c core smoothing potential), it might deviate at sub-mm scales. Experimentally, gravity has been tested down to mm or sub-mm; no deviation found yet, so SST must align with that.

- In orbital motion or frame dragging tests (Gravity Probe B, LIGO detection of gravitational waves), would SST predict the same phenomena? If gravitational waves in GR correspond to fluid vorticity waves in SST, maybe the polarization or speed might differ. LIGO sees GR-consistent polarization and speed = c . If SST's analog can be shown to replicate those, okay; if not, it's trouble. It might be hard to conceive how a fluid medium exactly reproduces the two polarization states of GR gravitational waves (which are quadrupolar, spin-2). Possibly, coherent excitations of swirl (coupled B and W fields) could mimic it. If not, gravitational wave observations could challenge SST.
- Matter vs. antimatter free-fall: Since SST says both have positive mass and attract, it predicts antimatter falls down just like matter (which is consistent with current best tests, e.g. the ALPHA experiment at CERN recently measured that antihydrogen falls with g consistent with normal within $\sim 10\%$ or so uncertainty). If a future experiment found antimatter falls differently (say anti-H gravitating with different acceleration), that would break GR but also either break SST or force an adjustment (SST currently states matter/antimatter chirality doesn't affect gravitational energy, only swirl orientation flips sign but energy depends on v^2 so it's same). Right now, evidence suggests antimatter gravity is normal, which is consistent with SST.
- Tests of preferred frame: Since SST has an absolute frame (the swirl medium rest frame), there might be subtle violations of Lorentz invariance if one could detect motion relative to that medium. SST aims to be empirically Lorentz invariant for kinematics (like modern LETs are typically so exactly consistent that it's undetectable). But perhaps extremely sensitive clock comparisons or cosmic ray anisotropies might hint at an absolute rest frame. All 20th century Michelson–Morley type experiments put tight limits on any ether wind. For SST to survive, the swirl medium must either be exactly dragging along with Earth locally (which could be a boundary condition, like local comoving frame) or the coupling of matter to medium is such that no first-order effects appear. Possibly swirl medium flows around Earth (giving emergent gravity) also means Earth's frame is the rest frame at its location. Still, maybe cosmic frame (CMB rest frame) is the swirl medium on large scale. If so, tiny anisotropies in particle maximum speeds or something could appear. Modern tests (like comparing speed of light from different directions, or isotropy of inertia (Hughes–Drever experiments)) have not found any anisotropy beyond 10^{-18} level or so. That constrains any difference between swirl frame and Earth frame, implying swirl medium either perfectly drags or the coupling yields Lorentz invariance to high precision. Perhaps the preferred frame is only visible in second-order effects (like time dilation differences, which are small and usually gaugeable away).

SST authors probably argue it's completely consistent with Lorentz symmetry in all currently tested regimes (like any Lorentz Ether can be if the dynamics mimics SR exactly). But future more precise tests might push this further, e.g. searching for Lorentz violation in neutrino oscillations or such could indirectly challenge any underlying fixed frame.

Other Predictions: Beyond the ones enumerated, SST might have other signals:

- Vacuum Cherenkov radiation: In some ether theories, if particles move faster than medium excitations, they'd radiate. If medium excitations speed = c , then normal matter at subluminal speeds no effect. But high-energy cosmic rays nearing c – any difference? If c

is exactly the medium wave speed, no vacuum Cherenkov (consistent with observation, no photon emission from cosmic ray in vacuum).

- Variation of constants: If SST’s medium properties (like density ρ_f or Φ_0 coupling) vary with environment (temperature of vacuum, presence of fields), might fundamental constants vary slightly? E.g. speed of light or vacuum permittivity vary in gravitational potential (some theories like emergent gravity predict that). Could test with atomic clock comparisons at different gravitational potentials or electromagnetic fields. Current limits show if any variation, it’s extremely small.
- Direct detection of the medium: It’s a fluid, can we see any dissipative effect at extremely high frequencies or short scales? Perhaps at very short wavelengths, propagation might deviate from pure Maxwell (like dispersion or cutoff frequency if medium granular). E.g. if m_{eff} or r_c imposes a limit, maybe photons above certain frequency (like beyond some huge value) behave differently. Hard to test with present tech, but maybe cosmic gamma rays or neutrinos could reveal if vacuum not perfectly transparent or if there’s friction. Observations of high-energy photons from GRBs etc. show they arrive within microseconds regardless of energy differences, up to TeV scale, limiting dispersion. That suggests if SST medium has any frequency-dependent wavespeed, it’s extremely small up to that energy (which likely it is, since Φ_0 being small implies extremely stiff medium at those scales).

Finally, it’s worth noting falsifiability: If none of these predicted effects are observed, SST’s parameter space shrinks to a trivial corner (where effectively it becomes just an interpretation with no new effects). If one effect is observed (like quantized impulses with no conventional explanation, or an unexplained decoherence or polarization effect), it would be a breakthrough supporting SST or similar ideas.

The philosophy here is commendable: SST does not hide behind “we can always adjust to match experiments”—instead it lays out specific, parameter-fixed predictions I–III in the abstract, meaning the theory as formulated can be disproven.

In the next section, we will compare SST to other frameworks to highlight differences and similarities, before concluding.

1.9 Comparison with Existing Frameworks

Swirl–String Theory draws inspiration from numerous historical and contemporary frameworks, yet it differs in crucial ways. Here we compare SST with several related approaches: Kelvin’s vortex atom model, Madelung/Bohm hydrodynamic interpretations of quantum mechanics, analog gravity in fluids, string theory in high-energy physics, and the Standard Model plus General Relativity themselves. The goal is to situate SST in the landscape of theories, clarifying what it tries to improve or replace, and acknowledging where it must confront challenges already known in those frameworks.

Kelvin’s Vortex Atoms and Early Ether Theories: Lord Kelvin’s 19th-century idea proposed that atoms were knotted vortex rings in a luminiferous ether fluid [?]. This remarkably prescient model shares with SST the notion of matter as stable vortex knots and of conservation of topology (Kelvin noted that vortex loops in an ideal fluid cannot break, which would explain atomic stability). SST can be seen as a modern resurrection of Kelvin’s idea, now informed by quantum theory and relativity. However, Kelvin’s model lacked a mechanism for different elements (how to get chemical properties, periodic table from knots was speculative) and was eventually abandoned as the electron and nucleus were discovered as more fundamental entities. SST addresses one shortcoming by providing a mapping of knots to the known particle zoo rather than to chemical atoms, and introduces the swirl medium as a Lorentzian ether (to

respect relativity’s empirical success). Kelvin’s ether was a mechanical incompressible fluid filling space, and SST’s medium is essentially the same concept with a century and a half of additional physics layered on. Modern LETs were once considered superfluous since Einstein’s relativity can do without; SST counters that by giving the ether an active role (mass spectrum predictions, new coupling $G_\odot$, etc.). Unlike static classical ethers, SST’s condensate is a dynamic field that reproduces all of relativistic kinematics (much like Lorentz and Poincaré’s formulation but with a way to hide the absolute frame practically). In summary, SST embraces the spirit of Kelvin’s vortex atoms but places them in a comprehensive effective field theory context, aiming to overcome the criticisms that doomed older ether theories (lack of explanatory power and conflict with Michelson-Morley results).

Hydrodynamic Interpretations of Quantum Mechanics: Louis de Broglie and Erwin Madelung in the 1920s showed that the Schrödinger equation for a particle could be rewritten as fluid equations (Madelung’s hydrodynamic form) with a quantum potential term, suggesting perhaps quantum behavior could be understood as a form of fluid flow [?]. Later, David Bohm’s pilot-wave theory also invoked a real wave guiding particles, analogous to a fluid wave. More recently, experiments with oil droplets bouncing on vibrating fluid surfaces (Couder’s experiments) produced pilot-wave-like behavior, giving credence to a “quantum analog” fluid picture. SST shares with these the notion that a real fluid underlies quantum phenomena and that wave–particle duality can be visualized (in SST, the R-phase vs T-phase is analogous to pilot-wave guiding fluid vs localized droplet). The difference is that SST is far more ambitious: it’s not just an interpretation of quantum mechanics, it’s a full field theory claiming to produce gravity and other forces too. Madelung’s equations still needed an external quantum potential to match quantum mechanics; SST instead tries to derive the effective quantum potential (for example, the “quantum pressure” term in Madelung is replaced by fluid pressure effects from vortices in SST). Another difference: in pilot-wave theory, particles are pointlike with an associated guiding wave; in SST, the particle is the wave (when unknotted) or becomes localized as a soliton (knotted). There is no separate guiding wave vs particle, it’s the same entity in two regimes. This possibly avoids some conceptual pitfalls of pilot-wave (like what fundamentally is the guiding wave if not a physical field). On the flip side, pilot-wave theory is explicitly nonlocal (the guiding wave responds instantaneously in some formulations to enforce quantum correlations), whereas SST’s interactions are all local in the medium (though topologically they can produce nonlocal correlations akin to entanglement—this area likely needs work in SST: can it naturally produce entanglement and Bell inequality violations? That’s a challenge for any realist theory. If two particles are two distant vortex loops that were once linked or have correlated medium excitations, maybe it can emulate entanglement. But ensuring no superluminal signaling while having a medium is tricky. The authors likely claim the medium’s excitations are limited to c , so any correlation must be established at co-creation of entangled state and then just manifest as correlated outcomes without message sending. This is plausible if the state of the medium that connects them is nonlocally correlated but not transferring info faster than c . In any case, SST has to reproduce all quantum correlation results that standard QM does, or it will be falsified).

Overall, SST can be viewed as a synthesis of the hydrodynamic interpretation (for quantum phenomena) and topological soliton ideas (for particle stability), taking them to a unified extreme.

Analogue Gravity and Emergent Spacetime: In the early 2000s, a program of “analogue gravity” studied how various condensed matter systems (superfluids, Bose–Einstein condensates, electromagnetic waveguides, etc.) can reproduce phenomena of curved spacetime and black holes [?]. For instance, a moving fluid can have sound horizons analogous to event horizons (as in the famous Unruh sonic black hole). SST aligns with analogue gravity in that it too

asserts gravity is not fundamental but emergent from something more basic (in SST, fluid pressure gradients). The difference is analogue gravity usually doesn't claim the fluid is real spacetime—only a laboratory model to gain intuition—whereas SST says the fluid is the real thing. Some emergent gravity proponents (e.g. Ted Jacobson [18] or Erik Verlinde [7]) have suggested that Einstein's equations might be equations of state of some unknown microstructure of spacetime (like thermodynamics emerging from atoms). SST provides a candidate for that microstructure. It achieves emergent Newtonian gravity clearly (Sec. 4), and presumably could recover aspects of GR in some limit (though to fully get Einstein's field equations may require more in SST, e.g. the swirl medium's perturbations might mimic gravitational waves only in linear approximation but not all strong-field phenomena—this remains to be validated). Compared to other emergent gravity models (entropic gravity, quantum graphity, etc.), SST is a classical continuum theory, which is an advantage in clarity but potentially a drawback since quantizing gravity is replaced by just the fluid (so quantum gravity puzzles shift to fluid dynamics puzzles). But SST doesn't attempt to quantize spacetime, rather spacetime is a stage for fluid dynamics.

One should mention also the classical “fluid-mechanical model of ether” that was once considered for GR (e.g. theories by Dirk Jan Brouwer or others who tried to get Einstein equations from a fluid). Those didn't fully succeed. SST might succeed at least in the Newtonian regime by design. But if confronted with tests like gravitational light bending (which in GR comes from spacetime curvature), will SST get the right deflection? Possibly yes, if one considers that light in SST is a wave in the medium, passing through a low pressure region around a mass, it might bend akin to a refractive index gradient. In fact, one can derive light bending in a potential by treating space as having an effective refractive index $n(r)$; SST's swirl potential $V_{\text{SST}}(r)$ might induce an $n(r) > 1$ inside gravitational wells (because local light speed might be lower where swirl energy density is high, due to S_t factor). This could yield lensing. That is an avenue to check: does SST predict correct lensing (which is twice the Newtonian deflection)? If the medium slows light by the right amount (given $S_t = \sqrt{1 - v^2/c^2}$ for local time, an observer at infinity would see time (and thus light speed) reduced near mass, causing bending equivalently to GR's). If it only got half the deflection, then it fails. But likely the pressure effect yields effectively a curved space metric (Painlevé-Gullstrand form maybe). The authors should ensure that: emergent gravity in SST beyond Newtonian.

Nevertheless, in spirit, SST supports the emergent spacetime paradigm strongly: geometry is an epiphenomenon of fluid flows, not fundamental.

Superstring Theory and Topological Solitons: It's somewhat ironic that SST shares the word “string theory” but is almost opposite in philosophy to the mainstream string theory. Superstring theory posits fundamental one-dimensional strings vibrating in a fixed background or in a dynamically curved spacetime, aiming to unify all particles and gravity in a quantum framework. SST, by contrast, has strings as literal vortex loops in a classical fluid; they are composite/excitations, not fundamental Planck-scale strings. The only commonality is usage of topology: both have various winding and vibrational modes. But SST's knots vs superstring's oscillation modes correspond to quantum numbers differently. Also, SST operates at low energy/low velocity for emergent gravity, whereas superstring works at ultra-high energy to unify gravity with quantum. In some sense, SST is a competitor to high-energy unification: it claims you don't need to quantize gravity or introduce Planck-scale strings to get a unified description of forces—just use classical fluid and emergent fields. This is radical, and many physicists would be skeptical that a low-energy EFT (effectively) could hold at arbitrarily high energy. For instance, what happens at LHC energies or beyond? Does the fluid picture still hold, or do we see evidence of substructure? If SST is correct, maybe at some high energy or small scale, new phenomena (like vacuum turbulence or dispersion of propagation) might show up. If nothing shows up up to 10^{19} GeV, one might then accept the fluid as fundamental. But if new

stuff appears (like string theory predicts new particles, supersymmetry, extra dimensions), that could contradict SST if not accounted for. So far, LHC hasn't found new fundamental particles beyond Standard Model. That might not support SST per se, but it doesn't contradict it either. If one day an electron substructure is observed (like form factor deviations), one might tie that to size of a vortex core (r_c maybe 10^{-15} m yields some effect at momentum $1/r_c$ scale $\sim$ a few TeV maybe). It would be interesting if future accelerators see evidence of finite size of leptons or internal structure that could align with a vortex model.

Topological solitons have appeared in various quantum field contexts (e.g. Skyrmions for nucleons, hopfion solutions in certain nonlinear sigma models, etc.). SST is akin to a Skyrme-like model but in a physical fluid rather than an abstract field. It benefits from that because fluids naturally allow knotted vortex solutions that are hard to achieve in simpler field models (usually requiring higher order terms or special conditions). So SST's success might hinge on leveraging known stable vortex phenomena. On the flip side, control of those solutions mathematically is tough; one may rely on numerics or analogies rather than closed-form solutions. But if one can numerically demonstrate a knotted vortex is stable in the SST equations with exactly the right mass, that would be a big milestone.

Standard Model and GR: Finally, how does SST compare to just using the Standard Model (SM) and General Relativity (GR)? The SM+GR is extremely well-tested in their domains. SST must reproduce their successes where applicable: quantum electrodynamics, nuclear forces (if SST has analogs for weak and strong interactions via the emergent gauge fields W_μ^a perhaps representing SU(3) color etc.), and gravitational tests. The main motivation for SST is to provide a more unified and explanatory picture (e.g. why masses are what they are, why quantum is weird, etc.), and to be a *testable* alternative (predicting new physics rather than just adding epicycles). It also aims to be a realist interpretation, removing the mystery of wavefunction collapse, etc.

One advantage: SM has 19+ free parameters (masses, mixing angles). SST tries to cut that down drastically (maybe to a handful: ρ_f , v_o , some geometric factors in mass functional). If indeed SST can predict masses within few percent, that's something SM cannot do inherently (it just inputs them from experiment). Similarly, if SST can naturally explain why we see 3 generations of fermions (knot complexity up to some limit yields 3 sets? Or maybe more but higher knots unstable or correspond to very heavy states that decayed early?), that addresses a puzzle SM leaves open.

However, SST also introduces new constants like κ , m_{eff} (though those combine to physical ones anyway), $G_\circ$ (which turned out to be $h/2e$ by identification, interestingly not a new independent constant but linking known ones), and medium properties (like ρ_f which might be chosen such that $c = 1/\sqrt{\mu_0 \varepsilon_0}$ emerges correctly, so ρ_f is effectively fixed by that requirement if we identify $\rho_f = \varepsilon_0$ from earlier). Possibly ρ_f is determined by requiring $G_{\text{swirl}} = G_N$ as well in combination with v_o and r_c . The table in snippet [7] lines 4027-4036 says ρ_f is calibrated to 10^{-7} (some units) for dimensional consistency and acts like ε_0 , and ρ_m is defined via it. So ρ_f might be 10^{-7} kg/m³? If that's in SI, it's an extremely low density (like one-tenth of a microgram per cubic meter), which is so small that being undetectable physically is plausible. A medium that thin would be harder to detect than interstellar gas, and if it's non-viscous, it wouldn't slow planets or satellites (the classic ether wind drag arguments) because the coupling is only through gravity or EM subtle terms. Good.

Comparing how SM does things: forces by exchange particles vs SST: forces by fluid flow patterns. For electromagnetic force, SM says photons carry force, SST says two charged knots create pressure/velocity field patterns that push/pull (like two swirl rings in a smoke might push each other via induced flows—one might recall vortex ring interactions are interesting: coaxial vortex rings can attract or repel depending on orientation, akin to electromagnetic forces if you interpret something as "charge" or "current loops"). Possibly SST's emergent W_μ gauge field

yields something similar to non-Abelian gauge boson exchange effectively.

One major difference: SM's weak and strong forces are short-range (weak has massive bosons, strong confines). In SST, short-range could come from medium effects like vortex cores and alignment fields (like the director fields U_2, U_3 introduced perhaps give Yukawa-like contact terms for knots linking/unlinking). The Lagrangian abstract did mention "parity-violating and contact terms enabling vertices analogous to Yukawa and weak couplings." So they have put in some explicit terms to mimic weak force (parity violation means the medium can have a small helical term, or the swirl connection picks out handedness? Possibly their χ_h was a term to allow that but set to 0 for canonical theory, though if nonzero would break parity slightly, that could mimic weak interaction which is parity-violating). And "contact term for Yukawa coupling" suggests a term that directly couples three fields to simulate how quark knots might exchange something (maybe an instanton-like term when three vortex strands meet, akin to baryon-quark interactions?).

If SST can incorporate all these, it basically becomes an alternate formulation of SM (plus gravity) with fewer free parameters and a clearer ontology. But the risk is it could become as baroque as SM if many extra terms are needed to fit all details (like adding by hand a term to mimic each gauge interaction might seem just an elaborate rewrite rather than a true reduction of complexity). The authors likely aimed to derive as much as possible from minimal assumptions. Still, they did allow some extra terms (like θ term, λ_1, λ_2 constraints, etc. in Lagrangian—some might worry these are analogous to adding potential by hand or fine-tuning to get right results).

In any event, from a pragmatic perspective, SST will ultimately be judged by:

- Does it predict new phenomena (and do those occur)?
- Does it reduce conceptual problems (like unify concepts, solve hierarchy of masses, clarify quantum weirdness) in a satisfying way?

We have enumerated how it attempts those. If experiments show, e.g., the flux impulse or a specific mass prediction pans out, it gains credibility.

Conversely, if an effect like no sign of vacuum-induced decoherence or a clear violation of isotropy at some level is seen, that could challenge SST.

One comparative note: It's somewhat like a classical or hidden-variable theory underneath quantum—such approaches historically face Bell's theorem issues. SST's nonlocal connections (via the medium's topological links) might circumvent Bell locality assumptions by having the "hidden variables" (like relative phases or linking in the medium) established at creation of entangled pair. If done carefully, it might reproduce QM's predictions while maintaining an objective reality of sorts.

No analog of fermion exchange antisymmetry has been explicitly discussed here except noting that knots might encode spin-statistics. If SST fails to produce the Pauli exclusion principle naturally, it falls down (because we see that principle in matter structure). Possibly having an odd linking with the θ term could give minus signs under 360° rotation or exchange—this is subtle to evaluate.

Finally, one should mention: a challenge for any ether theory is stability and isotropy of the cosmic rest frame. The cosmic microwave background (CMB) provides a reference frame (the frame in which CMB is isotropic). Earth moves at ~ 370 km/s relative to CMB. If swirl medium is at rest with CMB, Earth is moving a bit relative to it. That speed is $10^{-3}c$. Michelson–Morley experiments limit anisotropy in light speed to parts in 10^{-15} , which would correspond to any effect of order $(v/c)^2 \approx 10^{-6}$ suppressed by 10^{-9} more beyond naive expectation. Possibly SST says a uniform medium flow has no effect on experiments because only swirling (rotational) motion matters, or that electromagnetic wave equations are Lorentz invariant up to second order automatically (like in Lorentz ether, you can't detect first-order because matter contracts etc., only second order which might be too small to measure at 370 km/s with old experiments, but

modern ones might catch second order). However, modern experiments like LIGO or atomic clocks might be able to see second-order anisotropy. The fact none is reported suggests if the medium exists, either it’s exactly comoving with Earth locally (maybe entrained by gravitational potential—some ether theorists proposed gravitational entrainment of ether), or the effect is beyond current sensitivity.

In summary, compared to SM+GR, SST is more cumbersome in that it introduces a physical medium and complicated fluid dynamics, but it’s rewarding if it explains things SM+GR leave arbitrary (like mass values, quantum measurement). It’s up against very tight experimental scrutiny though.

1.10 Conclusion & Outlook

We have presented **Swirl–String Theory (SST)** as a comprehensive fluid-based reformulation of fundamental physics. In this model, all of particle physics and gravity emerge from the dynamics of a single classical medium permeating space. By treating particles as quantized vortex loops (*swirl strings*) in an incompressible condensate, SST provides intuitive accounts of formerly mysterious phenomena: quantum duality arises from knotted vs. unknotted phases of these loops, inertia and gravity stem from fluid flows and pressure, and discrete quantum numbers correspond to topological invariants. We have shown how a Lagrangian framework can be constructed for this picture, embedding the postulates into a field theory that reproduces known physics in appropriate limits.

Several key results stand out:

- **Emergent Newtonian Gravity:** A pair of chiral vortex rings attract each other with an inverse-square law, and by calibrating the circulation quantum κ and core size r_c , the effective gravitational constant matches G_N . This offers a concrete fluid mechanism underlying gravity: low pressure zones created by swirling matter pull masses together, in flat space but mimicking curved-space gravity to leading order (Appendix C).
- **Swirl-Induced Time Dilation:** Clocks tied to swirling motion tick slower by the factor $S_t = \sqrt{1 - v^2/c^2}$, exactly as in special relativity. We derived this “swirl clock” effect from kinetic arguments and the medium’s Lorentz-like symmetry (Appendix B). It unifies kinematic time dilation and gravitational redshift under one fluid phenomenon: both are due to local swirl energy content.
- **Modified Faraday’s Law and Flux Quanta:** Perhaps the most distinctive prediction, we derived that a changing density of swirl strings induces an electric field curl, adding a term $G_\circ \partial_t \varrho_\circ$ to Faraday’s law (Eq. 1.5.1). The coupling $G_\circ$ is identified with $h/2e$, linking this new effect to the quantum of magnetic flux. A single vortex crossing a loop yields a sharp EMF impulse of one flux quantum. This result (derived in Appendix D) points to experimental tests with SQUIDS and other systems.
- **Topological Spectrum of Matter:** The allowed stable configurations of swirl strings correspond one-to-one with observed particle species (as summarized in Fig. 1). We highlighted how leptons, quarks, and photons map to specific knot types, and how composite knots model baryons and nuclei. Using a calibrated topological mass functional, SST can predict particle masses within a few percent in favorable cases, a level of quantitative prediction unusual for a beyond-standard-model theory.
- **Chirality-Dependent Dynamics:** SST attributes parity violation and certain time asymmetries to the handedness of the underlying vortices. We discussed how the theory accounts for attosecond-scale differences in photoionization times for opposite enantiomers (as observed in experiments [28]) by invoking the concept of a swirl-clock orientation.

Matter and antimatter are mirror-swirl states, which in SST have identical bulk properties except for how they interact with chiral environments or fields.

- **Unification and Realism:** With a single substrate and a unified set of equations, SST strives to unify interactions that in the Standard Model are disparate. The emergent W_μ^a gauge field in our Lagrangian hints at a common origin for gauge forces in coarse-grained vorticity structures. Moreover, SST is deterministic and realist at its core: the wavefunction’s mysteries are traded for concrete fluid behavior. While this is conceptually satisfying, it forces confronting issues like nonlocality and Bell’s theorem in novel ways (e.g. through pre-existing topological linkages). We have not fully delved into those, but any future elaboration of SST must show how entanglement and spin-statistics arise naturally.

Looking forward, there are several avenues for further research and development of SST:

1. **Relativistic and Strong-Field Extensions:** Thus far, SST recovers special relativistic kinematics and Newtonian gravity. The next step is to demonstrate (or refute) that SST can produce the full phenomenology of general relativity in appropriate regimes. This means exploring whether the fluid equations (perhaps with the $B_{\mu\nu}$ field) contain an analog of Einstein’s equations or at least the post-Newtonian corrections (light bending, gravitational time dilation, frame dragging, gravitational waves). The formal analogy between fluid flow equations and curved-space metrics (e.g. the Painlevé-Gullstrand form of Schwarzschild metric corresponds to a draining flow) should be investigated. A concrete goal would be to derive the effective spacetime metric seen by perturbations in the medium and check if it satisfies Einstein’s field equations with an emergent stress tensor. If discrepancies are found (for instance, if SST predicts the wrong amount of light bending), modifications to the theory or identification of specific experimental conditions under which it holds would be necessary.
2. **Stability and Topology of Solutions:** We relied on qualitative arguments about knotted vortices. Rigorous work (likely numerical simulations) is needed to verify that a knotted vortex in an incompressible, inviscid fluid with the given additional terms can remain stable indefinitely. The inclusion of the helicity term $W\tilde{W}$ is presumably to stabilize knots against unraveling or shrinking. Simulations of vortex ring collisions, knot tying/unknotted in fluids, and their time dilation effects would provide tangible evidence for SST’s mechanisms. It would also help refine the particle-knot mapping: e.g. confirming that a trefoil loop carries one unit of charge and spin- $\frac{1}{2}$ in the theory, whereas an unknot loop carries spin-1, etc.
3. **Quantum Corrections and the Correspondence Limit:** While SST is a classical theory at heart, it must reproduce quantum mechanics as observed. That means demonstrating that small fluctuations (the phonon/torsion modes) on top of a background can be quantized in the usual way. One path is to quantize the effective field theory given by $\mathcal{L}$, treating $B_{\mu\nu}, W_\mu, \Phi$ fields quantum mechanically. This raises the possibility of new quanta: for example, quantized excitations of the B -field are like emergent photons (which we already consider), quantized W -field excitations might act like emergent gauge bosons (gluons, $W^\pm$, etc.), and quantized Φ excitations could be a new scalar (maybe related to the Higgs scale). SST might then incorporate a connection to the Higgs mechanism via Φ fluctuations (though Φ was not the SM Higgs, it could mix with it or explain electroweak symmetry breaking as a property of the medium’s anisotropy). Ensuring that this quantization leads to the correct commutation relations and uncertainty principles will either validate SST’s viability or reveal tension. The theory’s success will hinge on showing that no contradiction arises between having a fundamentally classical substrate and the observed irreducible quantum randomness—likely the randomness enters through

sensitivity to initial conditions of the fluid (chaos amplifying microscopic fluctuations). This is reminiscent of Bohmian mechanics, but here the complexity of the fluid could lead to effectively unpredictable outcomes, matching the Born rule.

4. **Integrating the Weak and Strong Interactions:** We have largely focused on gravity and electromagnetism. The outline of how weak and strong forces might emerge (via swirl gauge field W_μ^a and internal director fields U_2, U_3) has been hinted at but not fleshed out here. That is a major task: one must show that knotted vortices interacting in the fluid give rise to, say, $SU(2)_L$ gauge fields that mediate beta decay, and an $SU(3)$ color interaction that confines the three quark-knots into a baryon. Possibly, the linking of three vortices (baryon) is stabilized by a shared topological flux analogous to a glued color flux tube—this might mirror the Y-shaped color flux tubes in QCD. The parity-violating term in $\mathcal{L}$ could be exploited to make one chirality of knot interact differently (thus replicating the chiral nature of the weak force). Achieving this in a controlled fluid model is challenging; it may require adding dissipative or symmetry-breaking terms that are phenomenological. If too much has to be added by hand, SST’s elegance would suffer. But if it naturally yields these interactions from, e.g., the coupling of vorticity to internal orientation fields, that would be a profound unification. One concrete step: derive the analog of the electroweak mixing angle or coupling constants from medium properties, and check if one can get values close to the observed g, g' , etc. At least qualitatively, an explanation of why the weak force is weak (massive W, Z bosons) might come from twist stiffness or a cutoff length in the medium for certain excitations (the “swirl Coulomb” potential has a core r_c which could correspond to a Compton wavelength of heavy bosons).
5. **Experimental Searches for Predicted Effects:** On the experimental side, the list of tests in Sec. 8 serves as a guide for near-term investigations. In particular, verifying the quantized EM impulse prediction would be a landmark result. Setting up the described SQUID experiment with a single vortex injection, or its superfluid analog, should be pursued. Additionally, experiments in high vacuum with atom interferometers could check for any unusual decoherence when subjected to rotating frame or in proximity to large-scale vortex flows (perhaps using superconducting magnets to simulate swirl fields). High-precision optical polarization experiments could attempt to detect any self-rotation or birefringence in vacuum aligned with the presence of swirling matter (for example, light passing near a rotating superfluid might experience a tiny phase shift if SST coupling exists). Although challenging, these experiments benefit from modern techniques in quantum sensing and could reach the sensitivity needed to detect extremely subtle effects.
6. **Astrophysical and Cosmological Implications:** If the swirl medium is real, it might have consequences in astrophysics. For example, could astrophysical jets or vortex structures be manifestations of swirling vacuum? Or, could early-universe baryogenesis relate to an imbalance in swirl chirality (providing a tangible mechanism for matter–antimatter asymmetry by favoring one chirality of knots)? The “triadic closure” Axiom 0 which enforced something like a mod 2π sum for three circulations hints at a built-in baryon number conservation or asymmetry. It would be worthwhile to examine if SST can accommodate cosmic inflation or dark energy as properties of the medium (maybe Φ field variations or global rotation of the condensate). Similarly, dark matter might find an explanation as certain knot states that are stable but not interacting (like the figure-8 knot dark state we mentioned). One could predict, for instance, what fraction of medium excitations are in dark-sector knots given some conditions, and whether they could clump gravitationally—tying into structure formation.

In closing, Swirl–String Theory offers a daring and grand unification of concepts, converting the metaphors of fluid dynamics into literal physical mechanisms for the universe’s workings.

It brings the solidity of classical intuition to the ethereal realm of quantum fields and curved spacetime. Whether nature indeed operates like a cosmic superfluid remains to be seen. The theory does not shy from empirical risk: the coming years could see it triumph through confirmed predictions or face falsification. Regardless, the exploration of SST enriches the dialogue between disciplines—fluid mechanics, topology, quantum theory, and gravitation—illustrating how progress often re-emerges from reimagining old ideas (like Kelvin’s vortex) in the light of new knowledge. If nothing else, SST provides a fresh conceptual playground, one where one can picture the invisible: watching the fabric of reality swirl, knot, and unknot to craft the tapestry of particles and forces we observe.

Appendix A: Lagrangian Density and Master Equations

For completeness, we provide the full form of the SST Lagrangian density and some key field equations derived from it. As given in the main text (Eq. 1.5), the Lagrangian can be written as:

$$\begin{aligned}\mathcal{L}_{\text{SST}} = & -\frac{\kappa_\omega}{4}W_{\mu\nu}^aW^{a,\mu\nu} + \frac{\kappa_B}{12}H_{\mu\nu\rho}H^{\mu\nu\rho} + \frac{1}{2}(\partial_\mu\Phi)(\partial^\mu\Phi) - V(\Phi) \\ & + \frac{\theta}{4}W_{\mu\nu}^a\tilde{W}^{a,\mu\nu} + \lambda_1(u_\mu u^\mu + 1) + \lambda_2(\nabla_\mu u^\mu) \\ & + \sum_K \bar{\Psi}_K \left(i\gamma^\mu D_\mu - m_K^{(\text{sol})} \right) \Psi_K\end{aligned}\quad (12)$$

where the symbols have the meanings described earlier (briefly: $W_{\mu\nu}^a$ is the field strength of the swirl gauge field W_μ^a , $H_{\mu\nu\rho} = \partial_{[\mu}B_{\nu\rho]}$ is the 3-form field strength of $B_{\mu\nu}$, Φ is the condensate modulus, u_μ is the unit timelike 4-velocity of the medium with Lagrange multipliers $\lambda_{1,2}$ enforcing its normalization and incompressibility, θ is the coefficient of the topological $W\tilde{W}$ term, and Ψ_K are spinor fields representing quasiparticles for each knot type K with soliton mass $m_K^{(\text{sol})}$).

From this $\mathcal{L}$, one can derive the Euler–Lagrange equations for each field. We list a few central equations (without damping or external forces, all in SI units or with $c = 1$ in some cases):

- *Condensate 4-velocity (u^μ) equations:* Varying λ_1 and λ_2 yields the constraints

$$u_\mu u^\mu = -1, \quad \nabla_\mu u^\mu = 0 \quad (13)$$

ensuring u^μ is a divergence-free unit vector field (effectively a statement of constant density and no expansion in the incompressible limit). Varying u^μ itself (taking into account these constraints and the coupling to other fields via $h_{\mu\nu}$) leads to a Navier–Stokes-like equation. In the simplest form (neglecting shear viscosity, pressure gradients beyond those coming from Φ), it can be written as:

$$(u \cdot \nabla)u^\mu + \nabla^\mu \Phi_{\text{eff}} + \frac{1}{\rho_f} u_\nu H^{\nu\mu\sigma} u^\lambda H_{\lambda\sigma\alpha} u^\alpha = 0 \quad (14)$$

which is a generalized Euler equation for the flow. Here Φ_{eff} is an effective potential combining the pressure from fluid motion and any condensate potential $V(\Phi)$ contributions, and the last term represents the influence of vortex fields $H_{\mu\nu\rho}$ on the flow (it comes from the H^2 term and ensures Kelvin’s circulation theorem in the absence of other forces). Equation (14) in a non-relativistic approximation reduces to $\partial_t \mathbf{v} + (\mathbf{v} \cdot \nabla) \mathbf{v} = -\nabla \tilde{h}$ (where $\tilde{h}$ is enthalpy per unit mass), consistent with classical incompressible flow equations.

- *Swirl 2-form ($B_{\mu\nu}$) field equation:* Varying $B_{\mu\nu}$ yields (using $H^{\mu\nu\rho} = \kappa_B^{-1} \partial \mathcal{L} / \partial (\partial_\rho B_{\mu\nu})$):

$$\partial_\rho H^{\mu\nu\rho} = J^{\mu\nu} \quad (15)$$

where $J^{\mu\nu}$ is the source 2-form from any coupling to Ψ_K (if the knotted particles couple to B as string sources). In the absence of explicit string sources (treating the fluid vorticity as mostly continuous), this is simply $\partial_\rho H^{\mu\nu\rho} = 0$, which is analogous to vorticity conservation (Kelvin's theorem in differential form). In 3-vector language, one part of this equation is $\partial_t \boldsymbol{\omega} = \nabla \times (\mathbf{v} \times \boldsymbol{\omega})$ (the vorticity transport equation for an inviscid fluid). This shows the medium supports vortex lines moving with the flow, and no vorticity is created or destroyed except where string sources (the Ψ_K terms) might enter when a swirl string is explicitly introduced or ends (in our theory physical strings are closed loops so effectively $J^{\mu\nu} = 0$ everywhere; an open string would need an end which doesn't exist for a vortex in a true incompressible fluid).

- *Swirl gauge field (W_μ^a) equations:* Varying W_μ^a and using $W^{a\mu\nu} = \partial\mathcal{L}/\partial(\partial_\nu W_\mu^a) = -\kappa_\omega W^{a\mu\nu} + \theta \tilde{W}^{a\mu\nu}$ yields a Yang–Mills type equation with a topological current:

$$D_\nu W^{a,\mu\nu} + \frac{\theta}{2\kappa_\omega} \epsilon^{\mu\nu\rho\sigma} W_{\nu\rho}^a \partial_\sigma(\cdot) = J^{a,\mu} \quad (16)$$

where D_ν is the covariant derivative for the gauge group and $J^{a,\mu}$ is the Noether current from Ψ_K spinors (e.g., $\bar{\Psi}\gamma^\mu T^a \Psi$). The second term (with θ) is identically zero when taking divergence (Bianchi identity), but its variation yields the conserved helicity charge. If $\theta \neq 0$, the theory admits solutions with nonzero $W\tilde{W}$ that saturate certain bounds (resembling instantons; in our context they reflect knottedness conservation). Ignoring θ for the equation of motion, Eq. (16) reduces to $D_\nu W^{a,\mu\nu} = J^{a,\mu}$, a standard Yang–Mills equation stating that coarse-grained swirl vorticity ($W_{\mu\nu}$) is sourced by knotted spinor currents. In the Abelian case (analogue of Maxwell), this splits into a Coulomb-like law and Faraday's law. For example, setting a to correspond to electromagnetism and ignoring non-Abelian parts, the spatial components yield $\nabla \cdot \mathbf{E} = \rho_{\text{el}} + \rho_\circ$, where we have included both electric charge density ρ_{el} from conventional charges and an effective contribution $\rho_\circ$ from the time-varying swirl density via $G_\circ$ coupling (this comes from the modified Gauss's law including the new induction term). Likewise, the curl of the electric field gives Eq. 1.5.1 in the main text.

In short, Eq. (16) contains within it the modified Maxwell equations:

$$\begin{aligned} \nabla \cdot \mathbf{E} &= \frac{\rho_e}{\epsilon_0} + G_\circ \varrho_\circ \\ \nabla \times \mathbf{B} - \frac{1}{c^2} \partial_t \mathbf{E} &= \mu_0 \mathbf{J}_e + G_\circ \nabla \times \mathbf{M}_\circ \\ \nabla \cdot \mathbf{B} &= 0 \\ \nabla \times \mathbf{E} + \partial_t \mathbf{B} &= G_\circ \partial_t \mathcal{B}_\circ \end{aligned}$$

where ρ_e , $\mathbf{J}_e$ are electric charge and current of conventional sources, and $\varrho_\circ$, $\mathbf{M}_\circ$, $\mathcal{B}_\circ$ represent effective sources due to the swirl medium (with $G_\circ$ playing the role of coupling constant). In static conditions without vortices, these reduce to the usual Maxwell equations.

- *Condensate modulus (Φ) equation:* Varying Φ yields a Klein–Gordon-type equation for the condensate stiffness:

$$\nabla^\mu \nabla_\mu \Phi + \frac{\partial V}{\partial \Phi} = -\frac{1}{2} \frac{\partial \ln(\rho_f)}{\partial \Phi} H_{\mu\nu\rho} H^{\mu\nu\rho} + \dots \quad (17)$$

The right-hand side could include contributions from how ρ_f (the effective density) might depend on Φ . In many formulations, one would assume ρ_f is constant and then the equation is simply $\nabla^2\Phi - V'(\Phi) = 0$, which has the homogeneous solution $\Phi = \Phi_0$ as a stable vacuum. Small excitations $\delta\Phi$ would propagate with velocity determined by $V''(\Phi_0)$, potentially representing a massive scalar mode (which could be very high mass if the medium is very stiff, hence not easily excited except perhaps in extreme events like the Big Bang or neutron stars).

- *Soliton spinor (Ψ_K) equations:* Each quasiparticle field satisfies a Dirac equation minimally coupled to W_μ :

$$i\gamma^\mu D_\mu \Psi_K - m_K^{(\text{sol})} \Psi_K = 0 \quad (18)$$

Here $D_\mu = \partial_\mu + ig_{\text{sw}} W_\mu^a T^a$ as before. This equation is an effective one emerging after one has integrated out the detailed vortex structure, so $m_K^{(\text{sol})}$ already encodes the energy of the knotted vortex. Importantly, any solution Ψ_K must be supported by an underlying vortex configuration in the B and W fields. In practice, one might solve the classical equations for a knotted vortex (neglecting Ψ), then quantize small fluctuations around it to produce Ψ as a collective coordinate. The Dirac equation ensures these spinor solutions transform correctly under Lorentz symmetry and have the right statistics (assuming the usual anticommutation for spinor fields).

What is notable is that in the limit of a weakly coupled gauge field ($g_{\text{sw}} \rightarrow 0$ or medium very symmetric), Eq. 18 decouples and the spinor is free, revealing the physical interpretation: a free Ψ_K corresponds to a free knotted vortex propagating without interacting beyond carrying its rest mass.

This Appendix has laid out the formal underpinnings of SST's equations. While complex, they show consistency with known forms: Kelvin's circulation theorem, Navier–Stokes continuity, Yang–Mills equations, etc., are all embedded here. Many simplifying assumptions were used (inviscid fluid, symmetrical condensate, etc.); relaxing those will enrich the phenomenology (e.g. small viscosity could correspond to finite lifetimes of certain states or CPT violation due to small helicity-violation χ_h).

Finally, to connect with Table III in the main text (traceability of concepts), we mention some “Master” dimensionless parameters that appear in these equations and calibrations:

$$\text{Fr} = \frac{v^2}{gL}, \quad \text{Re} = \frac{\rho_f v L}{\eta}, \quad \frac{\rho_m}{\rho_f} = \frac{v^2}{c^2}, \quad G_{\text{swirl}} = \frac{\kappa^2 \rho_f}{8\pi^2 c^2} \quad (19)$$

where Fr is a Froude number (ratio of flow inertia to gravity, relevant in analog gravity contexts), Re is Reynolds number (which in our idealized theory is infinite due to $\eta = 0$ viscosity), ρ_m/ρ_f ratio relates swirl energy to rest energy density, and G_{swirl} is an example of how medium constants determine gravitational coupling. Plugging numbers (as done in the main text's Table III): if $\rho_f \sim 10^{-7} \text{ kg/m}^3$, κ corresponds to $\hbar/m_{\text{eff}}$ with m_{eff} chosen around some Planck-scale or particle-scale value, one can indeed get $G_{\text{swirl}} \approx 6.67 \times 10^{-11} \text{ SI}$. Thus, the abstract equations here can meet empirical scales.

Appendix B: Derivation of Swirl-Clock Time Dilation

In this appendix, we derive Eq. (1.2) of the main text from the dynamics of the swirl medium. The result, $dt_{\text{local}} = \sqrt{1 - v^2/c^2} dt_\infty$, can be obtained in multiple ways: (1) by analyzing the metric induced by moving with the fluid's 4-velocity u^μ , (2) by using an energy argument (equating kinetic and total energy for a comoving clock), or (3) by analogy to special relativity

with the existence of a preferred frame. We provide method (1) here, which is geometric and aligns with modern Lorentz ether interpretations.

Consider an observer (clock) attached to a small swirl string segment moving with velocity v relative to the rest frame of the medium. In the rest frame of the medium (the “lab frame” or chronos frame), the metric is just flat Minkowski: $ds^2 = -c^2 dt^2 + d\mathbf{x}^2$. The fluid provides a preferred slicing of spacetime into absolute time $T(x^\mu)$ surfaces (i.e. $T = \text{constant}$ defines a simultaneity in the medium’s rest frame). The clock moves with 4-velocity U_{clock}^μ . Because the clock is comoving with the local fluid, U_{clock}^μ is aligned with the fluid’s u^μ at that location (except perhaps small residual motions if not exactly stuck to a vortex line; here we assume it’s attached to the vortex).

Therefore, $U_{\text{clock}}^\mu = u^\mu$ (clock moves with the fluid element), which implies

$$U_{\text{clock}}^0 = \gamma, \quad U_{\text{clock}}^i = \gamma\beta^i, \quad (20)$$

where $\beta^i = v^i/c$ and $\gamma = 1/\sqrt{1 - \beta^2}$, in the coordinates of the lab frame. This is just a statement that if the fluid element moves at velocity $\mathbf{v}$, then the time-component of its 4-velocity is γ .

Now the proper time increment $d\tau$ measured by the clock between two events (e.g. ticks) on its worldline is given by the metric:

$$d\tau^2 = -\frac{ds^2}{c^2} = dt^2 - \frac{d\mathbf{x}^2}{c^2}, \quad (21)$$

for those events (assuming they are closely separated). If in time dt the clock moves $d\mathbf{x} = \mathbf{v}dt$, then

$$d\tau^2 = dt^2 \left(1 - \frac{v^2}{c^2}\right),$$

so $d\tau = \sqrt{1 - \frac{v^2}{c^2}} dt$. By identification, $dt_{\text{local}} = d\tau$ (proper time of clock) and $dt_\infty = dt$ (lab frame time, essentially the time coordinate T at infinity far from any swirl), yielding

$$dt_{\text{local}} = \sqrt{1 - \frac{v^2}{c^2}} dt_\infty. \quad (22)$$

This matches Eq. (1.2).

The interpretation is straightforward: the swirl medium’s rest frame is the frame of “aether” time t_∞ (which we called chronos time). A clock moving at speed v relative to this frame will be slowed by the usual Lorentz factor. What ensures consistency in SST (so that this is not in conflict with all relativity experiments) is that all physical processes including light propagation are also governed by the same metric and thus also experience the same time dilation and length contraction. For instance, light still moves at speed c in the medium (by construction of ρ_f as analogous to ε_0), and the laws of physics in a moving frame of the medium take the Lorentz-transformed form. This is essentially the Lorentz invariance of the equations built into our Lagrangian, albeit with a hidden absolute frame.

To connect this derivation to a more fluid-mechanical perspective: one can derive time dilation from Bernoulli’s principle. In steady flow (no explicit time dependence in lab frame), the quantity $E = \frac{1}{2}v^2 + w + \Phi_g$ is constant along flow lines (where w is specific enthalpy and Φ_g gravitational potential). In our case, w might include rest energy c^2 as well since we consider relativistic regime. If a fluid element’s total energy is constant, then if kinetic energy $\frac{1}{2}v^2$ increases, the internal energy (which influences clock rate) must decrease. A rough argument is that a clock’s tick rate is proportional to its internal energy transitions (say an atomic transition frequency). If that internal energy is effectively lower by a factor γ when moving (time dilation corresponds to lower frequency), then indeed $\nu_{\text{moving}} = \nu_{\text{rest}}/\gamma$. This is an informal reasoning consistent with the exact result we derived above.

Another approach uses the so-called “chronometric invariants” formalism (used by Lorentz ether theories [?]): define a global time t and a spatial metric h_{ij} on simultaneity surfaces. The proper time for an observer with velocity v^i relative to the ether is $d\tau = \sqrt{-g_{\mu\nu}dx^\mu dx^\nu}/c = \sqrt{dt^2 - h_{ij}dx^i dx^j}/c$. If $h_{ij} = \delta_{ij}$ (flat space) and $dx^i = v^i dt$, it reduces to the above. In curved or flow-distorted cases, one might get additional gravitational redshift factors (e.g. if Φ_g present, $d\tau = \sqrt{(1 + 2\Phi_g/c^2) - v^2/c^2} dt$ to first order). SST indeed predicts gravitational potential affects time similarly, since Φ_g arises from swirl flows as well.

In conclusion, the swirl-clock time dilation factor is a built-in feature of any Lorentz-covariant theory with a preferred frame and consistent dynamics. SST passes this test by construction. Notably, if one were to detect a deviation from $S_t = \sqrt{1 - v^2/c^2}$ at high v for moving clocks, that would falsify Lorentz invariance. Experiments (particle lifetime measurements, fast ion clocks) confirm the γ factor to high precision, thus also confirming that if a fluid ether exists, it must exactly implement this time dilation. Our derivation shows how SST does so: by encoding motion effects in the metric $g_{\mu\nu}$ and ensuring all matter fields (Ψ_K , etc.) couple to that metric in the standard way (through the minimal coupling and $h_{\mu\nu}$ projector).

Appendix C: Emergent Newtonian Gravity from Swirl Pressure

Here we derive the inverse-square law attraction between two swirl strings, focusing on the “Hydrogen-Gravity mechanism” described in Sec. 4. We work in the regime of non-relativistic flow ($v \ll c$, incompressible fluid) and show how a pair of vortex loops co-axial along an axis produce a force analogous to Newtonian gravity.

C1. Pressure field of a single vortex loop

Consider an isolated swirl string that, for simplicity, we treat as a thin straight vortex filament of circulation κ aligned with the z -axis (an infinite line, which can approximate a large loop section or a vortex through a torus representing a bound hydrogen-like system). In cylindrical coordinates (r, φ, z) , the flow is azimuthal: $\mathbf{v}(r) = v_\varphi(r)\hat{\varphi}$, with

$$v_\varphi(r) = \frac{\kappa}{2\pi r} f(r), \quad (23)$$

where $f(r)$ is some profile function that accounts for the vortex core (e.g. $f(r) = 0$ inside core of radius a , tend to 1 at $r \gg a$). For an ideal thin vortex, $f(r) = 1$ for $r > a$ and effectively v diverges as $1/r$ outside core, then drops to 0 at $r = 0$ (physical core size provides cutoff).

Under these conditions, the fluid’s Bernoulli equation (from momentum conservation) in steady state reads:

$$\frac{1}{2}v^2 + \frac{p}{\rho_f} = \text{const (along a streamline)}. \quad (24)$$

Far from the vortex ($r \rightarrow \infty$), we can set $p \rightarrow p_\infty$ (ambient pressure) and $v \rightarrow 0$, so the constant is p_∞/ρ_f . Thus at radius r ,

$$\frac{p(r) - p_\infty}{\rho_f} = -\frac{1}{2}v_\varphi(r)^2, \quad (25)$$

Plugging Eq. (23) for v_φ (and assuming r outside the core so $f(r) \approx 1$), we obtain

$$\Delta p(r) \equiv p(r) - p_\infty = -\frac{\rho_f \kappa^2}{8\pi^2 r^2}, \quad (26)$$

This is the pressure deficit as a function of radius. It falls off as $1/r^2$. Note that this is structurally similar to a Coulomb or gravitational potential profile (since in a static field,

$\nabla^2\Phi \propto \delta(\mathbf{r})$ leads to $\Phi \propto 1/r$ in 3D, and field $\propto 1/r^2$). Indeed, one can define an effective potential for this vortex:

$$\Phi_{\text{swirl}}(r) = \int^r \frac{\Delta p(r')}{\rho_f} dr' = -\frac{\kappa^2}{8\pi^2 r}, \quad (27)$$

where we integrated assuming core cutoff at small r . The form $-A/r$ emerges, with $A = \kappa^2/(8\pi^2)$.

This $\Phi_{\text{swirl}}(r)$ plays the role of a gravitational potential in SST. In particular, one can identify A with GM for some mass M . If one considers the energy of the vortex per unit length as $\rho_m c^2$ times volume, one can find an effective mass distribution. But a simpler route is to go directly to force.

C2. Force between two coaxial vortices

Now take two identical vortex filaments (circulation κ each) aligned on the z -axis, separated by a distance Z along z . We imagine them as representing two distant swirling objects (like two hydrogen atoms on an axis, each with a vortex line through it). The question is: what is the force between them?

In fluid statics, two parallel vortex lines with the same circulation sign actually rotate around each other (like a vortex pair), and two vortex rings on an axis either move apart or together depending on flow direction. But if they are steady and coaxial, one can calculate an interaction energy or pressure field superposition.

Since the flow is irrotational outside the cores except on the axis (no other vorticity introduced), we can approximate the pressure field as a superposition of each vortex's field:

$$p_{\text{total}}(r, z) \approx p_{\infty} - \frac{\rho_f \kappa^2}{8\pi^2} \left(\frac{1}{r^2 + (z - Z/2)^2} + \frac{1}{r^2 + (z + Z/2)^2} \right) / 2$$

The net force on one vortex due to the presence of the other can be found by integrating the pressure over its core cross-section area facing the other vortex. Equivalently, one can derive it from the gradient of an interaction energy.

By symmetry, the force will be along $\pm z$ axis. The pressure on the end of one vortex (facing the other) is lower than on the far side, causing it to be pulled. The difference in pressure Δp across a cross-section roughly of area A yields force $F = \Delta p, A$.

Using Eq. (26), the pressure due to vortex 2 at the location of vortex 1 (on-axis) is:

$$\Delta p_{2 \rightarrow 1} = -\frac{\rho_f \kappa^2}{8\pi^2 Z^2}$$

Multiplying by the cross-sectional area of the vortex core $A \sim \pi a^2$ (where a is core radius) gives the force magnitude:

$$F \approx \frac{\rho_f \kappa^2 a^2}{8\pi Z^2} \quad (28)$$

Now, $\rho_f \kappa^2 a^2/(8\pi)$ can be interpreted. If we identify $\mu = \rho_f \pi a^2$ as the “mass per unit length” of the vortex core (since $\rho_m = \rho_f v^2/2c^2$ and v^2 at core relates to κ/a , but let's assume non-relativistic so mass per length $\approx$ energy per length/ $c^2 = \rho_f v^2/2c^2$ · area?), it gets tricky to interpret directly. But qualitatively, $F \propto 1/Z^2$.

Comparing to Newton's law $F = G \frac{m_1 m_2}{Z^2}$, we can read off an effective gravitational constant times masses. We expect $m_1 \sim m_2 \sim$ (mass of vortex represented hydrogen atom). The derivation here is crude, but one can refine by considering the momentum flux in the fluid and using the vortex self-energy as surrogate for mass. A more rigorous approach is to compute the interaction energy via fluid kinetic energy:

$$E_{\text{int}} = \frac{1}{2} \rho_f \int (\mathbf{v}_1 \cdot \mathbf{v}_2) dV$$

To put it succinctly: the potential energy of two aligned vortices is

$$U(Z) \propto -\frac{\rho_f \kappa^2 \ell}{8\pi^2} \frac{1}{Z}$$

$$F = -\frac{dU}{dZ} \propto -\frac{\rho_f \kappa^2 \ell}{8\pi^2} \frac{1}{Z^2}$$

By identifying $\frac{\rho_f \kappa^2 \ell}{8\pi^2} = GM_1 M_2$, we can define the masses:

$$M = \frac{\rho_f \kappa^2 \ell}{8\pi^2 G}$$

Thus, the emergent gravity constant in SST is:

$$G_{\text{swirl}} = \frac{\rho_f \kappa^2}{8\pi^2} \quad (29)$$

as quoted earlier in Appendix A in different notation. Achieving $G_{\text{swirl}} = G_N$ is one calibration condition of the theory.

In summary, two swirling objects attract with $F \approx G_{\text{swirl}} \frac{M_1 M_2}{Z^2}$, thereby reproducing Newton's law in the appropriate regime. The “mass” here corresponds to fluid-dynamical mass (energy of swirl tied up in the object divided by c^2). Non-chiral swirl distributions (like opposite circulations bound together) yield little far-field $1/r$ effect and thus behave as having less active gravitational mass—this addresses why symmetric configurations might be gravitationally inert (or nearly so, akin to how two opposite vortices side by side produce near-cancellation of far field).

One must note limitations: our derivation assumed incompressibility and ignored relativistic corrections. If flows approach c , or if pressures become too large, modifications would occur (potentially connecting to GR deviations). However, for typical subsonic internal flows (e.g. electrons swirling at $v \sim c\alpha$ where α is fine structure constant, that's $\sim 0.007c$ —non-relativistic), these formulas hold.

$$V_{\text{SST}}(r) = -\Lambda / \sqrt{r^2 + r_c^2}$$

We have thus shown how Newtonian gravitational attraction emerges from fluid pressure deficits around swirl strings, making concrete the heuristic arguments of Sec. 4.

Appendix D: Electromotive Impulse from Swirl Density Change

This appendix provides a derivation for the modified Faraday's law and the resulting quantized EMF impulse that occurs when the linking number between a swirl string and an electrical loop changes by one.

D1. Integral form derivation of modified Faraday law

We begin with Faraday's law in integral form for a loop C bounding a surface S :

$$\oint_C \mathbf{E} \cdot d\boldsymbol{\ell} = -\frac{d}{dt} \int_S \mathbf{B} \cdot d\mathbf{A} \quad (30)$$

Now imagine that through the surface S , in addition to magnetic flux, there are $\mathcal{N}(t)$ swirl strings passing (each string here meaning a vortex line carrying quantized circulation κ). These

swirl strings are not electromagnetic in themselves, but SST posits that a time-varying $\mathcal{N}$ will induce an $\mathbf{E}$ -field circulation.

We hypothesize a modified law:

$$\oint_C \mathbf{E} \cdot d\boldsymbol{\ell} = -\frac{d\Phi_B}{dt} - \frac{d\Phi_\circ}{dt} \quad (31)$$

where $\Phi_B = \int_S \mathbf{B} \cdot d\mathbf{A}$ is the ordinary magnetic flux, and $\Phi_\circ$ is a new term proportional to the number of swirl strings linking C . Specifically, since each swirl string is associated with a circulation quantum κ and coupling constant $G_\circ$, the dimensional analysis and consistency with Maxwell's equations require $\Phi_\circ = -G_\circ \mathcal{N}(t)$ (the minus sign just to follow Lenz's law sign conventions conveniently).

To determine $G_\circ$, consider a single event: one swirl string appears through S (so $\Delta\mathcal{N} = +1$) over a short time Δt . Then Eq. (31) would predict a sudden jump in the left side (EMF) equal to $-\Delta\Phi_\circ/\Delta t = G_\circ(\Delta\mathcal{N}/\Delta t)$. Taking Δt short, effectively an impulse $\mathcal{E} = \int \mathbf{E} \cdot d\boldsymbol{\ell} dt = G_\circ \Delta\mathcal{N}$. For one vortex ($\Delta\mathcal{N} = 1$), $\mathcal{E} = G_\circ$. This $\mathcal{E}$ has units of Weber (since $\mathbf{E} \cdot d\boldsymbol{\ell} dt$ integrates to magnetic flux unit). Empirically, if such an impulse is observed and equals one flux quantum $\Phi_0 = h/2e$, we can equate:

$$G_\circ = \Phi_0 = \frac{h}{2e} \quad (32)$$

This was an assumption in our main text that the coupling is identified with the known quantum of flux. If one did not know Φ_0 from superconductivity, one could attempt to derive $G_\circ$ from consistency in dimensions or perhaps from a quantum argument: a change of one vortex linking corresponds to a phase change of 2π in a superconducting wavefunction around the loop, which by flux quantization implies the EMF must equal $h/2e$. In any case, taking $G_\circ = h/2e$ allows us to rewrite the modified Faraday law compactly as:

$$\nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t} + \frac{h}{2e} \frac{\partial \varrho_\circ}{\partial t} \hat{\mathbf{n}} \quad (33)$$

where $\varrho_\circ$ is the areal density of swirl strings (so $\int_S \partial_t \varrho_\circ dA = \partial_t \mathcal{N}$) and $\hat{\mathbf{n}}$ is oriented by right-hand rule to link orientation (thus giving direction to the induced field consistent with orientation conventions).

D2. Electromotive impulse calculation

We now consider the scenario in detail: A single vortex loop initially wholly outside loop C enters and passes through C , linking it once, then perhaps annihilates or leaves. During this process, what is the induced voltage $V(t)$ in a physical coil following C ?

From Eq. (31),

$$\oint_C \mathbf{E} \cdot d\boldsymbol{\ell} = -\frac{d\Phi_B}{dt} + \frac{h}{2e} \frac{d\mathcal{N}}{dt} \quad (34)$$

If no change in ordinary magnetic flux Φ_B happens (assume the vortex has no actual magnetic field tied to it; it might carry some effective vector potential coupling but no real magnetic flux in vacuum scenario), then

$$\oint_C \mathbf{E} \cdot d\boldsymbol{\ell} = \frac{h}{2e} \frac{d\mathcal{N}}{dt} \quad (35)$$

Integrate both sides over the entire event time (from before vortex entry to after vortex exit):

$$\int_{t_i}^{t_f} \oint_C \mathbf{E} \cdot d\boldsymbol{\ell} dt = \frac{h}{2e} \int_{t_i}^{t_f} \frac{d\mathcal{N}}{dt} dt \quad (36)$$

The right side is $\frac{h}{2e}(\mathcal{N}(t_f) - \mathcal{N}(t_i))$. If initially no vortex and finally one vortex passed (or vice versa if it annihilates), $\Delta\mathcal{N} = \pm 1$. So the integral is $\pm h/2e$. The left side is the net flux

change through the coil, which equals $\int V(t) dt$ (the time-integral of induced voltage around the loop). Thus:

$$\int_{-\infty}^{\infty} V(t) dt = \pm \frac{h}{2e} \quad (37)$$

This means the total time-integrated voltage (the area under the voltage vs. time curve, or equivalently the step change in flux linkage through the coil) is one flux quantum. The sign depends on whether a vortex enters (+) or leaves (−) and on relative orientation (Lenz’s law sign — an entering vortex will induce a polarity that opposes it in a certain orientation etc.). This is a crisp, binary prediction: one vortex event, one flux quantum impulse.

If multiple vortices passed in sequence, the impulses add, giving multiples of Φ_0 . If a vortex-antivortex pair nucleates and one goes through while the other goes the opposite side, one might get opposite sign impulses cancel if both link the loop, or separate signals if only one links, etc. The possibilities align with the idea that only linking number changes matter, not other details.

Notably, this result is independent of the coil’s shape or resistance; even if it’s not a superconductor, an EMF is induced and if open-circuited, it results in a certain integrated voltage. In a closed circuit, it would drive a current such that the flux quantum mostly ends up counteracted by induced current (in a superconductor, it would persist as current).

We can examine the consistency with energy conservation. A vortex moving through a loop generates an EMF which could do work on the circuit (like lighting a lamp briefly). Where does that energy come from? In SST, it comes from the kinetic energy of the vortex or medium. Essentially, as the vortex passes the loop, it experiences a drag from Lenz’s law: the induced current’s magnetic field exerts a force on the vortex, slowing it (or requiring work to push it through). This is analogous to how moving a magnet into a coil transfers energy from your work into electrical energy. In SST, the coupling $G_\circ$ is set so that one quantum of circulation yields exactly one quantum of flux, so the work done per event is (current) $\times$ (flux quantum) and so on, matching known quantum relations.

D3. Gauge field derivation

We can also derive the modified Faraday law via the W_μ gauge field formalism. The presence of a swirl string is akin to a quantized pseudomagnetic flux tube for the W field. In Maxwell’s equations with sources, a changing “magnetic” flux (here pseudomagnetic from swirl) yields an $\mathbf{E}$. In fact, in the W field Bianchi identity:

$$\partial_{[\sigma}(\nabla \times \mathbf{E})_{\mu\nu]} = \partial_{[\sigma}(\partial_\mu B_{\nu]} + G_\circ \partial_\mu \tilde{W}_{\nu]}) = 0 \quad (38)$$

one sees $\partial_t(\nabla \times \mathbf{E}) = -\nabla \times \partial_t \mathbf{B} + G_\circ \partial_t \nabla \times \tilde{\mathbf{W}}$. But $\nabla \times \tilde{\mathbf{W}}$ corresponds to swirl string areal density in that representation. Working through yields the same term.

In conclusion, the swirl–EM induction effect is a direct consequence of including the topological coupling between B -field (fluid vortex field) and the electromagnetic A -field (vector potential). One could show that an added term $L_{\text{int}} \sim G_\circ B_{\mu\nu} F^{\mu\nu}$ in the Lagrangian, when varied, produces the modified Faraday equation. However, such a term might break gauge invariance unless $B_{\mu\nu}$ is a curl of some potential (which it is). A safer way is to incorporate it via the W fields and the induction term $b_\circ = G_\circ \partial_t \varrho_\circ$ (like in Table III).

The upshot: whenever the topology of the swirl configuration changes (a linking number through a loop), an electromotive response of fixed magnitude results. This is a clear, falsifiable prediction that sets SST apart from conventional EM (which would predict no such effect without actual magnetic fields or charges present).

If an experiment sees, for example, a flux quantum entering a SQUID absent any applied magnetic flux but coincident with an engineered vortex event, this would strongly support SST’s concept. On the other hand, stringent null results would constrain $G_\circ$ to be much smaller than

$h/2e$, perhaps forcing it effectively to zero, which would push SST into a corner or exclude it. The experiment's feasibility and current status are beyond this appendix, but as noted, analogous effects in superconductors already inform us that if the vacuum acts similarly, Φ_0 is the relevant scale.

This completes our derivation of the modified Faraday law and the quantized impulse phenomenon.

Appendix A

SST-44 — Parallel capstone

Swirl–String Theory: Canonical Fluid Reformulation of Relativity and Quantum Structure Omar Iskandarani

Independent Researcher, Groningen, The Netherlands

ORCID: 0009-0006-1686-3961,

DOI: 10.5281/zenodo.xxxx April 19, 2026

Abstract

We present *Swirl-String Theory* (SST), a fluid-topological framework in which matter and radiation are modeled as quantized vortex loops (“swirl strings”) in an incompressible, non-dissipative condensate. Within SST, classical gravitational phenomenology *emerges* as a collective pressure effect in flat space, while local swirl flows *recover* relativistic time-dilation kinematics. We formulate a covariant effective field theory on a preferred foliation, show how topological quantization organizes a discrete particle spectrum, and outline a route by which gauge structures may be *derived* from orientational textures of the medium. A modified Faraday law is proposed in which time-varying swirl areal density sources an electromotive impulse; this yields geometry-independent, quantized flux signatures that serve as explicit falsification targets. Wave-particle duality is described via R/T phase dynamics (unknotted vs. knotted states), with measurement modeled as an $R \leftrightarrow T$ transition. We compare SST with Kelvin’s vortex lineage, analogue-gravity programs, and emergent-gauge constructions, emphasizing where SST *recovers* established limits (Newtonian gravity, Maxwell electrodynamics, quantum interference) and where it *predicts* deviations that are testable in BECs, superconducting films, and attosecond spectroscopy. All equations use SI units with dimensional checks. Our aim is a parameter-light, topological account that complements standard formulations and invites direct experimental appraisal. *Keywords:* vortex dynamics; topological fluid; emergent gauge theory; time dilation; quantum measurement; attosecond spectroscopy

A.1 Introduction

Reconciling General Relativity (GR) with Quantum Mechanics (QM) remains difficult because GR treats spacetime as dynamical and curved, whereas quantum field theories (QFT) typically operate on a fixed background. Swirl-String Theory (SST) approaches this tension by positing a unified substrate: a flat-space, incompressible condensate that supports and constrains all excitations.¹

Motivation: Recasting the GR–QM Mismatch

The long-standing mismatch between the Standard Model (SM) and GR suggests re-examining some starting assumptions. SST recasts curvature as effective fluid kinematics within a background Euclidean manifold. In this picture, what is interpreted as gravity *emerges* from pressure and flow gradients in the condensate rather than from a fundamental geometric interaction. Treating both gravitational and quantum effects within one hydrodynamic framework *aims* to clarify shared mechanisms while preserving empirical constraints from GR and the SM.

Historical Lineage: From Vortex Atoms to Analogue Gravity

SST fits within a broader tradition that links topological fluid structures to matter. Early ideas—such as Kelvin’s vortex atoms—anticipated the use of knotted, conserved circulations as building blocks. Modern developments in knot theory, quantized circulation, and analogue gravity deepen this thread: horizons and other curved-spacetime analogues can be *recovered* in superfluids and Bose–Einstein condensates. SST draws on this lineage by using the stability of vortex dynamics to *derive* a durable particle spectrum.

SST Proposition: A Topological Fluid Substrate for Particles and Interactions

SST posits a single universal medium—the “swirl” condensate—characterized by effective density, core swirl speed, and core radius. Physical observables such as mass, forces, and time dilation *emerge* from topologically protected excitations: knotted swirl strings. Formally, the framework resembles a modern Lorentz–Ether–style construction: it lives on a flat 4D manifold with an absolute time parameter and a preferred foliation set by the condensate’s unit timelike flow. The appeal of this background lies in what it may *predict*: topological constraints on mass generation and the *recovery* of SM-like gauge structure offer a path toward fewer free parameters than many effective models. At the same time, SST accepts empirical equivalences with standard formulations wherever they obtain and treats departures as testable claims rather than assumptions.

Relation to Emergent-Gravity Programs

Conceptually, SST aligns with approaches in which gravity *derives* from statistical or informational tendencies rather than standing as a separate fundamental interaction. In SST, a time-varying swirl density acts like an entropy- or information-density field. Resulting hydrodynamic gradients track the gradient of this swirl-based entropy and *recover* attractive behavior consistent with gravitational phenomenology. This analogy is used as an organizing principle rather than as a claim of superiority; points of agreement and potential deviation are presented as targets for calculation and experiment. While string theory often pursues unification through extra-dimensional vibrational modes [19], SST frames unification in terms of topological fluid structures in a fixed 4D background. Consistent with calls for sharper empirical criteria [20], we emphasize falsifiable consequences (Secs. A.9, A.14).

¹Several preprints by the author are cited to reflect distinct developments of the SST framework. Each addresses separate derivational or phenomenological components. Peer-reviewed validation is ongoing.

A.2 Core Postulates of Swirl–String Theory

SST is defined by six axioms that constrain the dynamics and properties of a universal condensate and its topological excitations. Core postulates follow from SST Canon v0.5 [1], with corresponding derivations in [4].

Axioms from the Canon v0.5

1. **Incompressible swirl condensate.** Physics is modeled on Euclidean $\mathbb{R}^3$ with an absolute time t . The background substrate is a frictionless, incompressible fluid ($\nabla \cdot \mathbf{v} = 0$) that supplies the kinematic stage on which all excitations evolve.
2. **Swirl strings as knotted topological excitations.** Particles and field quanta are represented by closed, stable vortex filaments (“swirl strings”). Their discrete quantum numbers *derive* from knot topology and linking invariants, providing a topological state space for matter and radiation.
3. **Quantized circulation** ($\Gamma = n\kappa$). The circulation of the swirl velocity $\mathbf{v}_\circ$ around any closed loop C is *quantized* in integer multiples of a fundamental quantum κ ,

$$\Gamma = n\kappa,$$

with $\kappa = h/m_{\text{eff}}$ linking topological class to a quantum scale, in parallel with Onsager–Feynman quantization in superfluids [9].

4. **Swirl clocks and local time dilation** (S_t). Local proper time is set kinematically by the local tangential swirl velocity v , with

$$S_t = \frac{dt_{\text{local}}}{dt_\infty} = \sqrt{1 - \frac{v^2}{c^2}}, \quad (\text{A.1})$$

which *recovers* the standard special–relativistic time-dilation factor. In the SST picture, high swirl speeds co-vary with deeper effective gravitational potentials, so clocks slow where the medium’s flow is strongest. The claim is kinematic equivalence, not empirical replacement: SST aims to match established relativistic tests while offering a medium-based account of the same effects.

5. **Dual R-phase and T-phase states.** Swirl strings exhibit a two-phase description: an extended, unknotted R-phase (radiative, wave-like, effectively massless) and a localized, knotted T-phase (tangible, particle-like, mass-carrying). Quantum duality is modeled as a dynamical $R \leftrightarrow T$ transition.
6. **Canonical knot–particle correspondence (illustrative mapping).** Specific knot classes are proposed to *correspond* to particle species. A representative assignment places charged leptons on torus knots (e.g., electron $\leftrightarrow$ trefoil 3_1) and quarks on chiral hyperbolic knots (e.g., up quark $\leftrightarrow 5_2$, down quark $\leftrightarrow 6_1$). Massless bosons (e.g., photons) are modeled as unknotted R-phase torsional excitations. These mappings are presented as testable, calculable hypotheses anchored in topological invariants rather than as presupposed identities.

A.3 Lagrangian and Field-Theoretic Framework

The dynamical content of SST is organized as a covariant effective field theory (EFT) defined on a preferred time foliation supplied by the condensate flow [4]. The formalism is intended to *recover* standard phenomenology where required while making distinct, testable predictions where the medium’s structure is consequential.

Preferred foliation via clock field, projectors, and Khronon sector

A scalar “clock” field selects a unit timelike 4-velocity that defines the preferred frame. Spatial dynamics are confined to leaves orthogonal to this flow via a projector construction. The action includes a Khronon sector with gradient terms for the time field and associated couplings. Constraints from multimessenger observations (e.g., GW170817) motivate parameter choices that *fix* the tensor-mode speed to be luminal, aligning the EFT with gravitational-wave propagation bounds.

Two-form vorticity, non-Abelian swirl connection, and emergent gauge structure

Topological degrees of freedom are described by two complementary ingredients. A two-form potential captures coherence and topological charge with a topologically conserved field strength. In parallel, an emergent non-Abelian “swirl connection” encodes coarse-grained orientational textures of the string network and takes values in a compact Lie algebra; its curvature measures defect density in the medium. Upon integrating out short-distance structure, the EFT *recovers* a Yang–Mills sector whose modes correspond to the medium’s internal excitations, offering a route to SM-like gauge interactions as emergent phenomena [1, 4]. The emphasis is on derivation and empirical equivalence, not on asserting prior superiority.

Canonical Lagrangian formalism

A minimal consistent Lagrangian includes kinetic terms for the clock and vorticity sectors, matter couplings to the swirl connection, and protected topological terms. Stability follows from conserved charges and topological invariants, with contributions such as a Chern–Pontryagin density enforcing the requisite conservation laws in the action.

Summary of the Canonical Lagrangian (see Appendix .1)

On a flat 4D background with a preferred foliation selected by a scalar clock field Φ , we define the unit timelike vector $u^\mu \propto \partial^\mu \Phi$ and the spatial projector $P^{\mu\nu} = \eta^{\mu\nu} + u^\mu u^\nu$. The SST action decomposes into the following sectoral terms:

$$\mathcal{L} = \mathcal{L}_\Phi + \mathcal{L}_B + \mathcal{L}_{\text{YM}} + \mathcal{L}_{\text{mat}} + \mathcal{L}_{\text{top}} + \mathcal{L}_{\text{bridge}} \quad (\text{A.2})$$

- **Clock (Khronon) sector $\mathcal{L}_\Phi$.** Fixes the preferred foliation via the unit field u^μ (constructed from Φ). Couplings are chosen such that $c_{13} \equiv c_1 + c_3 = 0$, yielding a luminal tensor mode consistent with multimessenger constraints.
- **Two-form (vorticity) sector $\mathcal{L}_B$.** Encodes coherent vorticity with a two-form potential $B_{\mu\nu}$ and conserved field strength $H_{\mu\nu\rho} = 3\partial_{[\mu}B_{\nu\rho]}$; a topological density $\mathcal{L}_{\text{top}}$ (e.g., Chern–Pontryagin-type) protects the associated charge.
- **Emergent gauge (swirl connection) sector $\mathcal{L}_{\text{YM}}$.** Introduces a non-Abelian connection $A_\mu \in \mathfrak{g}$ with curvature $G_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu + [A_\mu, A_\nu]$; the Yang–Mills form governs the coarse-grained orientational textures of the swirl network.
- **Matter (knot–soliton) sector $\mathcal{L}_{\text{mat}}$.** Dirac fields ψ_K minimally coupled via D_μ to the emergent connections on the spatial leaves ($P^\mu_\nu D_\mu$), with solitonic rest mass $m_K^{(\text{sol})} = \mathcal{M}_0 \Xi_K$ as in Eq. (A.3). [16, 15]
- **Bridge term $\mathcal{L}_{\text{bridge}}$.** Realizes the modified Faraday law by coupling the time variation of the swirl areal density $\partial_t \rho_\circ$ to the U(1) field strength, yielding $\nabla \times \mathbf{E} = -\partial_t \mathbf{B} - \mathbf{b}_\circ$ with $\mathbf{b}_\circ \propto \mathcal{G}_\circ \partial_t \rho_\circ \hat{n}$ (Sec. A.9) [2].

Explicit densities, coefficient normalizations, and symmetry restrictions (foliation-preserving diffeomorphisms; topological charge conservation) are listed in Appendix .1.

Clock-Rescaled Schrödinger Evolution

Let the local clock factor follow the postulate (A.1):

$$\tau(\mathbf{x}) = \sqrt{1 - \frac{|\mathbf{v}_\perp(\mathbf{x})|^2}{c^2}} \in [0, 1], \quad \mathbf{v}_\perp \text{ tangential to the swirl sheet.}$$

Then laboratory-time evolution of a bound mode Ψ is

$$\boxed{i\hbar \partial_{t_\infty} \Psi(\mathbf{x}, t_\infty) = \tau(\mathbf{x}) H[\mathbf{v}] \Psi(\mathbf{x}, t_\infty)},$$

with single-mode reduction $i\hbar \dot{\Psi} = \langle \tau \rangle H \Psi$. Small-speed expansion yields

$$\tau \approx 1 - \frac{|\mathbf{v}_\perp|^2}{2c^2} \Rightarrow \frac{\Delta\nu}{\nu} \approx -\frac{\langle |\mathbf{v}_\perp|^2 \rangle}{2c^2}.$$

This is the Hamiltonian-fluid realization of the clock law (cf. Hamiltonian fluids [47, 48]). **Appendix .8** gives the variational derivation and core-averaging details.

Mass via a solitonic knot energy functional

A central working hypothesis in SST is that fermion rest masses *emerge* as non-perturbative soliton energies of stable knotted excitations [4]. This aims to reduce reliance on freely tuned Yukawa parameters by tying masses to topological data through a canonical scaling law:

$$m_K^{(\text{sol})} = \mathcal{M}_0 \Xi_K(m, n, s, k; V_K, \phi_{\text{DSI}}), \quad (\text{A.3})$$

where the universal scale $\mathcal{M}_0$ is fixed by swirl-fluid parameters ($\mathbf{v}_\zeta, \rho_f, r_c$) and calibrated to the electron mass m_e . The dimensionless multiplier Ξ_K encodes knot-specific invariants—e.g., crossing number m , symmetry class s , chirality index k , and (for quark knots) hyperbolic volume V_K [4].

To incorporate helicity and torsional twist, a discrete-scale-invariance factor ϕ_{DSI}^{-2k} enters Ξ_K , with

$$\phi_{\text{DSI}} \equiv \exp\left(\text{asinh} \frac{1}{2}\right),$$

providing a canonical suppression of higher-chirality configurations.

Normalization and canonical limits.

- **Electron anchor:** Assign $\Xi_{3_1} = 1$ for the trefoil (3_1), fixing $\mathcal{M}_0 = m_e$.
- **R/T phase limit:** The R-phase (unknot) satisfies $\Xi_{\text{unknot}} = 0$ (massless), while T-phase knots yield $\Xi_K > 0$.
- **Quark-sector monotonicity:** For chiral hyperbolic knots, impose $\partial \Xi_K / \partial V_K > 0$ to preserve topological ordering across species.

Once $\mathcal{M}_0$ is calibrated, the fermion spectrum is *recovered* from knot class via Ξ_K , yielding a parameter-light account subject to direct comparison with observed masses.

A.4 Internal Gauge Symmetry in SST: $SU(3) \times SU(2) \times U(1)$

Postulate (Multi-director swirl). At each spacetime point, the coarse-grained swirl state carries a unit complex frame $\mathcal{D}(x) = (\hat{d}_1(x), \hat{d}_2(x), \hat{d}_3(x))$ with $\hat{d}_a \in \mathbb{C}^3$, orthonormal under the local inner product induced by the swirl metric. Phase rephasings and unitary rotations of this frame leave observable swirl-energy densities invariant.

Symmetry. Global rephasings $\hat{d}_a \rightarrow e^{i\alpha} \hat{d}_a$ generate a $U(1)$. Independent unitary rotations among the three directors generate an $SU(3)$. A two-component subframe $(\hat{d}_1, \hat{d}_2)$ supports an $SU(2)$ action. Promoting these to local transformations yields an internal gauge group $G = SU(3) \times SU(2) \times U(1)$.

Matter fields (knot excitations). A swirl string of topological type K supports a complex multiplet Ψ_K transforming in a representation $R_3(K) \otimes R_2(K)$ with $U(1)$ charge $Y(K)$. We define a minimal, computable map using knot descriptors:

$$t(K) = (L(K) \bmod 3, S(K) \bmod 2, \chi(K)),$$

where L is minimal crossing number, S is Seifert circle parity, and $\chi \in \{\pm 1, 0\}$ is a chirality index. Then

$$R_3(K) = \begin{cases} \mathbf{3} & \text{if } L \bmod 3 = 1, \\ \bar{\mathbf{3}} & \text{if } L \bmod 3 = 2, \\ \mathbf{1} & \text{if } L \bmod 3 = 0, \end{cases} \quad R_2(K) = \begin{cases} \mathbf{2} & \text{if } S \bmod 2 = 1, \\ \mathbf{1} & \text{if } S \bmod 2 = 0, \end{cases}$$

and the hypercharge is assigned by a linear functional of knot data,

$$Y(K) = \alpha_Y \chi + \beta_Y \text{wr}(K) + \gamma_Y g(K),$$

with $\text{wr}(K)$ the writhe and $g(K)$ the genus. The constants $(\alpha_Y, \beta_Y, \gamma_Y)$ are fixed by calibrating to the electron, up/down quarks, and neutrino benchmarks while satisfying anomaly cancellation (see App. .3).

Covariant derivative and field strengths. Introduce gauge bosons G_μ^A ($A = 1, \dots, 8$), W_μ^i ($i = 1, 2, 3$), and B_μ with couplings g_3, g_2, g_1 . With Gell-Mann λ^A and Pauli σ^i matrices,

$$D_\mu = \partial_\mu - ig_3 G_\mu^A \frac{\lambda^A}{2} - ig_2 W_\mu^i \frac{\sigma^i}{2} - ig_1 Y B_\mu, \quad (\text{A.4})$$

$$\begin{aligned} G_{\mu\nu}^A &= \partial_\mu G_\nu^A - \partial_\nu G_\mu^A + g_3 f^{ABC} G_\mu^B G_\nu^C, & W_{\mu\nu}^i &= \partial_\mu W_\nu^i - \partial_\nu W_\mu^i + g_2 \epsilon^{ijk} W_\mu^j W_\nu^k, \\ B_{\mu\nu} &= \partial_\mu B_\nu - \partial_\nu B_\mu. \end{aligned} \quad (\text{A.5})$$

Gauge sector Lagrangian (dimensionally consistent).

$$\mathcal{L}_{\text{gauge}} = -\frac{1}{4} G_{\mu\nu}^A G^{A\mu\nu} - \frac{1}{4} W_{\mu\nu}^i W^{i\mu\nu} - \frac{1}{4} B_{\mu\nu} B^{\mu\nu}. \quad (\text{A.6})$$

$[\mathcal{L}]$ = energy density; each term has mass dimension 4.

Matter–gauge coupling. For a knot species K with field Ψ_K ,

$$\mathcal{L}_{\text{matter}} = \bar{\Psi}_K (i\gamma^\mu D_\mu - m_K) \Psi_K + \dots, \quad (\text{A.7})$$

where m_K is determined by the SST mass functional (see main text; details in its appendix).

Physical charges from knot data. The electric charge follows $Q = T_3 + Y$ after electroweak mixing. Here T_3 is the eigenvalue of $\sigma^3/2$ acting on $R_2(K)$ and $Y = Y(K)$ above. Once $(\alpha_Y, \beta_Y, \gamma_Y)$ are fixed by a minimal set, all other Q become predictions tied to (χ, wr, g) of K .

Consistency checks (see App. .2, .3). (i) Abelian limit $(g_2, g_3 \rightarrow 0)$ reduces to QED with $A_\mu = s_w W_\mu^3 + c_w B_\mu$. (ii) Anomaly cancellation constrains the (Y, R_2, R_3) assignment; our knot map admits a unique fit matching the SM pattern within the chosen calibration set.

SST interpretation. The multi-director frame encodes internal “swirl phases.” Local reorientation is compensated by gauge connections (G_μ, W_μ, B_μ) that enter the energy via (A.6). Knot-type fixes representation content; chirality/writhe/genus fix hypercharge.

A.5 Swirl–Braid Correspondence and Energy Functional

In the Swirl–String framework, each quantized filament is represented by a smooth, oriented embedding

$$\Gamma : S^1 \rightarrow \mathbb{R}^3, \quad \Gamma(s) = \mathbf{r}(s), \quad s \in [0, 1),$$

with local swirl velocity field $\mathbf{v}_\zeta(\mathbf{r})$ and conserved fluid helicity

$$\mathcal{H} = \int_{\mathbb{R}^3} \mathbf{v}_\zeta \cdot (\nabla \times \mathbf{v}_\zeta) d^3x,$$

which is preserved in ideal (nondissipative) flow up to boundary terms.

Recent work [41] demonstrates that the connected sum of two nontrivial knots can collapse to the unknot via five explicit band moves. This supports the SST view of topological relaxation occurring through reconnection cascades, without requiring inverse braid unwinding.

In the SST framework, this implies that swirl-string configurations composed of multiple nontrivial loops may undergo collective reconnection to yield a trivial topological state, such as the vacuum or a null-charge excitation. Crucially, the process need not mirror the algebraic inverse of each braid component, highlighting a path toward irreversible helicity dissipation that bypasses local symmetry undoing. The five-move bound further suggests a topological transition cost in the energy landscape defined by Eq. (A.8).

Implication of Knot Fusion in SST

Two nontrivial swirl strings can dynamically reconnect into a swirl-loop with trivial topology (the unknot), even though neither component is individually trivial. This suggests:

- Topological charge may vanish without stepwise reversal.
- Swirl-string decay may proceed via collective reconnection cascades.
- The number of required reconnections (five) may bound the helicity dissipation channel.

This insight strengthens the SST model of quantized decay without requiring symmetry-based subtraction.

Braid closure for knotted filaments. A classical result ensures every oriented knot or link is the closure of a braid:

Theorem 1 (Alexander [21]). *Every oriented knot or link in $\mathbb{R}^3$ is equivalent to the closure of some n -strand braid $w \in B_n$.*

Thus each swirl string can be associated with an element of the Artin braid group

$$B_n = \langle \sigma_1, \dots, \sigma_{n-1} \mid \sigma_i \sigma_j = \sigma_j \sigma_i \ (|i - j| > 1), \ \sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1} \rangle,$$

where σ_i denotes a right-handed crossing of strands i and $i+1$ (and σ_i^{-1} a left-handed crossing).

Trefoil as a braided swirl loop

The canonical electron configuration in SST—the trefoil knot—admits a 3-strand braid closure, e.g.

$$w_e = \sigma_1^3,$$

so dynamical tightening can be viewed as a braid-word contraction driven by swirl tension. Each σ_i represents a localized swirl crossing; closure enforces the single-valued phase condition $\Gamma(0) = \Gamma(1)$.

Swirl–braid energy functional

To formalize contraction dynamics, we use a canonical effective functional on a braid representative w :

$$\mathcal{E}_{\text{eff}}[w] = \alpha C_{\min}(w) + \beta L(w) + \gamma \mathcal{H}(w). \quad (\text{A.8})$$

Here $C_{\min}(w)$ is the minimal crossing number over equivalent braid representatives of the same link type (to avoid projection dependence), $L(w)$ is the physical core-line length, and $\mathcal{H}(w)$ is the helicity associated with the configuration. Coefficients scale with swirl-fluid parameters as

$$\alpha \sim \frac{\rho_f r_c^3}{\|\mathbf{v}_\odot\|^2}, \quad \beta \sim \rho_E r_c, \quad \gamma \sim \frac{1}{2} \rho_f r_c^2,$$

where ρ_f is the effective fluid density, r_c the core radius, and ρ_E the energy per unit length of the filament.

Markov moves as physical updates. Alexander’s theorem plus Markov’s theorem implies that closures of braids related by conjugation and stabilization represent the same link type:

$$w \rightarrow \sigma_i w \sigma_i^{-1} \quad (\text{conjugation}), \quad w \rightarrow w \sigma_n^{\pm 1} \quad (\text{stabilization}).$$

We interpret these as local reconnections/reorderings and strand creation/annihilation on the coarse-grained worldsheet. In ideal swirl flow, helicity is conserved (Moffatt) and changes only through controlled dissipative channels; $\mathcal{E}_{\text{eff}}$ decreases monotonically under allowed relaxation.

Topological quantization and swirl charge

The statistical treatment of knotted vortex dynamics finds lineage in early work on quantized circulation [9], anchoring SST’s helicity-based energy model. For thin, isolated filaments, the Calugăreanu–White–Fuller relation connects link, twist, and writhe of a framed curve: $L_k = T + W$ [24, 25, 26]. In fluid dynamics, helicity decomposes into self- and mutual-linking contributions with circulation weights (Moffatt) [23]. In the single-filament limit with fixed circulation, we use the schematic form

$$\mathcal{H} \propto T + W,$$

noting that the precise coefficients depend on circulation and framing. The associated quantized swirl charge is

$$Q_\odot = \frac{\rho_f}{2\pi} \int_\Sigma (\nabla \times \mathbf{v}_\odot) \cdot d\mathbf{S},$$

which, in SST, maps algebraically to braid data (e.g., braid index n) and to the particle-family tuple used in Sec. A.12.

Connection to Chern–Simons field theory

In a continuum description, braid closures correspond to Wilson loops in three-dimensional Chern–Simons theory:

$$S_{\text{CS}} = \frac{k}{4\pi} \int A \wedge dA + \frac{2}{3} A \wedge A \wedge A, \quad \langle W(\Gamma) \rangle \propto V_K(q),$$

where V_K is the Jones polynomial of the knot K evaluated at $q = \exp(\frac{2\pi i}{k+2})$ for the $\text{SU}(2)$ theory [27]. This furnishes a formal bridge between the swirl-string energy landscape and topological quantum invariants.

Physical Picture

A braided swirl string behaves like multiple helical vortices twisting around each other and closing into a loop. As tension tightens the bundle, crossings smooth while preserving helicity, and the configuration settles into a quantized knotted state (e.g., the trefoil) corresponding to an elementary excitation.

A.6 Local Swirl Stress and Lab Traction

Define $\boldsymbol{\omega} = \nabla \times \mathbf{v}$. A symmetric local Cauchy stress consistent with the surrogate energy of the filament core is

$$\sigma_{ij} = \rho_f v_i v_j + \rho_f r_c^2 \left(\omega_i \omega_j - \frac{1}{2} \delta_{ij} |\boldsymbol{\omega}|^2 \right) - \delta_{ij} \frac{1}{2} \rho_f |\mathbf{v}|^2 \quad [\text{Pa}],$$

cf. vortex-tension analogies in fluids [42, 43, 44]. **Consequences.** (i) Traction on a probe at radius r scales $\propto r^{-2}$. (ii) Sign flips with filament chirality. (iii) Area integration reproduces the $\ln(R/\xi)$ per-length energy with the thermal prefactor from Sec. A.10. **Appendix .11** derives σ_{ij} from the local Lagrangian and lists boundary conditions for lab probes.

A.7 Emergent Gravity and Time Dilation

In SST, gravity is reinterpreted as a hydrodynamic attraction resulting from conserved circulation in a flat background [3].

Swirl-Induced Pressure Gradients as Gravitational Attraction

Massive particles, represented by stable chiral knotted strings, maintain a persistent, non-vanishing circulation around a central axis. This circulation induces a radial pressure deficit along the axis, governed by the Euler fluid balance equation [1]. When two neutral, composite systems (e.g., two proton cores within an H_2 molecule) share this central line, their circulations add, intensifying the pressure well (for two protons). This shared pressure deficit draws the systems together, producing the observed long-range inverse-square gravitational attraction in flat space, known as the Hydrogen-Gravity Mechanism. We treat this as a constructive mechanism with testable scaling relations; it is not assumed to replace post-Newtonian fits *a priori*.

Derivation of Matching Newton’s Constant

The effective gravitational coupling G_{swirl} is derived from the core swirl parameters of the medium:

$$G_{\text{swirl}} = \mathcal{G}_{\mathcal{O}} = \frac{v_{\mathcal{O}} c^5 t^2}{2 F_{\mathcal{O}}^{\text{max}} r_c^2} \approx G_N \quad (\text{A.9})$$

By calibrating the foundational constants ($v_{\mathcal{O}}, r_c$) and the maximum emergent electromagnetic force ($F_{\mathcal{O}}^{\max} \approx 2.9 \times 10^1$ N), G_{swirl} is numerically consistent with Newton’s constant G_N within calibration uncertainty [1]. This relation frames the gravitational coupling strength as constrained by the medium’s dynamics and the maximal electromagnetic tension it can support.

Composite Baryons as Merged Vortex Tubes

Baryons, such as the proton, are realized as composite swirl tubes formed by the merging of three quark knots (e.g., two up 5_2 and one down 6_1 twist-knots) at a Y-junction [3]. Due to Kelvin’s circulation theorem, the circulation is additive. Since each constituent quark carries circulation around the central axis, the baryon core possesses a total circulation. This increased circulation translates to a significant increase in the effective tangential core velocity and, consequently, a much deeper pressure well, correlating with the baryon’s larger rest mass.

Swirl Clock Effects Explain Gravitational Redshift

The Swirl Clock factor (Equation A.1) dictates that time runs slower in regions of higher swirl velocity [1]. Regions of concentrated mass (knotted strings) generate intense swirl flows. Consequently, intense gravitational potential wells are simply high-swirl-velocity regions where time dilation is pronounced. Gravitational redshift is therefore interpreted as a kinematic frequency shift arising from the difference in local time rates between a source located in a deep swirl region and an observer located in a quiescent, faster-ticking region of the medium [1].

A.8 Cosmological Branch: de Sitter Background in Swirl–String Theory

Canonical Context

Section A.7 derived the Hydrogen–Gravity Mechanism as a local manifestation of swirl–pressure wells in flat space, with the effective coupling $G_{\mathcal{O}}$ of Eq. (5) linking gravitational attraction to the medium’s intrinsic parameters. Here we extend that reasoning to the cosmological scale and show that the same mechanism, applied to a uniform positive swirl–pressure offset, reproduces the de Sitter expansion branch without invoking spacetime curvature.

A.8.1 From Local Swirl Wells to Global Expansion

The local pressure deficit responsible for attraction between knotted composites can be generalized to a uniform, sign-reversed bias. When the mean swirl pressure $p_{\mathcal{O}}$ of the medium exceeds its equilibrium value by a small positive constant,

$$p_{\mathcal{O}} = p_{\mathcal{O}}^{(0)} + \delta p_{\mathcal{O}}, \quad \delta p_{\mathcal{O}} > 0, \quad (\text{A.10})$$

the condensate experiences an isotropic expansion that, in General Relativity, would be attributed to a positive cosmological constant Λ .

A.8.2 Canonical Identification with the Cosmological Constant

For a spatially uniform medium with $\nabla \cdot \mathbf{v}_{\mathcal{O}} = 0$, define the mean volumetric expansion rate

$$H_{\mathcal{O}} = \frac{1}{3} \nabla \cdot \mathbf{v}_{\mathcal{O}}, \quad (\text{A.11})$$

and require dynamical balance between kinetic and pressure terms:

$$H_{\odot}^2 = \frac{p_{\odot}^{(0)}}{\rho_m c^2}. \quad (\text{A.12})$$

Comparison with the relativistic Friedmann relation $H^2 = \Lambda c^2/3$ yields the canonical mapping

$$\Lambda = \frac{3 p_{\odot}^{(0)}}{\rho_m c^4}. \quad (\text{A.13})$$

Thus, the cosmological constant is reinterpreted as the ratio of a uniform swirl–pressure bias to the medium’s mass–equivalent density.

A.8.3 Numerical Evaluation

Using the canonical constants

$$\rho_f = 7.0 \times 10^{-7} \text{ kg/m}^3, \quad \|\mathbf{v}_{\odot}\| = 1.09384563 \times 10^6 \text{ m/s},$$

the mass–equivalent density is

$$\rho_m = \frac{\frac{1}{2} \rho_f \|\mathbf{v}_{\odot}\|^2}{c^2} \simeq 4.2 \times 10^{-6} \text{ kg/m}^3.$$

Matching the observed $\Lambda \simeq 1.1 \times 10^{-52} \text{ m}^{-2}$ via Eq. (A.13) gives

$$p_{\odot}^{(0)} \simeq \frac{\Lambda \rho_m c^4}{3} \simeq 3.7 \times 10^{-9} \text{ Pa}. \quad (\text{A.14})$$

Hence, the “dark-energy” term corresponds to an exceedingly small positive swirl–pressure bias in the cosmic medium.

A.8.4 Unified Interpretation with the Hydrogen–Gravity Mechanism

Both the local gravitational wells of the Hydrogen–Gravity Mechanism and the global de Sitter expansion follow from the same constitutive law of the swirl medium:

$$\nabla p_{\odot} = -\rho_m \mathbf{a}_{\odot},$$

with $\mathbf{a}_{\odot}$ the swirl-induced acceleration field. Negative $\nabla p_{\odot}$ produces attraction (bound systems); a positive uniform offset produces large-scale repulsion (cosmic acceleration). Equation (5) for $G_{\odot}$ and Eq. (A.13) are therefore complementary limits of the same flat-space hydrodynamics.

A.8.5 Physical Picture and Testable Consequences

The de Sitter branch represents the flat, positive-pressure equilibrium of the swirl condensate. Every parcel of the medium carries a faint residual rotation that exerts an outward kinematic tension, driving the observed acceleration of cosmic expansion. If the swirl–pressure bias relaxes over time, the effective Λ_{eff} evolves as

$$\frac{d\Lambda_{\text{eff}}}{dt} = \frac{3}{c^4} \frac{d(p_{\odot}^{(0)})}{dt}, \quad (\text{A.15})$$

predicting small but potentially measurable deviations from a constant dark-energy density.

Summary

In Swirl–String Theory, the de Sitter universe is not a curved spacetime but a flat medium under a uniform swirl–pressure bias. Local gravitational attraction (Eq. 5) and global cosmic acceleration (A.13) arise as opposite-sign branches of the same constitutive relation of the swirl condensate.

A.9 Electromagnetic Emergence: Modified Faraday Law

We model electromagnetism as an emergent response of the swirl medium to topological dynamics [2]. In this view, non-adiabatic events—nucleation, annihilation, or reconnection of swirl strings—produce localized changes in the swirl areal density $\rho_{\odot}$ and thereby *drive* an electromotive impulse in the surrounding region.

Swirl String Nucleation/Annihilation Produces EM Impulses

A central prediction is that each integer change ΔN in the number of linking swirl strings yields a geometry-independent flux impulse,

$$\Delta\Phi = \mathcal{G}_{\circlearrowright} \Delta N,$$

with the sign fixed by chirality [2]. **(Testable)** This provides a sharp, falsifiable signature designed for SQUIDS or fast pickup loops; independence from detector geometry is part of the prediction rather than an assumption.




Figure A.1: **Canonical Swirl–Electromagnetic Coupling Diagram.** Causal and dimensional structure of the electromagnetic sector within the Swirl–String framework. The top layer extends Faraday’s law with a swirl-induced backreaction term $\mathbf{b}_\mathcal{G} = \mathcal{G}_\mathcal{G} \partial_t \mathbf{q}_\mathcal{G}$, encoding the electromotive response to time-varying swirl density in the medium. The middle layer represents the constitutive closure: $\mathbf{D} = \epsilon \mathbf{E}$ and $\mathbf{B} = \mu \mathbf{H}$, together with the mechanical correspondence $\mathbf{q}_\mathcal{G} = \chi_H \mathbf{H}$. The bottom layer completes the circuit with areal accumulation $\boldsymbol{\eta}$, source current $\mathbf{j}$, and the modified Ampère curl. All dimensionalities are shown for canonical homology between mechanical (swirl) and electromagnetic sectors, establishing the *Swirl–Electromagnetic Bridge* that underlies the flat-space emergence of Maxwellian dynamics.

The canonical coupling between swirl densities and electromagnetic response is depicted in Fig. A.1. The key mediator is the coupling constant $\mathcal{G}_\mathcal{O}$, which translates temporal changes in the swirl areal density $\rho_\mathcal{O}$ into observable electromotive responses. Strikingly, this same constant governs gravitational emergence within SST. When the swirl–electromagnetic system saturates its allowed flux bounds, the effective gravitational coupling becomes:

$$\mathcal{G}_\Omega \longrightarrow G_{\text{swirl}} \approx G_N,$$

numerically consistent with Newton’s constant (see Eq. A.9).

This canonical equivalence suggests a unifying physical principle: both electromagnetic induction and gravitational attraction arise from topological dynamics in the underlying swirl medium [2]. Swirl string nucleation events, which modify $\rho_{\mathbf{O}}$, thereby drive both classical forces—via a shared coupling.

Coupling via Induced Electric Field

This topological coupling modifies the standard Faraday law by introducing an additional source term $\mathbf{b}_\odot$, which is proportional to the time rate of change of the swirl areal density $\rho_\odot$ along the local orientation $\hat{n}$ [2]:

$$\nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t} - \mathbf{b}_\odot, \quad (\text{A.16})$$

where

$$\mathbf{b}_\odot = \mathcal{G}_\odot \frac{\partial \rho_\odot}{\partial t} \hat{n}.$$

We treat $\mathbf{b}_\odot$ as an effective source term that *recovers* standard Faraday induction when $\partial_t \rho_\odot = 0$.

Identification with Flux Quantum

The constant $\mathcal{G}_\odot$ is the universal topological transduction constant and, by dimensional analysis, carries units of magnetic flux (Weber) [2]. By normalizing the predicted impulse to observed quantum phenomena, it is *hypothesized* to coincide with the fundamental flux quantum $\Phi_\star = h/2e$ [2]. The experiments below are designed to adjudicate this identification.

Experimental Predictions

Attosecond spectroscopy now resolves the time scales on which chirality leaves a dynamical imprint [29, 30]. In methyloxirane, Han *et al.* [28] find robust forward-backward (FB) photoemission delays whose sign tracks molecular handedness. The swirl-EM coupling leads to clear experimental avenues [2]:

1. **SQUID loops detecting quantized impulses [6, 5, 10, 11]:** Quantized flux impulses $\Delta\Phi = n\Phi_\star$ during swirl-string creation or annihilation, with geometry independence and chirality-determined sign, are the primary electromagnetic test [2]. **(Testable)**
2. **Analog simulations in BECs [13, 14]. or type-II superconductors:** Controlled entry/exit of $h/2e$ vortices in superconducting films, and single-vortex dynamics in ultracold BECs, supply platforms to monitor voltage impulses and verify geometry independence and chirality sign flip [2]. **(Testable)**

A.10 Spectral Thermodynamics of Swirl Excitations

Define the geometric core factor and natural frequency scale

$$u_0(r) = \frac{\rho_f |\mathbf{v}_\odot|^2 r_c^2}{2 r^2} \quad [\text{J m}^{-3}], \quad \Omega_c = \frac{|\mathbf{v}_\odot|}{r_c} \quad [\text{s}^{-1}].$$

A Planck-like spectral energy density for swirl excitations is

$$u_{\text{vortex}}(r, \omega, T) = u_0(r) \lambda_{BE} \left(\frac{\omega}{\Omega_c} \right)^3 \frac{1}{e^{\hbar\omega/k_B T} - 1},$$

with dimensionless calibration λ_{BE} . Frequency integration yields

$$u_{\text{tot}}(r, T) = u_0(r) \lambda_{BE} \frac{\pi^4}{15} \left(\frac{k_B T}{\hbar \Omega_c} \right)^4, \quad P = \frac{1}{3} u_{\text{tot}}, \quad s = \frac{4}{3} \frac{u_{\text{tot}}}{T},$$

i.e. T^4 and T^3 scalings, with near-core r^{-2} localization (cf. blackbody thermodynamics [45, 44]). For a straight filament, $E/L \propto \ln(R/\xi)$ arises from $\int_\xi^R u_{\text{tot}} 2\pi r dr$ (classical vortex energetics [42]). **Appendix .9** provides density-of-states choices and finite-core regularization.

Photoelectric Mapping in SST Units

Let the EM/swirl-sector maximum force be $F_{\text{swirl}}^{\text{max}} = 29.053507$ N. Define the emergent action

$$h_{\text{SST}} = 4\pi \frac{F_{\text{swirl}}^{\text{max}} r_c^2}{|\mathbf{v}_{\odot}|} \quad [\text{J s}],$$

yielding the standard photoelectric law

$$E_{\text{max}} = h_{\text{SST}} f - \Phi, \quad f_0 = \Phi/h_{\text{SST}}.$$

Thus the slope of $E_{\text{max}}(f)$ is fixed by $(F_{\text{swirl}}^{\text{max}}, r_c, |\mathbf{v}_{\odot}|)$, while intensity controls current, not slope [46]. **Appendix .10** supplies the ring-emission geometry that produces the 4π factor and the unit checks mapping to Planck's h .

A.11 Chirality and Quantum Measurement Dynamics ($\text{R} \leftrightarrow \text{T}$)

SST provides kinematic origins for chiral time asymmetry and a dynamical alternative to the collapse postulate [1].

Interpretation of Attosecond Chiral Delays in SST

Attosecond spectroscopy has shown that molecular chirality introduces dynamical delays in photoionization, such as the forward/backward (FB) delay observed in methyloxirane [12]. SST interprets molecular chirality as a geometric constraint that fixes the orientation of the local swirl circulation, which directly dictates the orientation of the local Swirl Clock [6].

Enantiomers Exhibit Orientation-Dependent Swirl Clocks

Opposite enantiomers (L and D configurations) correspond to mutually reversed swirl-clock orientations [6]. The propagation phase of an emitted electron is therefore dependent on its direction relative to this fixed, orientation-sensitive time field [6].

Delay Sign Flips with Molecular Handedness

This kinematic interpretation leads to the prediction that reversing molecular handedness (swapping enantiomers) flips the sign of the measured forward/backward delay asymmetry [6]. This relies only on the geometric reversal of the kinematic time asymmetry and is consistent with reported experimental results, providing a direct test of the Swirl Clock principle [6]. (**Testable**)

$\text{R} \leftrightarrow \text{T}$ Phase Transitions as Measurement

SST models quantum measurement as a dynamical R-phase (wave-like) to T-phase (particle-like) transition with rate

$$\Gamma_{RT} \propto \frac{P}{A},$$

where P is incident power and A is area. The transition rate increases with incident power per area, aligning with environment-induced decoherence bounds and treating compatibility with such bounds as an empirical constraint [1].

A.12 Canonical Quantization and Topological Spectrum

SST provides a geometric derivation of quantum numbers based on knot topology [1, 4, 17].

SST Assigns Quantum Numbers via Knot-Theory Invariants

Discrete quantum properties are determined by topological invariants of the corresponding swirl strings. Electric charge (Q), weak isospin (I), and hypercharge (Y) are modeled as linear functions of linking, self-linking parity, and chirality:

$$(Q, I, Y) = f(\text{linking, parity, chirality}) \text{ [4, 1].}$$

Color representation (C) is set by net linking number modulo 3, assigning leptons ($C = 1$) as color singlets and quarks ($C = 3$) as color triplets [4].

Soliton-Based Mass Quantization Reduces Yukawa Freedom

The invariant mass of a fermion m_K is the soliton energy stored in the knot’s configuration (Eq. A.3) [4]. By defining the mass scale using the electron and anchor knots (up/down quarks) and relating Ξ_K to measures like hyperbolic volume V_K , SST reduces reliance on free Yukawa parameters by tying masses to topological invariants; quantitative fits are presented as tests of the scaling law [4]. **(Testable)**

Electroweak Parameters as Medium Stiffness Ratios

Two SM parameters become computable within the medium description. The electroweak mixing angle θ_W is determined by the ratio of stiffnesses (κ_1, κ_2) of emergent U(1) and SU(2) director fields,

$$\theta_W = \arctan\left(\frac{\kappa_1}{\kappa_2}\right) [1].$$

The Higgs VEV v_Φ is derived from the condensate’s bulk swirl energy density u_{swirl} , yielding $v_\Phi \approx 259.5 \text{ GeV}$, consistent with the empirical 246 GeV scale [1]. These are offered as concrete targets for refinement and comparison.

Knot Taxonomy: Visual Table of Particle Knot Mappings

Particle	Knot Type (Tangle Measure)	Topological Class	Notes
Photon (γ)	Unknot (0_1)	R-Phase Excitation	Massless, torsional pulse [1, 4]
Electron (e^-)	Trefoil (3_1)	Torus knot	Baseline lepton [1, 4]
Muon (μ^-)	Cinquefoil (5_1)	Torus knot	Higher-generation lepton [4]
Up quark (u)	5_2 knot	Chiral hyperbolic knot	Canonical baseline quark [1, 4]
Down quark (d)	6_1 knot	Chiral hyperbolic knot	Canonical baseline quark [1, ?]
Neutrino (ν)	Amphichiral link	Two-component link	Neutral, amphichiral linkage [4]
Proton (p)	Composite linkage	Baryonic composite tube	Additive circulation [3]

Table A.1: Canonical knot–particle correspondence.

A.13 Real-Time Tunneling and the Kappa Event

A further falsifiable prediction of Swirl–String Theory concerns *Kappa events*—defined as real-time, phase-coherent alignments of swirl dynamics across a potential barrier. In this picture, quantum tunneling is not an instantaneous phase advance but a finite-rate coherence transfer governed by the characteristic swirl speed $\|\mathbf{v}_\odot\|$. The measurable consequence is a dwell time that grows *linearly* with barrier length:

$$\tau_d^{(\text{SST})}(L) = \frac{L}{\|\mathbf{v}_\odot\|} \Xi, \quad (\text{A.17})$$

where $\Xi \geq 1$ is a dimensionless alignment overhead determined by the Kappa-transition rate. This contrasts sharply with the Hartman-type saturation of standard quantum mechanics, in which the phase (Wigner) delay tends toward a constant $\tau_H \sim \hbar/(V_0 - E)$ for opaque barriers [35, 36, 37].

A direct test uses a Larmor-clock geometry: a weak magnetic field confined to the barrier induces spin precession proportional to the in-barrier dwell time $\tau_d = \theta/\omega_L$. Scanning the barrier length L at fixed energy and measuring $d\tau_d/dL$ isolates the causal slope $\Xi/\|\mathbf{v}_\odot\|$ predicted by SST. A statistically significant positive slope would falsify Hartman saturation and confirm the real-time Kappa alignment mechanism.

The full derivation of the dwell-time scaling law, extraction of the Kappa transition rate Γ_κ , and numerical validation using the canonical constants ($\|\mathbf{v}_\odot\| = 1.09384563 \times 10^6 \text{ m s}^{-1}$, $r_c = 1.40897017 \times 10^{-15} \text{ m}$) are presented in Appendix .7.

A.14 Experimental Implications & Falsifiability

SST is presented with empirical testability as a design constraint [20]. The most direct signatures are (i) geometry-independent, quantized flux impulses from topological transitions and (ii) chirality-dependent attosecond delays in photoemission, both following from the swirl-EM coupling and Swirl Clock kinematics (Secs. A.9, A.11) [2]. Accordingly, SST offers clear avenues for experimental validation that target topological effects in emergent fields using SQUID loops, BEC/superconducting platforms, and ultrafast spectroscopy [2].

A.14.1 Identification from Slow and Non-Uniform Samples

We linearize the SST rig as a continuous-time descriptor system with LFT parameters and exploit steady-state plateaus and low-frequency sinusoids to estimate $H(0)$, $H(j\omega)$, and $\partial_s H|_{s=0}$ without uniform sampling. These frequency-point constraints feed the calibration of SST couplings (e.g., $\mathcal{G}_\odot$) while enforcing dimensional and passivity checks. Full details and estimators are provided in Appx. .6 (cf. the slow/non-uniform identification framework of [34]).

Key Falsifiable Predictions

1. **EM impulses of $\Delta\Phi = \pm\Phi_\star$ upon topological transitions:** Quantized flux impulses $\Delta\Phi = n\Phi_\star$ during swirl-string creation or annihilation, with geometry independence and chirality-determined sign [2]. **(Testable)**
2. **Attosecond delay sign flip with chirality reversal:** The sign of the measured forward/backward photoemission delay $\Delta\tau_{\text{FB}}$ flips strictly upon molecular enantiomer reversal, reflecting Swirl-Clock orientation [6, 12]. **(Testable)**
3. **Mass spectrum tied to knot taxonomy:** The scaling law in Eq. A.3, anchored by a single universal scale, yields mass ratios for higher generations and composite hadrons based on Ξ_K alone [4]. **(Testable)**

Suggested Platforms: BECs, Superconductors, Chiral Molecules

Controlled vortex manipulation in type-II superconducting films or ultracold BECs provides analogues for testing the quantized EMF impulse using SQUID magnetometers [2]. High-resolution attosecond spectroscopy offers a platform to probe the Swirl Clock's influence on chiral dynamics [6].

A.15 Experimental Testability and Falsifiable Setups

A.15.1 Swirl-Clock Dilation under Chiral Driving

SST predicts

$$\frac{dt_{\text{local}}}{dt_{\infty}} = \sqrt{1 - \frac{\|\mathbf{v}_{\odot}\|^2}{c^2}}.$$

Numerics: $\|\mathbf{v}_{\odot}\|/c \approx 3.6487 \times 10^{-3} \Rightarrow dt_{\text{local}}/dt_{\infty} \approx 0.9999933436$ (6.66×10^{-6} fractional dilation). **Protocol:** attosecond pump–probe on chiral targets; flip light helicity shot-to-shot; measure helicity-odd delay Δt . **Thresholds:** < 100 as pulses; 15–60 eV; avoid saturation. **Falsifier:** no helicity-odd component within error after fluence normalization.

A.15.2 $R \leftrightarrow T$ Transition Map in Knotted Coils

Drive a trefoil-like current path; track reconnection-induced topology flips in B-dot arrays and SQUID flux steps $\Delta\Phi = n\Phi_{\star}$. **Falsifier:** absence of discrete $\Delta\Phi$ statistics correlated with near-contact events.

A.15.3 Swirl-Pressure Force Proxy

$\mathbf{a}_{\text{eff}} = -\nabla p/\rho_f$ with $[\mathbf{a}] = \text{m s}^{-2}$. **Protocol:** 3-phase coil below a torsion balance plate; compare $F_z = \int_A \Delta p dA$ to CFD. **Falsifier:** null result tracking $\Delta p \rightarrow 0$.

Status. *Axiomatically required.* This section lists quantitative SST predictions with thresholds and protocols.

A.15.4 Swirl-Clock Time Dilation under Chiral Driving

SST predicts a local clock ratio

$$\frac{dt_{\text{local}}}{dt_{\infty}} = \sqrt{1 - \frac{\|\mathbf{v}_{\odot}\|^2}{c^2}} \tag{A.18}$$

under incompressible, inviscid conditions. Using $\|\mathbf{v}_{\odot}\| = 1.09384563 \times 10^6 \text{ m s}^{-1}$,

$$\frac{\|\mathbf{v}_{\odot}\|}{c} \approx 3.65 \times 10^{-3}, \quad \frac{dt_{\text{local}}}{dt_{\infty}} \approx 0.99999335,$$

a fractional dilation of $\sim 6.65 \times 10^{-6}$. **Test:** Attosecond pump–probe on chiral targets with counter-rotating circular polarizations; differential delay Δt must scale with (A.18) when the same fluence reorients vorticity. **Thresholds:** Pulse duration < 100 as; center frequency 15–60 eV; helicity flip every other shot; intensity tuned below target ionization saturation. **Observable:** helicity-odd delay $\Delta t_{\odot} - \Delta t_{\ominus} \neq 0$ at fixed fluence.

A.15.5 Quantized EM Impulse from Swirl Reconnection

For a closed swirl string undergoing reconnection, SST predicts discrete EM flux impulses

$$\Delta\Phi = n\Phi_{\star}, \quad n \in \mathbb{Z}, \tag{A.19}$$

coupled to knot topology. **Test:** SQUID loop near a driven knotted-conductor coil; detect telegraphic $\Delta\Phi$ coincident with reconnection signatures (high-speed B-dot probes). **Thresholds:** drive frequency near coil eigenmode; current ramp below thermal softening; low- T to reduce $1/f$ noise.

A.15.6 Swirl-Pressure Gravity Proxy

Effective gravitational acceleration in SST reduces to

$$\mathbf{a}_{\text{eff}} = -\frac{\nabla p}{\rho_f}, \quad [\mathbf{a}] = \text{m s}^{-2}, \quad (\text{A.20})$$

with $\rho_f = 7.0 \times 10^{-7} \text{ kg m}^{-3}$. **Test:** Table-top vertical force on a suspended plate above a 3-phase swirl coil: predict $F_z = \int_A \Delta p dA$ from CFD of the experimental geometry; **falsifier:** no force within error once $\Delta p(\text{phase}) \rightarrow 0$.

A.15.7 GW Sector Sanity Check (Propagation Speed)

Linearized transverse excitations must propagate at c within $|\delta c/c| < 10^{-15}$ to satisfy multi-messenger bounds. **Test:** See Sec. A.20 for a dispersion bound; **falsifier:** any measurable frequency-dependent GW delay exceeding that bound.

A.16 Running Couplings and Topological β -Functions

We propose

$$\mu \frac{dg_i}{d\mu} = -\frac{b_i}{2\pi} g_i^3 + \kappa_i \frac{\mathcal{H}(K) + \alpha \mathcal{C}(K)}{(\mu r_c)^2} g_i + \mathcal{O}(g_i^5),$$

with $\mathcal{H}$ helicity (dimensionless), $\mathcal{C}$ near-contacts (dimensionless count). As $\mu r_c \rightarrow \infty$ the SST correction decouples, recovering SM running (asymptotic freedom for SU(3), screening for U(1)). **Falsifier:** precision extractions of $g_i(\mu)$ show no universal $(\mu r_c)^{-2}$ tail.

A.17 Canonical Quantization of $\mathbf{v}_\odot$ and Directors

Fields. Incompressible $\mathbf{v}_\odot(\mathbf{x}, t)$ with $\nabla \cdot \mathbf{v}_\odot = 0$, directors $\hat{d}_a(\mathbf{x}, t)$ with $\hat{d}_a^\dagger \hat{d}_a = 1$. Hamiltonian density (prototype):

$$\mathcal{H} = \frac{\rho_f}{2} \|\mathbf{v}_\odot\|^2 + \frac{\lambda}{2} \sum_a \|\nabla \hat{d}_a\|^2 + \gamma (\nabla \times \mathbf{v}_\odot) \cdot \mathbf{S}(\hat{d}) + \mathcal{V}_{\text{topo}}.$$

Equal-time algebra. Let Π be conjugate to $\mathbf{v}_\odot$. With transverse projector $P_{ij} = \delta_{ij} - \partial_i \partial_j / \nabla^2$,

$$[v_i(\mathbf{x}), \Pi_j(\mathbf{y})] = i\hbar P_{ij} \delta^{(3)}(\mathbf{x} - \mathbf{y}), \quad [\hat{d}_a(\mathbf{x}), \hat{d}_b^\dagger(\mathbf{y})] = \delta_{ab} \delta^{(3)}(\mathbf{x} - \mathbf{y}).$$

Linearization yields transverse quanta $a_{\mathbf{k}, \lambda}$ and internal modes $b_{\mathbf{k}, r}^{(a)}$. **Falsifier:** measured dispersion violating the projector-enforced spectrum.

A.18 Mass Generation: Scalar Condensate vs. Topological Tension

A.18.1 Scalar Swirl Condensate (Higgs-analogue)

Introduce a real/complex scalar $S(x)$ with symmetry-breaking potential

$$V(S) = \lambda(|S|^2 - S_0^2)^2, \quad [\lambda] = 1, \quad [S] = \text{energy}^{1/2} \text{length}^{-3/2}$$

and Yukawa-like couplings to knot species Ψ_K ,

$$\mathcal{L} \supset |D_\mu S|^2 - V(S) - y_K S \Psi_K^\dagger \Psi_K + \text{h.c.},$$

producing $m_K = y_K S_0$ after $\langle S \rangle = S_0$. *Dimensional check:* $[y_K S_0] = \text{energy} \Rightarrow m_K$ has mass units. *Falsifier:* a single S_0 and $\{\Psi_K\}$ fail to fit leptons and quarks simultaneously.

A.18.2 Topological Tension Mechanism

Define a swirl tension scale from fluid pressure times core radius

$$\mathcal{T}_0 \equiv \rho_f \|\mathbf{v}_\odot\|^2 r_c, \quad [\mathcal{T}_0] = \text{N m}^{-1}.$$

Numerically, $\mathcal{T}_0 \approx 7.0 \times 10^{-7} \cdot (1.09384563 \times 10^6)^2 \cdot 1.40897017 \times 10^{-15} \text{ N m}^{-1} \approx 1.18 \times 10^{-9} \text{ N m}^{-1}$. Assuming a slender-tube knot of length $\mathcal{L}(K)$, with contact/helicity corrections,

$$m_K c^2 = \beta \mathcal{T}_0 \mathcal{L}(K) + \alpha \mathcal{C}(K) + \gamma \mathcal{H}(K),$$

with $\mathcal{C}$ dimension energy, $\mathcal{H}$ dimensionless. *Dimensional check:* $\mathcal{T}_0 \mathcal{L}$ is energy. *Falsifier:* a global fit fails across $(e, \mu, \tau, u, d, s, \dots)$ with one common $\{\alpha, \beta, \gamma\}$.

A.19 Charge and Quantum Numbers from Knot Data: Benchmark Table

Table A.2: SST benchmark predictions (prototype). Fill with full spectrum in revisions.

Species	Knot K	(χ, writhe, g)	$SU(3)$	$SU(2)$	Y	$Q = T_3 + \frac{Y}{2}$
e^-	trefoil 3_1	(χ_e, w_e, g_e)	1	1	-2	-1
u	5_2	(χ_u, w_u, g_u)	3	2	$+\frac{1}{3}$	$+\frac{2}{3}$
d	6_1	(χ_d, w_d, g_d)	3	2	$-\frac{1}{3}$	$-\frac{1}{3}$
γ	unknot	$(0, 0, 0)$	1	1	0	0

Falsifier. Any row where Q computed from the SST mapping disagrees with the SM value rules out that mapping.

A.20 Gravitational Waves and Lorentz-Breaking Bounds

Linearized transverse swirl obeys

$$\partial_t^2 \mathbf{u}_\perp - c^2 \nabla^2 \mathbf{u}_\perp + \eta_2 c^2 r_c^2 \nabla^4 \mathbf{u}_\perp + \dots = 0, \quad \Rightarrow \quad \omega^2 = c^2 k^2 (1 + \eta_2 (r_c k)^2 + \dots).$$

At $f = 100 \text{ Hz}$, $k = 2\pi f/c \approx 2.095 \times 10^{-6} \text{ m}^{-1}$, $r_c k \approx 2.953 \times 10^{-21}$, $(r_c k)^2 \approx 8.72 \times 10^{-42}$. GW170817 enforces $|v_g - c|/c \lesssim 10^{-15} \Rightarrow |\eta_2| (r_c k)^2 \lesssim 10^{-15}$, hence $|\eta_2| \lesssim 1.1 \times 10^{26}$ (very weak at LIGO band). **Implication:** high-frequency probes (if available) are needed; any measurable $v_g(f)$ beyond this expansion falsifies the linear SST wave sector.

A.21 Cosmological Embedding: Dark Matter and Dark Energy

A.21.1 Swirl-Foam Dark Matter

A gas of weakly radiative knotted excitations (“swirl foam”) with negligible pressure acts as DM: $w \approx 0$. Effective radial acceleration

$$a_r(r) = a_{\text{Newt}}(r) - \frac{1}{\rho_f} \partial_r p_{\text{foam}}(r).$$

Falsifier: inability to fit high-surface-brightness galaxies with one universal foam parameter.

A.21.2 Network-Tension Dark Energy

A percolating large-scale string network contributes $p_{\text{eff}} \simeq -\rho_{\text{tension}}$ ($w \approx -1 + \varepsilon$). **Falsifier:** SNe/BAO/CMB joint fits excluding any constant-tension component.

A.22 Cosmology: Baryogenesis and Dark Sector in SST

A.22.1 Chiral Swirl Condensation and Baryogenesis

SST permits chirality imbalance during early condensation. A minimal set of Sakharov-like conditions arises via (i) CP-odd helicity functional in $\mathcal{V}_{\text{topo}}$, (ii) out-of-equilibrium reconnection cascades, (iii) effective baryon number violation through knot fusion/splitting maps. **Falsifier:** no net leptogenesis/baryogenesis in kinetic simulations with realistic cooling ramps.

A.22.2 Dark Matter as Swirl Foam

Sub-radiative, weakly coupled knotted excitations form a *swirl foam* with equation of state $w \approx 0$. Predict rotation curves via

$$a_r(r) = a_{\text{Newt}}(r) - \frac{1}{\rho_f} \partial_r p_{\text{foam}}(r).$$

Falsifier: inability to fit high-surface-brightness galaxies with one universal foam parameter.

A.22.3 Dark Energy as Network Tension

A percolating string-network tension gives $p_{\text{eff}} \approx -\rho_{\text{tension}}$, i.e., $w \approx -1 + \varepsilon$. **Falsifier:** SNe/BAO/CMB fit excludes any constant-tension component.

A.23 Meta-Topology and Cosmogenetic Hierarchy in SST

A.23.1 Swirl Genesis and Knot Nucleation

We hypothesize that the early swirl condensate underwent a vacuum topology shift, spontaneously nucleating closed string loops—knotted excitations—through a stochastic tangle-unwinding process. Topological charge conservation implies metastable knot species K_i , which self-assemble into higher complexity via reconnection cascades and chirality bifurcation.

Initial condition: a high-temperature isotropic swirl field with $\|\mathbf{v}_\odot\|/c \sim \mathcal{O}(1)$, cooling into knotted sectors.

A.23.2 Golden-Ratio Mass Ladder

A recursive meta-knot hierarchy emerges via geometric self-similarity of swirl modes. The scaling law is governed by the *hyperbolic golden ratio*

$$\varphi_H = \frac{1 + \sqrt{5}}{2}, \quad \log m_n \propto n \log \varphi_H.$$

We propose the mass of the n -th family knot obeys:

$$m_n = m_0 \cdot \varphi_H^n,$$

with m_0 fixed by a benchmark species (e.g. electron or neutrino). This generates approximately exponential spacing in log-mass consistent with Standard Model families.

Falsifier: experimental mass ratios incompatible with any $\log m_n \propto n$ model, or inability to fit generations under any φ_H -based ladder.

A.23.3 Hierarchy from Contact and Helicity Correction

Total knot energy receives corrections from near-contact terms $\mathcal{C}(K)$ and helicity terms $\mathcal{H}(K)$, as in:

$$m_K c^2 = \beta \mathcal{T}_0 \mathcal{L}(K) + \alpha \mathcal{C}(K) + \gamma \mathcal{H}(K),$$

linking topological complexity directly to the mass scale.

Implication: The *meta-knot* tree acts as an ultraviolet completion for the mass hierarchy, replacing fine-tuned Yukawa couplings with a topological recursion.

A.23.4 Interpretation and Cosmological Implications

This cosmogenetic program frames knot species as frozen memory of early-universe swirl dynamics. Their distributions imprint both the dark sector (via knot abundance) and visible matter (via meta-hierarchy). Spontaneous symmetry breaking is topological, not field-theoretic.

Falsifier: Absence of any observable imprint of knot quantization in high-energy scattering or dark sector dynamics.

A.24 Numerical Program: Swirl–Knot Dynamics and EM Coupling

Algorithm. (1) Incompressible solver with Rankine-core regularization at r_c . (2) Vortex-line advection; adaptive remeshing; reconnection when separation $< \chi r_c$. (3) Topological diagnostics: helicity $\mathcal{H}$, writhe, linking; near-contacts $\mathcal{C}$. (4) EM coupling via Biot–Savart; predict SQUID steps $\Delta\Phi = n\Phi_*$. (5) Sweep drive helicity/frequency to map $R \leftrightarrow T$ transitions.

Dimensionless groups. $\text{He} = \|\mathbf{v}_\odot\|/c$, $\text{Kn} = r_c/L$, $\text{St} = \omega_{\text{drive}} L / \|\mathbf{v}_\odot\|$. **Falsifier:** failure to reproduce $\Delta\Phi$ quantization and helicity-odd delay maps.

A.25 Comparative Positioning

Table A.3: SST vs. alternative frameworks (sketch).

Feature	SST	LQG	Twistor	String Theory
Background	Flat + clock	Background-free	Conformal null	10D/AdS common
DoF	Swirl strings (hydro)	Spin networks	Twistors	1D strings
Mass origin	Topology/tension or S	Geometric/unknown	Model-dependent	Higgs + moduli
Testability	Lab hydro+EM	Cosmology/IR	Scattering	High scale

A.26 Comparison with Existing Frameworks

Scope of comparison. String theory’s unification program typically relies on extra dimensions and vibrational spectra of extended objects [19]. By contrast, SST seeks comparable reach using topological excitations in a single, incompressible medium. Our aim is not to displace established programs but to present a medium–based route that foregrounds concrete, falsifiable signatures [20].

SST vs. General Relativity: Reformulates Gravity via Swirl-Mediated Dynamics

Rather than postulating spacetime curvature as fundamental, SST models gravitation as an emergent kinematic effect arising from conserved fluid circulation in a flat background [3]. While GR attributes gravitational phenomena to metric curvature, SST *recovers* the same kinematic predictions—such as time dilation (Eq. A.1) and frame-dragging—through dynamics of a quantized swirl medium [1]. As shown in Fig. A.1, the same coupling constant $\mathcal{G}_\circ$ that governs electromagnetic impulse response also underlies gravitational emergence. When the system saturates,

$$G_{\text{swirl}} = \frac{v_\circ c^5 t^2}{2 F_\circ^{\text{max}} r_c^2}$$

converges numerically to G_N under the stated calibration, suggesting a shared topological origin.

SST vs. Standard Model: Topological Mass Spectrum, Emergent Gauge Structure

SST offers a topological and geometric framework that *recovers* several SM features. Fermion mass scaling arises from solitonic knot energies, reducing reliance on externally imposed Yukawa couplings [4]. Gauge structures and coupling ratios emerge from internal symmetry and elasticity within the swirl medium [4]. Quantum wave–particle duality and measurement are described dynamically via R $\leftrightarrow$ T phase transitions [1].

Analogy with Hydrodynamic Quantum Analogues

SST elevates empirically established phenomena—such as quantized circulation in superfluids [1]—to a universal principle, positing the vacuum as a structured, quantized superfluid. This connection underwrites the stability of swirl strings.

Connections to Emergent Gravity (Verlinde), Skymion Models, LET

The entropic interpretation of gravity aligns with Verlinde’s program [3, 7, 8, 18]. The use of topological solitons and emergent gauge fields parallels Skymion/Hopfion models [4, 15, 16]. Formally, SST resembles a modern Lorentz–Ether Theory (LET) due to its preferred time foliation, but seeks to justify this structure through parameter-light predictions grounded in topology [4].

A.27 Conclusion & Outlook

SST is offered as a *candidate* unification in which gravity, gauge fields, and matter *emerge* from the kinematics and topology of a single incompressible medium. The framework is constructed to *recover* established limits (relativistic time dilation, Maxwell induction, quantum interference) while supplying parameter-light, topologically anchored predictions that can be confronted with experiment.

Proposes Testable Dynamics Beyond Standard QFT

The theory delivers specific, numerically quantifiable predictions, notably quantized EMF impulses during vortex topological transitions [2] and chirality-dependent sign flips in attosecond delays [6]. These predictions make SST suitable for direct experimental appraisal in BECs, superconductors, and ultrafast spectroscopy.

Positioning SST Within the Theoretical Landscape

Rather than positioning itself in opposition to established theories, SST is framed as a candidate framework that **extends** their explanatory reach by grounding mass scaling, gauge structure, and gravitational phenomenology in a unified topological–fluid substrate. Its testable predictions and internal consistency motivate further study alongside broader efforts toward quantum gravity and unification.

Future Work

Future research aims to complete the particle spectrum and extend SST to cosmology [1].

1. **Neutrino sector from nested knots:** Analyze amphichiral multi-component links to derive neutrino masses and mixing [4].
2. **Loop entanglement entropy for black-hole analogues:** Establish thermodynamic analogues via entanglement in loop configurations [1].
3. **Cosmological inflation via swirl density fields:** Derive cosmic acceleration from global kinematics and vorticity variance Q_D ; aim to recover $w \approx -1$ from decay of length density $L(t)$ [1].

Bibliography

- [1] O. Iskandarani, *Swirl-string theory (sst) canon v0.5.9: Core postulates, constants, and boxed master equations*, Zenodo (2025).
- [2] O. Iskandarani, *Rotating-frame unification in the sst canon: From swirl density to swirl-emf, and a canonical derivation of the coupling $\mathcal{G}_\odot$* , Zenodo (2025).
- [3] O. Iskandarani, *Long-Distance Swirl Gravity from Chiral Swirling Knots with Central Holes*, Zenodo (2025).
- [4] O. Iskandarani, *A Hydrodynamic Lagrangian Framework for Swirl-String Theory*, Preprint (2025).
- [5] B. S. Deaver and W. M. Fairbank, *Experimental evidence for quantized flux in superconducting cylinders*, Phys. Rev. Lett. 7 (2), 43–46 (1961).
- [6] O. Iskandarani, *Chirality as Time Asymmetry: A Swirl-String Theory Interpretation of Attosecond Photoionization Delays*, Zenodo (2025).
- [7] E. Verlinde, *On the origin of gravity and the laws of newton*, JHEP 2011(4), 29 (2011).
- [8] E. P. Verlinde, *Emergent gravity and the dark universe*, SciPost Phys. 2(3), 016 (2017).
- [9] L. Onsager, *Statistical hydrodynamics*, Il Nuovo Cimento (Supplemento), 6, 279–287 (1949).
- [10] R. Doll and M. Näbauer, *Experimental Proof of Magnetic Flux Quantization in a Superconducting Ring*, Phys. Rev. Lett. 7, 51 (1961).
- [11] J. Clarke and A. I. Braginski (eds.), *The SQUID Handbook*, Wiley-VCH (2004).
- [12] L. Nahon et al., *Determining Molecular Chirality via Attosecond Photoionization Delay Measurements*, Nature Photonics 14, 491–496 (2020).
- [13] C. Barceló, S. Liberati, M. Visser, *Analogue Gravity*, Living Rev. Relativity 14, 3 (2011).
- [14] D. R. Scherer, C. N. Weiler, T. W. Neely, and B. P. Anderson, *Vortex Formation by Merging of Multiple Trapped Bose-Einstein Condensates*, Phys. Rev. Lett. 98, 110402 (2007).
- [15] T. H. R. Skyrme, *A Unified Field Theory of Mesons and Baryons*, Nucl. Phys. 31, 556–569 (1962).
- [16] N. S. Manton and P. Sutcliffe, *Topological Solitons*, Cambridge University Press (2004).
- [17] L. H. Kauffman, *Knots and Physics*, World Scientific (3rd ed., 2001).
- [18] T. Jacobson, *Thermodynamics of Spacetime: The Einstein Equation of State*, Phys. Rev. Lett. 75, 1260–1263 (1995).
- [19] L. Susskind, *The Anthropic Landscape of String Theory*, arXiv:hep-th/0302219 (2003).

- [20] S. Hossenfelder, *Lost in Math: How Beauty Leads Physics Astray* (Basic Books, 2018).
- [21] J. W. Alexander, A lemma on systems of knotted curves, *Proc. Natl. Acad. Sci. USA* **9**, 93–95 (1923).
- [22] A. A. Markov, Über die freie Äquivalenz geschlossener Zöpfe, *Rec. Math. [Mat. Sbornik]* N.S. **1(43)**, 73–78 (1935).
- [23] H. K. Moffatt, The degree of knottedness of tangled vortex lines, *J. Fluid Mech.* **35**, 117–129 (1969).
- [24] G. Călugăreanu, L’intégrale de Gauss et l’analyse des nœuds tridimensionnels, *Rev. Math. Pures Appl.* **4**, 5–20 (1959).
- [25] J. H. White, Self-linking and the Gauss integral in higher dimensions, *Am. J. Math.* **91**, 693–728 (1969).
- [26] F. B. Fuller, The writhing number of a space curve, *Proc. Natl. Acad. Sci. USA* **68**, 815–819 (1971).
- [27] E. Witten, Quantum field theory and the Jones polynomial, *Commun. Math. Phys.* **121**, 351–399 (1989).
- [28] Han, M., Ji, J.-B., Blech, A., Goetz, R. E., Allison, C., Greenman, L., Koch, C. P., Wörner, H. J., Attosecond control and measurement of chiral photoionization dynamics, *Nature*, 2025, doi:10.1038/s41586-025-09455-4, online 27 Aug 2025.
- [29] Krausz, Ferenc, Ivanov, Misha, Attosecond physics, *Rev. Mod. Phys.*, 81, 163–234, 2009, doi:10.1103/RevModPhys.81.163.
- [30] Beaulieu, S. et al., Attosecond-resolved photoionization of chiral molecules, *Science*, 358, 1288–1294, 2018, doi:10.1126/science.aao5624.
- [31] S. Lyu, V. Mathai, Y. Wang, B. Sobac, P. Colinet, D. Lohse, and C. Sun, “Final fate of a Leidenfrost droplet: Explosion or takeoff,” *Science Advances* **5**, eaav8081 (2019), doi:10.1126/sciadv.aav8081.
- [32] F. Celestini, T. Frisch, and Y. Pomeau, “Take Off of Small Leidenfrost Droplets,” *Phys. Rev. Lett.* **109**, 034501 (2012).
- [33] F. Celestini and G. Kirstetter, “Effect of an Electric Field on a Leidenfrost Droplet,” *arXiv:1203.4799* (2012).
- [34] Tong Zhou, “Identification of LFT Structured Descriptor Systems with Slow and Non-uniform Sampling,” *IEEE Transactions on Automatic Control*, 2025. arXiv:2407.00629
- [35] M. Büttiker, Larmor precession and the traversal time for tunneling, *Phys. Rev. B* **27**, 6178–6188 (1983). doi:10.1103/PhysRevB.27.6178
- [36] E. H. Hauge and J. A. Støvneng, Tunneling times: a critical review, *Rev. Mod. Phys.* **61**, 917–936 (1989). doi:10.1103/RevModPhys.61.917
- [37] H. G. Winful, Tunneling time, the Hartman effect, and superluminality: A proposed resolution, *Physics Reports* **436**(1-2), 1–69 (2006). doi:10.1016/j.physrep.2006.09.002
- [38] M. Gell-Mann, A Schematic Model of Baryons and Mesons, *Phys. Lett.* **8**(3), 214–215 (1964). doi:10.1016/S0031-9163(64)92001-3

- [39] G. Zweig, An $SU(3)$ model for strong interaction symmetry and its breaking, CERN Reports, 8182/TH.401 (1964). Preprint.
- [40] M. P. Brenner, S. Hilgenfeldt, and D. Lohse, Single-bubble sonoluminescence, *Rev. Mod. Phys.* **74**(2), 425–484 (2002). doi:10.1103/RevModPhys.74.425
- [41] Krzysztof K. Putyra and Paul Turner, “Unknotting Two Knots by Five Band Moves,” arXiv preprint arXiv:2408.04510, 2024. <https://arxiv.org/abs/2408.04510>
- [42] Saffman, P. G., *Vortex Dynamics*, Cambridge Univ. Press, 1992.
- [43] Batchelor, G. K., *An Introduction to Fluid Dynamics*, Cambridge Univ. Press, 1967.
- [44] Landau, L. D. and Lifshitz, E. M., *Fluid Mechanics*, Course of Theoretical Physics 6, Pergamon, 1987.
- [45] Planck, M., “On the Law of Distribution of Energy in the Normal Spectrum,” *Ann. Phys.*, 4, 553–563, 1901.
- [46] Einstein, A., “On a Heuristic Point of View...,” *Ann. Phys.*, 17, 132–148, 1905.
- [47] Salmon, R., “Hamiltonian Fluid Mechanics,” *Annu. Rev. Fluid Mech.*, 20, 225–256, 1988.
- [48] Morrison, P. J., “Hamiltonian Description of the Ideal Fluid,” *Rev. Mod. Phys.*, 70(2), 467–521, 1998.
- [49] de Sitter, Willem, “On the relativity of inertia: Remarks concerning Einstein’s latest hypothesis,” *Proc. Roy. Neth. Acad. Arts Sci.* **19**, 1217–1225 (1917).
- [50] Weinberg, Steven, “The cosmological constant problem,” *Rev. Mod. Phys.* **61**(1), 1–23 (1989). doi:10.1103/RevModPhys.61.1
- [51] C. N. Yang and R. L. Mills, “Conservation of Isotopic Spin and Isotopic Gauge Invariance,” *Phys. Rev.* **96**, 191–195 (1954). doi:10.1103/PhysRev.96.191
- [52] M. E. Peskin and D. V. Schroeder, *An Introduction to Quantum Field Theory*, Westview Press, 1995. ISBN: 978-0201503975
- [53] S. Weinberg, *The Quantum Theory of Fields, Vol. 2: Modern Applications*, Cambridge University Press, 1996. doi:10.1017/CBO9781139644174
- [54] M. Gell-Mann, “Symmetries of Baryons and Mesons,” *Phys. Rev.* **125**, 1067–1084 (1962). doi:10.1103/PhysRev.125.1067
- [55] H. Georgi, *Lie Algebras in Particle Physics*, 2nd ed., Westview Press, 1999. ISBN: 978-0738202334
- [56] B. P. Abbott et al., “GW170817: Observation of Gravitational Waves from a Binary Neutron Star Inspiral,” *Phys. Rev. Lett.* **119**, 161101 (2017). doi:10.1103/PhysRevLett.119.161101
- [57] K. Popper, *The Logic of Scientific Discovery*, Hutchinson, 1959.
- [58] D. Gross and F. Wilczek, “Ultraviolet Behavior of Non-Abelian Gauge Theories,” *Phys. Rev. Lett.* **30**, 1343 (1973). doi:10.1103/PhysRevLett.30.1343
- [59] H. D. Politzer, “Reliable Perturbative Results for Strong Interactions?” *Phys. Rev. Lett.* **30**, 1346 (1973). doi:10.1103/PhysRevLett.30.1346

- [60] C. G. Callan and K. Symanzik, “Infrared and Ultraviolet Behavior of Green’s Functions,” *Phys. Rev. D* **2**, 1541 (1970).
- [61] P. A. M. Dirac, *Lectures on Quantum Mechanics*, Yeshiva University, 1964.
- [62] S. Weinberg, *The Quantum Theory of Fields, Vol. I*, Cambridge University Press, 1995.
- [63] P. W. Higgs, “Broken Symmetries and the Masses of Gauge Bosons,” *Phys. Rev. Lett.* **13**, 508 (1964). doi:10.1103/PhysRevLett.13.508
- [64] S. O. Bilson-Thompson, “A Topological Model of Composite Preons,” arXiv:hep-ph/0503213 (2005).
- [65] A. D. Sakharov, “Violation of CP Invariance, C Asymmetry, and Baryon Asymmetry of the Universe,” *JETP Lett.* **5**, 24 (1967).

.1 Full Canonical Lagrangian for SST

We work on Minkowski space with metric $\eta_{\mu\nu} = \text{diag}(-, +, +, +)$ and Levi-Civita tensor $\varepsilon^{0123} = +1$. A scalar clock field Φ defines

$$X \equiv \eta^{\alpha\beta} \partial_\alpha \Phi \partial_\beta \Phi, \quad u_\mu \equiv \frac{\partial_\mu \Phi}{\sqrt{-X}}, \quad P^{\mu\nu} \equiv \eta^{\mu\nu} + u^\mu u^\nu,$$

so that $u^\mu u_\mu = -1$ and $P^{\mu\nu} u_\nu = 0$. The total action splits into sectoral pieces:

$$S = \int d^4x \mathcal{L} = \int d^4x \left(\mathcal{L}_\Phi + \mathcal{L}_B + \mathcal{L}_{\text{YM}} + \mathcal{L}_{\text{mat}} + \mathcal{L}_{\text{top}} + \mathcal{L}_{\text{bridge}} \right). \quad (21)$$

(i) **Khronon / clock sector $\mathcal{L}_\Phi$** [4].

$$\mathcal{L}_\Phi = \frac{M_\Phi^2}{2} \left[c_1 (\partial_\mu u_\nu) (\partial^\mu u^\nu) + c_2 (\partial_\mu u^\mu)^2 + c_3 (\partial_\mu u_\nu) (\partial^\nu u^\mu) + c_4 (u^\mu \partial_\mu u_\nu) (u^\alpha \partial_\alpha u^\nu) \right] + \lambda (u^\mu u_\mu + 1), \quad (22)$$

where λ enforces $u^2 = -1$. We adopt the Canon constraint $c_{13} \equiv c_1 + c_3 = 0$ to yield a luminal tensor mode (GW170817 consistency).

(ii) **Two-form (vorticity) sector $\mathcal{L}_B$ and topological density $\mathcal{L}_{\text{top}}$** [4]. With an antisymmetric potential $B_{\mu\nu} = -B_{\nu\mu}$ and $H_{\mu\nu\rho} \equiv 3 \partial_{[\mu} B_{\nu\rho]}$,

$$\mathcal{L}_B = -\frac{\kappa_B}{12} H_{\mu\nu\rho} H^{\mu\nu\rho}, \quad \mathcal{L}_{\text{top}} = \frac{\vartheta_B}{24} \varepsilon^{\mu\nu\rho\sigma} H_{\mu\nu\alpha} H_{\rho\sigma}{}^\alpha. \quad (23)$$

Here $\kappa_B > 0$ ensures stability; ϑ_B multiplies a (parity-odd) topological density that is a total derivative classically but controls global sectors and selection rules.

(iii) **Emergent gauge (swirl connection) sector $\mathcal{L}_{\text{YM}}$** [1, ?]. For $A_\mu \in \mathfrak{g}$ with curvature $G_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu + g[A_\mu, A_\nu]$,

$$\mathcal{L}_{\text{YM}} = -\frac{\kappa_{\text{YM}}}{4} \text{Tr}(G_{\mu\nu} G^{\mu\nu}) + \frac{\theta_{\text{YM}}}{32\pi^2} \text{Tr}(G_{\mu\nu} \tilde{G}^{\mu\nu}), \quad \tilde{G}^{\mu\nu} \equiv \frac{1}{2} \varepsilon^{\mu\nu\rho\sigma} G_{\rho\sigma}. \quad (24)$$

$\kappa_{\text{YM}} > 0$ and the θ_{YM} term is topological (CP-odd) at the classical level.

(iv) **Matter (knot-soliton) sector $\mathcal{L}_{\text{mat}}$** [4]. With Dirac field ψ and $D_\mu = \partial_\mu + igA_\mu$,

$$\mathcal{L}_{\text{mat}} = \bar{\psi} (i\gamma^\mu D_\mu - m_K^{(\text{sol})}) \psi, \quad m_K^{(\text{sol})} = \mathcal{M}_0 \Xi_K \text{ (Eq. A.3)}. \quad (25)$$

Couplings to the preferred frame may be restricted to the spatial leaves via $P^\mu{}_\nu D_\mu$ when required by foliation symmetry.

(v) **Swirl-electromagnetic bridge $\mathcal{L}_{\text{bridge}}$ (modified Faraday)** [2]. Let $F_{\mu\nu}$ be the emergent U(1) field strength. To make the source covariant and foliation-aware we take a spacelike unit field n^μ orthogonal to u^μ (i.e., $n^\mu u_\mu = 0$, $n^\mu n_\mu = +1$) built from the vorticity sector (e.g., a principal direction of H).

$$\mathcal{L}_{\text{bridge}} = -\frac{\mathcal{G}_\odot}{2} n_\alpha \varepsilon^{\alpha\mu\nu\rho} (\partial_\mu \rho_\odot) F_{\nu\rho}, \quad (26)$$

which is U(1) gauge invariant. Projecting the equations of motion onto the spatial leaves (P) yields, in 3+1 form,

$$\nabla \times \mathbf{E} = -\partial_t \mathbf{B} - \mathbf{b}_\odot, \quad b_\odot^i = \mathcal{G}_\odot \partial_t \rho_\odot n^i \text{ (Sec. A.9).}$$

Dimensional consistency (SI). $[\mathcal{L}] = \text{J m}^{-3}$; $[\mathcal{G}_\mathcal{O}] = \text{Wb}$; $\rho_\mathcal{O}$ is an areal swirl density, so (26) carries energy-density units after contraction with $F_{\mu\nu}$.

Canon checks. (i) $c_{13} = 0$ in (22) (GW170817 consistency); (ii) $\kappa_B, \kappa_{\text{YM}} > 0$ (stability); (iii) R-phase (unknot) implies $\rho_\mathcal{O} \rightarrow 0$ so $\mathcal{L}_{\text{bridge}} \rightarrow 0$ (standard Faraday recovered); (iv) masses follow Eq. A.3 with $\Xi_{3_1} = 1 \Rightarrow \mathcal{M}_0 = m_e$ after calibration.

Analogy (brief). Four Lego plates (clock, vorticity, gauge, matter) lock together; a slim adapter brick (the bridge term) lets the swirl plate drive the EM plate.

.2 Electroweak mixing and charge operator

Define $A_\mu = s_w W_\mu^3 + c_w B_\mu$ and $Z_\mu = c_w W_\mu^3 - s_w B_\mu$ with $s_w = \sin \theta_w$, $c_w = \cos \theta_w$. Then $Q = T_3 + Y$ and the photon couples as $e Q A_\mu$ with $e = g_2 s_w = g_1 c_w$.

.3 Anomaly constraints and hypercharge calibration

Impose $[\text{SU}(3)]^2 - U(1)$, $[\text{SU}(2)]^2 - U(1)$, $[U(1)]^3$, and gravitational- $U(1)$ anomalies to vanish generation by generation. Solve for $(\alpha_Y, \beta_Y, \gamma_Y)$ given a seed identification (electron trefoil, up/down knot types). This fixes $Y(K)$ for all K and thereby predicts Q .

.4 Representative knot $\rightarrow$ rep assignments

Provide a table listing $(K, L, S, \chi, \text{wr}, g) \mapsto (R_3, R_2, Y, Q)$ for the benchmark set.

.5 Swirl Cushion Stability and Leidenfrost Analogy

Canonical Context

In Swirl-String Theory (SST), short-range repulsion and levitation arise from pressure wells sustained by the outer irrotational envelope of a swirl string. The stability of this envelope determines whether a structure “floats” above its substrate or collapses into direct contact. An experimentally accessible analogue exists in the Leidenfrost effect, where a liquid droplet levitates on its own vapor cushion. The breakdown of that film, leading either to smooth takeoff or explosive contact, maps directly onto the stability limits of the SST swirl cushion.

.5.1 Empirical Leidenfrost Scaling

Experiments by Lyu *et al.* [31] demonstrate that a Leidenfrost droplet of radius R transitions from stable levitation to explosion at a critical size $R_\times$ that depends on the surface contaminant fraction ϕ as

$$\boxed{\frac{R_\times}{R_0} \propto \phi^{-1/3}}, \quad (27)$$

with R_0 a reference radius. Clean, small droplets ($\phi \rightarrow 0$) lift off smoothly, whereas contaminated or larger droplets collapse violently. This one-third power law defines a dimensionless invariant that can be translated into the swirl-string formalism.

.5.2 SST Translation: Pressure–Film Criterion

In SST, the levitating envelope is modeled as a thin region where the swirl–pressure difference

$$\Delta p_{\text{film}} \simeq \rho_f \|\mathbf{v}_\odot\|^2 \Theta(\phi, R) \quad (28)$$

balances the combined capillary and hydrostatic loads,

$$\Delta p_{\text{film}} \gtrsim \frac{\gamma}{R} + \rho_\ell g R. \quad (29)$$

Here $\Theta(\phi, R) \in (0, 1]$ is an effective *transmissivity* representing partial blockage of swirl flow through the thin layer. When contaminants or phase disorder increase ϕ , Θ decreases and the cushion fails once Eq. (29) can no longer be satisfied.

Combining Eqs. (27)–(29) yields a canonical stability condition

$$\Theta(\phi, R) \simeq 1 - \kappa \left(\phi \frac{R}{R_0} \right)^m, \quad m \simeq 1, \quad (30)$$

so that the empirical 1/3 exponent corresponds to the point where Δp_{film} drops below the restoring swirl–pressure $\rho_f \|\mathbf{v}_\odot\|^2$.

.5.3 Numerical Consistency

Using canonical constants $\rho_f = 7.0 \times 10^{-7} \text{ kg m}^{-3}$ and $\|\mathbf{v}_\odot\| = 1.09384563 \times 10^6 \text{ m s}^{-1}$, one obtains

$$\rho_f \|\mathbf{v}_\odot\|^2 \approx 8.4 \times 10^5 \text{ Pa},$$

which exceeds typical capillary pressures ($\gamma/R \sim 70 \text{ Pa}$ for $R = 1 \text{ mm}$) by four orders of magnitude. Thus a reduction of Θ below 10^{-3} suffices to collapse the swirl cushion, consistent with the measured instability threshold.

.5.4 Interpretation and Analogy

The vapor film of a Leidenfrost drop functions as a miniature *swirl cushion*: it stores mechanical energy in a confined, pressurized layer sustained by circulation. Contamination or local disorder reduces the transmissivity Θ , analogous to phase decoherence in a swirl envelope. When Θ falls below the critical value set by Eq. (29), the envelope implodes, releasing the stored pressure in a rapid outburst. The same criterion governs collapse of larger SST structures or levitating devices when the outer swirl envelope becomes phase–clogged or thermally decoherent.

Physical Analogy

A Leidenfrost droplet is like a hovercraft riding on a swirl of vapor. Clean air holes keep it afloat; a few crumbs block them, and it crashes.

.6 Identification with Slow and Non-Uniform Sampling (LFT–DAE Form)

Model. We represent a linearized SST rig as a continuous-time descriptor system

$$E \dot{x}(t) = A(\theta) x(t) + B(\theta) u(t), \quad (31)$$

$$y(t) = C(\theta) x(t) + D(\theta) u(t), \quad (32)$$

with parameters $\theta \in \Theta$ entering through a linear fractional transformation (LFT):

$$H(s, \theta) = \mathcal{F}_l(M(s), \Delta(\theta)),$$

where H is the transfer function matrix (TFM). We assume well-posedness, stability, and a known upper bound on the settling time t_s .

Key relation (from steady states). For inputs u generated by a stable LTI generator $G(s)$, the system response admits an explicit transient/steady-state decomposition. The steady-state segments determine, in principle, finite sets of linear constraints on $\{H(s_k, \theta), \partial_s H(s_k, \theta)\}$ and right-tangential data $\{H(s_k, \theta)v_k\}$ at user-selected points $s_k \in \mathbb{C}$.²

Estimator. Choose $\{s_k, v_k\}$ and form the least-squares problem

$$\min_{\theta} \sum_k \left\| \begin{bmatrix} \hat{H}(s_k) - H(s_k, \theta) \\ \partial_s \hat{H}(s_k) - \partial_s H(s_k, \theta) \\ \hat{H}(s_k)v_k - H(s_k, \theta)v_k \end{bmatrix} \right\|_2^2 + \lambda \Phi_{\text{SST}}(\theta),$$

where hats denote steady-state-derived estimates and Φ_{SST} imposes SST constraints (dimensionality, passivity at $s = 0$, bounds using $F_{\text{swirl}}^{\max}$, r_c , ρ_f). This avoids nonconvex direct time-fit traps and is valid for non-uniform, sub-Nyquist sampling.

Targets. Include the swirl-EMF coupling $\mathcal{G}_{\text{S}}$ within θ ; estimate it primarily from $s = 0$ and low- $|s|$ constraints, then cross-check on withheld s_k .

.7 Kappa-Event Dwell-Time Experiment (Larmor-Clock Protocol)

We outline the complete protocol for measuring the dwell time associated with Kappa events. The setup, derived from the Larmor-clock technique of Büttiker [35], measures the precession angle θ of a spin-polarized particle in a weak magnetic field B localized within the barrier. The dwell time is $\tau_d = \theta/\omega_L$, with $\omega_L = g\mu_B B/\hbar$.

SST prediction. Each successful traversal corresponds to a Kappa event, a phase-coherent alignment propagating through the barrier at speed $u_{\kappa} = \|\mathbf{v}_{\text{C}}\|/\Xi$. Hence

$$\tau_d^{(\text{SST})}(L) = \frac{L}{u_{\kappa}} = \frac{L}{\|\mathbf{v}_{\text{C}}\|} \Xi. \quad (33)$$

Using the canonical constants gives $\tau_d^{(\text{SST})} \approx \{4.6, 18.3, 45.7\}$ fs for $L = \{5, 20, 50\}$ nm.

Standard-QM benchmark. For comparison, a rectangular barrier of height $(V_0 - E) = 0.2$ eV predicts a Hartman delay $\tau_H \approx 3.3$ fs, nearly independent of L .

Observable falsifier. Measure $\tau_d(L)$ via the Larmor precession angle and extract the slope $s = d\tau_d/dL$. SST requires $s = \Xi/\|\mathbf{v}_{\text{C}}\|$ (constant, positive), while standard QM gives $s \approx 0$. The Kappa transition rate follows as $\Gamma_{\kappa} = 1/(s r_c)$, yielding $\Gamma_{\kappa} = 7.76 \times 10^{20} \text{ s}^{-1}$ and confirming $\Gamma_{\kappa} t_c = 1$ for the core time $t_c = r_c/\|\mathbf{v}_{\text{C}}\| = 1.29 \times 10^{-21}$ s.

Extensions. Slow acoustic or optical modulation of the barrier at frequency $f \sim 10$ MHz may be used to gate Kappa events on/off without altering τ_d , providing a phase-locked demonstration of causal alignment.

²See Zhou (2025) for explicit formulas and consistency conditions.

.8 Variational Derivation of the Clock–Rescaled Schrödinger Equation

Setup. Let the medium be incompressible and inviscid; denote the swirl velocity by $\mathbf{v}(\mathbf{x})$, vorticity $\boldsymbol{\omega} = \nabla \times \mathbf{v}$, and the SST clock speed by $|\mathbf{v}_\odot|$. Consider the local surrogate energy density for an isolated filament core [42, 43, 44]:

$$\mathcal{H}_{\text{swirl}}(\mathbf{x}) = \frac{1}{2} \rho_f |\mathbf{v}|^2 + \frac{1}{2} \rho_f r_c^2 |\boldsymbol{\omega}|^2, \quad [\mathcal{H}_{\text{swirl}}] = \text{J m}^{-3}. \quad (34)$$

Define the proper-time factor by kinematics

$$\tau(\mathbf{x}) = \sqrt{1 - \frac{|\mathbf{v}_\perp(\mathbf{x})|^2}{|\mathbf{v}_\odot|^2}} \in [0, 1], \quad (35)$$

with $\mathbf{v}_\perp$ the tangential speed on the swirl sheet (Hamiltonian-fluid kinematics [47, 48]).

Variational principle and generator rescaling. Let Ψ be a mode supported by the filamental region V with Hermitian operator $H[\mathbf{v}]$ constructed from $\mathcal{H}_{\text{swirl}}$. Local proper time satisfies $dt_{\text{loc}} = \tau(\mathbf{x}) dt_\infty$. The action

$$S[\Psi] = \int dt_\infty \int_V \left(\frac{i\hbar}{2} \tau \Psi^\dagger \overset{\leftrightarrow}{\partial}_{t_\infty} \Psi - \tau \Psi^\dagger H[\mathbf{v}] \Psi \right) d^3x$$

yields, upon variation in $\Psi^\dagger$,

$$i\hbar \partial_{t_\infty} \Psi(\mathbf{x}, t_\infty) = \tau(\mathbf{x}) H[\mathbf{v}] \Psi(\mathbf{x}, t_\infty). \quad (36)$$

For a single collective coordinate, average τ over the core: $\langle \tau \rangle = V^{-1} \int_V \tau d^3x$, giving $i\hbar \dot{\Psi} = \langle \tau \rangle H \Psi$.

Dimensional and limit checks. The argument $|\mathbf{v}_\perp|^2/|\mathbf{v}_\odot|^2$ is dimensionless; the prefactor τ multiplies an energy operator H and therefore rescales the frequency (clock) as expected. For $|\mathbf{v}_\perp| \ll |\mathbf{v}_\odot|$, $\tau \simeq 1 - \frac{|\mathbf{v}_\perp|^2}{2|\mathbf{v}_\odot|^2}$ and the fractional frequency shift is $\Delta\nu/\nu \simeq -\langle |\mathbf{v}_\perp|^2 \rangle / (2|\mathbf{v}_\odot|^2)$.

Remarks. This is the Hamiltonian-fluid realization of the SST clock law compatible with [47, 48] and with local vortex energetics [42, 43, 44].

.9 Spectral Thermodynamics: Density of States and Core Regularization

Model. Let the near-core energy density localize geometrically as $u_0(r) = \frac{1}{2} \rho_f |\mathbf{v}_\odot|^2 (r_c^2/r^2)$ and define the natural frequency scale $\Omega_c = |\mathbf{v}_\odot|/r_c$ ($[\Omega_c] = \text{s}^{-1}$). Assume a bosonic density of states akin to Planck radiation with calibration factor λ_{BE} :

$$u_{\text{vortex}}(r, \omega, T) = u_0(r) \lambda_{BE} \left(\frac{\omega}{\Omega_c} \right)^3 \frac{1}{e^{\hbar\omega/k_B T} - 1}. \quad (37)$$

Integration over frequency. Using $x = \hbar\omega/(k_B T)$ and $\int_0^\infty \frac{x^3}{e^x - 1} dx = \pi^4/15$ (Planck’s law [45]),

$$u_{\text{tot}}(r, T) = u_0(r) \lambda_{BE} \frac{\pi^4}{15} \left(\frac{k_B T}{\hbar \Omega_c} \right)^4. \quad (38)$$

Thermodynamic closures (massless-mode gas) give $P = \frac{1}{3} u_{\text{tot}}$, $s = \frac{4}{3} u_{\text{tot}}/T$, with T^4 and T^3 scaling.

Radial and far-field behavior. Near core, $u_{\text{tot}} \propto r^{-2}$. For a straight filament of length L in a cylinder $\xi \leq r \leq R$,

$$\frac{E}{L} = \int_{\xi}^R u_{\text{tot}} 2\pi r dr = 2\pi u_0(\xi) \lambda_{BE} \frac{\pi^4}{15} \left(\frac{k_B T}{\hbar \Omega_c} \right)^4 \ln \frac{R}{\xi},$$

exhibiting the classical $\ln(R/\xi)$ dependence of vortex energetics [42, 43, 44].

Dimensional check. u_0 has units J m^{-3} , $(\omega/\Omega_c)^3$ is dimensionless, and the Bose–Einstein factor is dimensionless; hence u_{vortex} and u_{tot} are J m^{-3} .

Numerical consistency (room temperature). With $|\mathbf{v}_{\odot}| = 1.09384563 \times 10^6 \text{ m s}^{-1}$, $r_c = 1.40897017 \times 10^{-15} \text{ m}$, $\rho_f = 7.0 \times 10^{-7} \text{ kg m}^{-3}$, $\Omega_c \simeq 7.763 \times 10^{20} \text{ s}^{-1}$. At $T = 300 \text{ K}$ and $\lambda_{BE} = 1$, $u_0(r_c) = \frac{1}{2} \rho_f |\mathbf{v}_{\odot}|^2 \approx 4.19 \times 10^5 \text{ J m}^{-3}$, but $u_{\text{tot}}(r_c, 300 \text{ K}) \approx 1.78 \times 10^{-23} \text{ J m}^{-3}$, showing strong thermal suppression because $k_B T \ll \hbar \Omega_c$.

References. Classical vortex energetics and logarithmic line energy: [42, 43, 44]. Quantum thermodynamic integral: [45].

.10 Photoelectric Mapping in SST Units: Geometry and Units

Ring-emission geometry. Model a vortex “photon” as a translating ring of circumference $2\pi r_c$ that couples one full swirl rotation per half-turn of its transverse frame, giving a 4π geometric factor (solid-angle coverage of a closed framed loop). The impulse scale combines (i) the EM/swirl maximum boundary force $F_{\text{swirl}}^{\text{max}}$ and (ii) the core size r_c set by the elastic boundary.

SST action scale and photoelectric slope. Define

$$\boxed{h_{\text{SST}} = 4\pi \frac{F_{\text{swirl}}^{\text{max}} r_c^2}{|\mathbf{v}_{\odot}|}} \quad [\text{J s}], \quad E_{\text{max}} = h_{\text{SST}} f - \Phi, \quad f_0 = \Phi/h_{\text{SST}}. \quad (39)$$

Units: $F \times r_c^2/|\mathbf{v}| = (\text{N})(\text{m}^2)/(\text{m s}^{-1}) = \text{J s}$.

Numerical calibration. With $F_{\text{swirl}}^{\text{max}} = 29.053507 \text{ N}$, $r_c = 1.40897017 \times 10^{-15} \text{ m}$, $|\mathbf{v}_{\odot}| = 1.09384563 \times 10^6 \text{ m s}^{-1}$,

$$h_{\text{SST}} = 4\pi \frac{(29.053507) (1.40897017 \times 10^{-15})^2}{1.09384563 \times 10^6} = 6.6260695 \times 10^{-34} \text{ J s},$$

numerically matching CODATA $h = 6.62607015 \times 10^{-34} \text{ J s}$ to within $< 10^{-40}$ in absolute units. Hence the photoelectric slope is recovered while intensity controls current (not slope) as in Einstein’s law [46].

Discussion. The equality $h_{\text{SST}} \approx h$ is an output of the SST constants ($F_{\text{swirl}}^{\text{max}}, r_c, |\mathbf{v}_{\odot}|$), not an input. The factor 4π is geometric (closed framed loop), and the units/limits align with the standard photoelectric phenomenology [46].

.11 Derivation of the Local Swirl Stress Tensor

Lagrangian and Euler–Lagrange form. Use the local surrogate (incompressible) Lagrangian density

$$\mathcal{L}_{\text{swirl}} = \frac{1}{2}\rho_f |\mathbf{v}|^2 - \frac{1}{2}\rho_f r_c^2 |\boldsymbol{\omega}|^2, \quad \boldsymbol{\omega} = \nabla \times \mathbf{v}. \quad (40)$$

This is the lowest-order local truncation of the Biot–Savart nonlocal energy [42, 43, 44]. Stationarity under $\mathbf{v} \mapsto \mathbf{v} + \delta\mathbf{v}$ yields the vector Helmholtz form (for incompressible fields): $\nabla^2 \mathbf{v} + \mathbf{v}/r_c^2 = 0$.

Symmetric Cauchy stress. By standard Noether/Belinfante symmetrization for spatial translations, the local, symmetric stress consistent with $\mathcal{L}_{\text{swirl}}$ reads

$$\sigma_{ij} = \rho_f v_i v_j + \rho_f r_c^2 \left(\omega_i \omega_j - \frac{1}{2} \delta_{ij} |\boldsymbol{\omega}|^2 \right) - \delta_{ij} \frac{1}{2} \rho_f |\mathbf{v}|^2 \quad [\text{Pa}], \quad (41)$$

cf. vortex-tension analogies ($\propto \omega_i \omega_j$) in classical fluids [42, 43, 44].

Checks. (i) *Dimensions:* $\rho_f v_i v_j$ and $\rho_f r_c^2 \omega_i \omega_j$ are $\text{kg m}^{-1} \text{s}^{-2} = \text{Pa}$. (ii) *Rigid rotation:* for $\mathbf{v} = \boldsymbol{\Omega} \times \mathbf{r}$, $\boldsymbol{\omega} = 2\boldsymbol{\Omega}$, the tensor splits into kinetic (Reynolds) and “swirl-tension” parts. (iii) *Radial scaling:* near a thin filament with $v_\theta \propto r^{-1}$ outside the core, traction on a probe scales as r^{-2} , consistent with classical line-vortex behavior [42].

Order-of-magnitude at $r=r_c$. Using $|\mathbf{v}_\odot| = 1.09384563 \times 10^6 \text{ m s}^{-1}$, $\rho_f = 7.0 \times 10^{-7} \text{ kg m}^{-3}$, and a representative tangential speed $v_\theta \approx |\mathbf{v}_\odot|/e \approx 4.02 \times 10^5 \text{ m s}^{-1}$, the dynamic pressure term is $\frac{1}{2}\rho_f v_\theta^2 \approx 5.67 \times 10^4 \text{ Pa}$. With $\omega \sim 2v_\theta/r_c$, the swirl-tension magnitude $\rho_f r_c^2 \omega^2 \approx 4.53 \times 10^5 \text{ Pa}$, showing that the ω -tension dominates near core—consistent with line-energy localization [42, 43, 44].

References. Classical derivations and analogies: [42, 43, 44]. Hamiltonian structure background: [47, 48].

.12 Benchmark Knot–Representation Table

Table 4: Benchmark map $(K, L, S, \chi, \text{wr}, g) \mapsto (R_3, R_2, Y, Q)$ with $Y = \alpha_Y \chi + \beta_Y \text{wr} + \gamma_Y g$ and $(\alpha_Y, \beta_Y, \gamma_Y) = (\frac{2}{3}, 0, -\frac{1}{2})$. Electric charge $Q = T_3 + Y$ with $T_3 = \pm\frac{1}{2}$ for $R_2 = \mathbf{2}$. Lepton candidate K_ℓ has $(L, S, \chi, \text{wr}, g) = (6, 3, 0, 0, 1)$; quark candidate K_q has $(4, 3, +1, 0, 1)$.

Species	K	L	S	χ	wr	g	R_3	R_2	Y	Q
ν_{eL}	K_ℓ	6	3	0	0	1	1	2	$-\frac{1}{2}$	$+\frac{1}{2} + (-\frac{1}{2}) = 0$
e_L	K_ℓ	6	3	0	0	1	1	2	$-\frac{1}{2}$	$-\frac{1}{2} + (-\frac{1}{2}) = -1$
$u_L^{(r)}$	K_q	4	3	+1	0	1	3	2	$+\frac{1}{6}$	$+\frac{1}{2} + (+\frac{1}{6}) = +\frac{2}{3}$
$u_L^{(g)}$	K_q	4	3	+1	0	1	3	2	$+\frac{1}{6}$	$+\frac{1}{2} + (+\frac{1}{6}) = +\frac{2}{3}$
$u_L^{(b)}$	K_q	4	3	+1	0	1	3	2	$+\frac{1}{6}$	$+\frac{1}{2} + (+\frac{1}{6}) = +\frac{2}{3}$
$d_L^{(r)}$	K_q	4	3	+1	0	1	3	2	$+\frac{1}{6}$	$-\frac{1}{2} + (+\frac{1}{6}) = -\frac{1}{3}$
$d_L^{(g)}$	K_q	4	3	+1	0	1	3	2	$+\frac{1}{6}$	$-\frac{1}{2} + (+\frac{1}{6}) = -\frac{1}{3}$
$d_L^{(b)}$	K_q	4	3	+1	0	1	3	2	$+\frac{1}{6}$	$-\frac{1}{2} + (+\frac{1}{6}) = -\frac{1}{3}$

.13 Swirl–Electromagnetic Unit Bridge Table

Swirl Quantity	EM Analogue	Symbol	Units	Meaning
Swirl field	Electric field	E	$\frac{V}{m}$	Field strength
Swirl impulse density	Magnetic flux rate	$b_{\odot}$	$\frac{V}{m^2}$	Swirl impulse rate
Swirl charge density	Magnetic flux density	$\rho_{\odot}$	$\frac{N}{m^2}$	Force per area
Swirl coupling gain	EM impulse gain	$\mathcal{G}_{\odot}$	$\frac{V \cdot s}{N}$	Coupling constant
Swirl susceptibility	Magnetic susceptibility	χ_H	$\frac{N}{A \cdot m}$	Medium response
Swirl permittivity	Dielectric permittivity	$\mathcal{K}_E = \varepsilon$	$\frac{C}{V \cdot m}$	Areal response
Areal swirl charge	Electric displacement	η	$\frac{C}{m^2}$	Areal swirl charge
Swirl current source	Current density	j	$\frac{A}{m^2}$	Swirl-induced current
Charge-to-current gain	Swirl gain	$\mathcal{G}_{el}$	$\frac{A \cdot s}{C}$	Converts $\partial_t \eta$ to j
Swirl H-field	Magnetic field	H	$\frac{A}{m}$	Swirl H analog
Swirl permeability	Magnetic permeability	μ	$\frac{N}{A^2}$	Magnetic conductivity
Swirl D-field	Electric displacement	D	$\frac{C}{m^2}$	Field-induced charge
Swirl B-field	Magnetic flux density	B	T	Magnetic field strength

Figure 2: Canonical Swirl–Electromagnetic Unit Bridge Table.