


An introduction to Physically-based animation: ODEs and PDEs


ODEs

- ODEs implicitly define an unknown function through its derivative
- Only a single parameter, e.g., time



Solving ODEs Numerically

- Given initial conditions $q(0), \dot{q}(0)$, find function $q(t)$
- Replace time-continuous function $q(t)$ with discrete samples q_i at time t_i





Solving ODEs Numerically

- Numerical time integration approximates

$$q(t+h) = q(t) + \int_t^{t+h} \dot{q}(t) dt$$

- Different 'quadrature rules' lead to different behaviors
 - Explicit Euler is simple but only conditionally stable
 - Implicit Euler is unconditionally stable but expensive

$$\dot{q} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} q$$



Partial Differential Equations (PDEs)

Differential Equations

An Ordinary Differential Equation (ODE) describes an unknown function through its derivatives with respect to a single variable.

Example ODE: $m \frac{d^2 x(t)}{dt^2} = F(x(t))$


A Partial Differential Equation (PDE) describes an unknown function through its partial derivatives with respect to **multiple** variables.

Example PDE: $\frac{\partial u(t, x)}{\partial t^2} = c^2 \frac{\partial u(t, x)}{\partial x^2}$

Why PDEs?

*The book of nature is written in
the language of mathematics.*

Galileo Galilei



- Many physical phenomena and processes can be described by partial differential equations (PDEs)
- PDEs arise naturally from
 - conservation laws (mass, momentum, energy)
 - equilibrium conditions, ...

Fluid Simulation in Graphics

Incompressible Navier Stokes Equations

$$\nabla \cdot \mathbf{u} = 0$$

$$\frac{\partial \mathbf{u}}{\partial t} + (\mathbf{u} \cdot \nabla) \mathbf{u} - \nu \nabla^2 \mathbf{u} = -\nabla w + \mathbf{g}.$$

Fluid Simulation in Graphics



[Losasso, F., Shinar, T. Selle, A. and Fedkiw, R., "*Multiple Interacting Liquids*"](#)

Fluid Simulation in Graphics


<https://jzehnder.me/publications/advectionReflection/>

Elasticity in Graphics in Graphics

Governing Equations of Continuum Mechanics

$$\nabla \cdot \sigma + f = m \cdot a$$

Elasticity in Graphics



[Irving, G., Schroeder, C. and Fedkiw, R., "Volume Conserving Finite Element Simulation of Deformable Models"](#)

Cloth Simulation in Graphics



Zhili Chen, Renguo Feng and Huamin Wang, *"Modeling friction and air effects between cloth and deformable bodies"*


Magnetism in Graphics

Maxwell Equations (static case)

$$\nabla \cdot \mathbf{B} = 0, \quad \nabla \times \mathbf{H} = \mathbf{J}$$

$$\mathbf{H} = \frac{1}{\mu_0} \mathbf{B} - \mathbf{M},$$


Magnetism in Graphics



<https://www.youtube.com/watch?v=NTIbGGcsYek>

A Simple Example: 1D Advection

Weather forecast: simulate temperature evolution.




Given: initial temperature distribution $T_0(x) = T(x, 0)$
and wind speed c .

Find: temperature distribution $T(x, t)$ for any t .

A Simple Example: 1D Advection

- How does the temperature change over a time interval Δt ?



$$T(x, t + \Delta t) = T(x - c\Delta t, t)$$

$$\Delta T = T(x, t + \Delta t) - T(x, t)$$

$$T(x - c\Delta t, t) = T(x, t) - \frac{\partial T}{\partial x} c\Delta t + O(\Delta t^2) = T(x, t + \Delta t)$$

1D advection equation

$$\frac{\Delta T}{\Delta t} \approx -c \frac{\partial T}{\partial x} \quad \Delta t \rightarrow 0 \quad \Rightarrow \quad \frac{\partial T}{\partial t} = -c \frac{\partial T}{\partial x}$$


Analytical Solution

- Any $T(x,t)$ of the form $T(x,t) = f(x - ct)$ solves $\frac{\partial T}{\partial t} = -c \frac{\partial T}{\partial x}$
- The solution also needs to satisfy the initial condition $T(x,0) = T_0(x)$
- The solution is thus $T(x,t) = T_0(x - ct)$

Note: *only simple PDEs can be solved analytically!*

Numerical Solution

- Sample temperature $T(x,t)$ on 1D grids $T^t[i] = T(i \cdot h, t \cdot \Delta t)$ with $i \in (1, \dots, n)$, $t \in (0, 1, 2, \dots)$



- Discretize derivatives with **finite differences** (*space & time*)

$$\frac{\partial T}{\partial t} = -c \frac{\partial T}{\partial x} \quad \Rightarrow \quad \frac{T^{t+1}[i] - T^t[i]}{\Delta t} = -c \frac{T^t[i] - T^t[i-1]}{h}$$

- Solving for $T^{t+1}[i]$ yields update rule

$$T^{t+1}[i] = T^t[i] - \Delta t \cdot c \frac{T^t[i] - T^t[i-1]}{h}$$

- Provide initial values $T^0[i]$
- Set boundary conditions, e.g. *periodic*

$$T^t[0] = T^t[n]$$

Some Notation

- Abbreviation

$$u_{tt} = \frac{\partial^2}{\partial t^2} u(t, \dots), \quad u_{xy} = \frac{\partial^2}{\partial x \partial y} u(x, y, \dots)$$

- Spatial variables

$$\mathbf{x} = (x_1, \dots, x_d)^t$$

- Nabla operator

$$\nabla = \left(\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_d} \right)^t \quad \nabla s = \left(\frac{\partial s}{\partial x_1}, \dots, \frac{\partial s}{\partial x_d} \right)^t$$

- Laplace operator

$$\Delta = \nabla^t \cdot \nabla = \nabla_{\mathbf{x}}^2 = \sum_{k=1}^d \frac{\partial^2}{\partial x_k^2}$$

(in d dimensions)

PDE Classification

- **Order** of PDE = order of highest partial derivative
- A PDE is **linear** if the unknown function u and its partial derivatives only occur linearly

Linear example

$$u_t + c \cdot u_x = 0$$

Advection equation

Nonlinear example


$$u_t + u \cdot u_x = 0$$

Burgers' equation

- Coefficients of linear PDEs can be nonlinear functions

$$y^2 \cdot u_{yy} + x^2 \cdot u_{yy} = 0$$

PDE Classification

- 2nd order linear PDEs are of high practical relevance
  *dedicated classification*

- A 2nd order linear PDE in 2 variables has the form

$$Au_{xx} + 2Bu_{xy} + Cu_{yy} = F(x, y, u, u_x, u_y)$$

- A 2nd order linear PDE in 2 variables is
 - **Hyperbolic** $B^2 - AC > 0$ *(Wave equation)*
 - **Parabolic** $B^2 - AC = 0$ *(Heat equation)*
 - **Elliptic** $B^2 - AC < 0$ *(Laplace equation)*

Model Equations

- Fundamental behavior of many important PDEs is well-captured by three model linear equations:

“Laplacian” (more later!)

LAPLACE EQUATION (“ELLIPTIC”) $\Delta u = 0$

“what’s the smoothest function interpolating the given boundary data”

HEAT EQUATION (“PARABOLIC”) $\dot{u} = \Delta u$


“how does an initial distribution of heat spread out over time?”

WAVE EQUATION (“HYPERBOLIC”) $\ddot{u} = \Delta u$

“if you throw a rock into a pond, how does the wavefront evolve over time?”


[NONLINEAR + HYPERBOLIC + HIGH-ORDER]

Solve numerically?



Elliptic PDEs / Laplace Equation



What's the smoothest function interpolating given boundary data?



- Conceptually: each value is at the average of its “neighbors”
- Very robust to errors: just keep averaging with neighbors!

Parabolic PDEs / Heat Equation

How does an initial distribution of heat spread out over time?



- After a long time, solution is same as Laplace equation!
- Models damping / viscosity in many physical systems

Hyperbolic PDEs / Wave Equation

If you throw a rock into a pond, how does the wavefront evolve over time?



- No steady state solution
- Errors made at the beginning will persist for a long time

How can we solve PDEs?


Numerical Solution of PDEs—Overview

- Like ODEs, many interesting PDEs are difficult or impossible to solve analytically
- Must instead use numerical integration
- Basic strategy:
 - pick a spatial discretization
 - pick a time discretization (forward Euler, backward Euler...)
 - as with ODEs, run a time-stepping algorithm

Real Time PDE-Based Simulation (Fire)



Real Time PDE-Based Simulation (Water)




[Nuttapong Chentanez](#), [Matthias Müller](#), *"Real-time Eulerian water simulation using a restricted tall cell grid"*

Lagrangian vs. Eulerian


- Two basic ways to discretize space: Lagrangian & Eulerian (particle-based & grid-based)
- Suppose you want to keep track of the weather...

LAGRANGIAN



track moving particles and
read what they are measuring

EULERIAN




record temperature at
fixed locations in space

Lagrangian vs. Eulerian—Trade-Offs


■ Lagrangian

- conceptually easy (like polygon soup!)
- resolution/domain not limited by grid
- good particle distribution can be tough
- finding neighbors can be expensive



■ Eulerian

- fast, regular computation
- easy to represent, e.g., smooth surfaces
- simulation “trapped” in grid



Mixing Lagrangian & Eulerian

- Of course, no reason you have to choose just one!
- Many modern methods mix Lagrangian & Eulerian:
 - PIC/FLIP, particle level sets, mesh-based surface tracking, Voronoi-based, arbitrary Lagrangian-Eulerian (ALE), ...
- *Pick the right tool for the job!*

Maya Bifrost



How do we solve *easy* PDEs?

Numerical PDEs—Basic Strategy

- Pick PDE that models phenomenon of interest
 - Which quantity do we want to solve for?
- Pick spatial discretization
 - How do we approximate derivatives in space?
- Pick time discretization
 - How do we approximate derivatives in time?
 - When do we evaluate forces?
 - Forward Euler, backward Euler, symplectic Euler, ...
- Finally, we have an *update rule*
- Repeatedly solve to generate an animation

The Laplace Operator

- All of our model equations used the Laplace operator

Nabla operator

$$\nabla = \left(\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_d} \right)^t \quad \nabla u = \left(\frac{\partial u}{\partial x_1}, \dots, \frac{\partial u}{\partial x_d} \right)^t$$

Laplace operator


$$\Delta = \nabla \cdot \nabla = \sum_{k=1}^d \frac{\partial^2}{\partial x_k^2} \quad \Delta u = \overset{\text{div}}{\nabla} \cdot \overset{\text{grad}}{\nabla} u$$

$$\Delta u = \frac{\partial^2 u}{\partial x_1^2} + \dots + \frac{\partial^2 u}{\partial x_n^2}$$

Discretizing the Laplacian


- How do we approximate the Laplacian?
- Depends on discretization (Eulerian, Lagrangian, grid, mesh, ...)
- Two extremely common ways in graphics:

GRID h




$$\frac{4u_{ij} - u_{i+1,j} - u_{i-1,j} - u_{i,j+1} - u_{i,j-1}}{h^2}$$

(actually, this becomes that)



TRIANGLE MESH



$$\frac{1}{2} \sum_j (\cot \alpha_{ij} + \cot \beta_{ij}) (u_j - u_i)$$

Numerically Solving the Laplace Equation

- Want to solve $\Delta u = 0$
- Plug in one of our discretizations, e.g.,

	c	
d	a	b
	e	

$$\frac{4a - b - c - d - e}{h} = 0$$
$$\iff a = \frac{1}{4}(b + c + d + e)$$

- At solution that solves the Laplace Equation, each value is the average of neighboring values.
- How do we solve this?
- One idea: keep averaging with neighbors! (“Jacobi method”)
- Correct, but *slow* convergence

Boundary Conditions for Discrete Laplace

- What values do we use to compute averages near the boundary?


	c	
$?$	a	b
	e	

$$a = \frac{1}{4}(b + c + ? + e)$$

- Two basic boundary conditions:
 1. *Dirichlet*—boundary data always set to fixed values
 2. *Neumann*—specify derivative (difference) across boundary

Dirichlet Boundary Conditions


- *Dirichlet* means “prescribe values”
- E.g., $\Phi(0)=a$, $\Phi(1) = b$



- Many possible functions “in between”!

Neumann Boundary Conditions


- *Neumann* means “prescribe derivatives”
- E.g., $\Phi'(0)=u$, $\Phi'(1) = v$



- Again, many possible functions

2D Laplace w/ Dirichlet BCs

- 2D Laplace: $\Delta \Phi = 0$
- Q: Can satisfy any Dirichlet BCs? (given data along boundary)



- Yes: Laplace is long-time solution to heat flow
- Data is “heat” at boundary. Then just let it flow...

Solving the Heat Equation

- Back to our three model equations, want to solve *heat diffusion equation*

$$\dot{u} = \Delta u$$

- Just saw how to discretize Laplacian
- Also know how to do time (forward Euler, backward Euler, ...)
- E.g., forward Euler:

$$u^{k+1} = u^k + \Delta u^k$$

- Q: On a grid, what's our overall update now at $u_{i,j}$?

$$u_{i,j}^{k+1} = u^k + \frac{\tau}{h^2} (4u_{i,j}^k - u_{i+1,j}^k - u_{i-1,j}^k - u_{i,j+1}^k - u_{i,j-1}^k)$$

- *Not* hard to implement! Loop over grid, add up some neighbors.

Solving the Wave Equation

- Finally, wave equation:

$$\ddot{u} = \Delta u$$

- Not much different; now have 2nd derivative in time
 - Convert to two 1st order (in time) equations:

$$\dot{u} = v, \quad \dot{v} = \Delta u$$

- Evaluate spatial derivative:


$$\frac{u^{k+1} - 2u^k + u^{k-1}}{\tau^2} = \Delta u^k$$

- Integrate forward in time using, for example, symplectic Euler
- This is just one way to solve this PDE. There are *many* other choices!

Want to Know More?

- There are some good books:
- And papers:

<http://www.physicsbasedanimation.com/>




Biomechanical Simulation and Control of Hands and Tendinous Systems

Prashant Sachdeva, Shinjiro Sueda, Susanne Bradley, Mikhail Fain, Dinesh K. Pai

Search...

Contact

This site is managed by
Christopher Batty from the
University of Waterloo



- Also, what did the folks who *wrote* these books & papers read?

