

FLEXIBLE IMPUTATION OF INCOMPLETE NETWORK DATA ^{*}

Ge SUN[†] Weisheng ZHANG [‡]

April 4, 2026

Abstract

Sampled network data are common in empirical research because collecting full network information is costly, but using sampled networks can lead to biased estimates. We propose a non-parametric imputation method for sampled networks and show that empirical analysis based on imputed networks yields consistent parameter estimates. Our approach imputes missing network links by combining a projection onto covariates with a local two-way fixed-effects regression, which avoids parametric assumptions, does not rely on low-rank restrictions, and flexibly accommodates both observed covariates and unobserved heterogeneity. We establish entrywise convergence rates for the imputed matrix and prove the consistency of GMM estimators based on the imputed network. We further derive the convergence rate of the corresponding estimator in the linear-in-means peer-effects model. Simulations show strong performance of our method both in terms of imputation accuracy and in downstream empirical analysis. We illustrate our method with an application to the microfinance network data of [Banerjee et al. \(2013\)](#).

Keywords: Sampled network, network formation model, dyadic nonparametric regression, GMM

^{*}We thank Aureo de Paula, Chen-Wei Hsiang, Andrei Zeleneev, and the participants of the UCL Econometrics Brownbag Seminar for their valuable comments.

[†]University of Notre Dame: gsun4@nd.edu.

[‡]University College London: weisheng.zhang.21@ucl.ac.uk.

1 Introduction

There is a large and growing literature studying how social networks shape individuals’ behavior and economic outcomes. Examples include how social interactions affect information diffusion (Banerjee et al., 2013; Beaman and Dillon, 2018), test scores (Sacerdote, 2001), and demand for financial assets (Conley and Udry, 2010; Cai, Janvry and Sadoulet, 2015). While network structure is central to these analyses, collecting complete network data is often expensive (Breza et al., 2020), so applied work typically relies on sampled or partially observed networks instead. However, it has been noted that using sampled networks in place of the full network may lead to biased regression coefficients and GMM estimators; see, for example, Chandrasekhar and Lewis (2011). As a result, many empirical conclusions may be undermined by incomplete network data.

To overcome this limitation, we propose a flexible imputation method for sampled networks and show that downstream empirical analysis using the imputed network delivers consistent parameter estimators. Our method is designed for networks generated by dyadic formation processes and observed through egocentric sampling, a widely used design in which researchers randomly sample¹ a subset of individuals and record all of their links in the network, not only those within the sample. The proposed procedure imputes missing links by combining two parts: (i) one explained by observable covariates, recovered via projection onto the observed covariates, and (ii) a residual part estimated using a local two-way fixed-effects regression. The method has several advantages over existing approaches, making it particularly well suited to empirical studies of social networks in economics.

First, the proposed method avoids parametric assumptions. Many imputation approaches, such as Chandrasekhar and Lewis (2011), Breza et al. (2020), and Boucher and Houndetoungan (2025), require specifying the functional form of the network formation model. In practice, this is a demanding requirement, as applied researchers rarely have reliable prior information about the correct functional form. Our approach circumvents this problem by using a fully nonparametric procedure and is therefore robust to misspecification of the network formation process.

Second, our imputation method does not rely on a low-rank assumption. Assuming a low-rank network formation model and exploiting this structure for imputation is a common approach in the statistical network literature (Li et al., 2023; Cai, Cai and Zhang, 2016), but this assumption may be violated when the network formation process is nonlinear, as in the specification in Graham (2017). Building on ideas from Freeman and Weidner (2023) and Feng (2023), our method instead approximates network formation using *local* two-way fixed-effects regressions, making it applicable to a broad class of *nonlinear* dyadic formation processes.

Third, our procedure jointly exploits observed covariates and accommodates unobserved heterogeneity in link formation. Social networks typically exhibit homophily—meaning that individuals with similar characteristics are more likely to form links—and degree heterogeneity—individuals differ systematically in the number of links. An effective imputation method should therefore exploit information from observed covariates and accommodate latent features. The method proposed in this paper achieves both objectives without imposing additional structure on the joint distribution of latent factors and observed covariates.

The general nonparametric network formation setting leads to a technical challenge: because the underlying formation model is unknown and potentially nonlinear, we cannot directly estimate latent factors, as is common in linear factor models, and then use them to recover the formation process and impute missing links. To address this problem, we draw on the insight of Zhang, Levina and Zhu (2017) that similarity in observed connections can reveal similarity in latent factors, and use the sampled

¹Our method also applies to more general sampling schemes in which individuals are randomly sampled with heterogeneous probabilities that depend on both observed and unobserved characteristics.

subnetwork to construct a pseudo-distance that serves as a proxy for latent distance. For each unsampled individual, we then use this pseudo-distance to locate sampled individuals who are “close” to her and use their observed connections to impute her missing links.

We establish a convergence rate for the imputed link probabilities as the network size and the number of sampled individuals go to infinity. The convergence is uniform over all missing links, which implies that the imputation error is controlled at the link level and is particularly useful for applications that require accurate imputation of each link probability. We further provide a bias-variance decomposition of the imputation error, which, to the best of our knowledge, is new in this context. The decomposition shows that the approximation error introduced by using pseudo-distance has an asymmetric effect: it is asymptotically negligible for the bias, but contributes to the variance.

We then establish the consistency of the GMM estimator constructed from the imputed network. This result is particularly relevant for empirical applications, where interest typically centers on the consistency of estimators based on the imputed network rather than on the accuracy of the imputed links themselves. In addition, because our imputation procedure does not require specifying a parametric network formation model and does not rely on low-rank assumptions, the resulting GMM estimator is correspondingly more robust than those obtained using existing approaches.

It is technically challenging to characterize how imputation error affects the asymptotic distribution of the GMM estimator. This is because the moment conditions rely on network statistics, which aggregate link-level imputation errors and thereby induce nonclassical measurement error.

It is technically challenging to characterize how imputation error affects the asymptotic distribution of the GMM estimator. This difficulty arises because (i) imputation error is non-classical and correlated across links, and (ii) the moment conditions rely on network statistics, which aggregate these errors and thereby further complicate the analysis. In this paper, we focus on the widely used linear-in-means peer-effects model and derive the asymptotic orders of the bias and variance of the resulting GMM estimator. We show that the optimal bandwidth for link-level imputation is generally not optimal for downstream estimation. We also provide guidance to help applied researchers assess the valid inference of parameter estimates obtained using our imputation approach.

We provide extensive simulation evidence to evaluate the finite-sample performance of our imputation method. First, we compare its imputation accuracy with alternative approaches and show that our method outperforms them across a wide range of sampling rates and network sparsity levels. We then evaluate the performance of our method in empirical analysis, including regressions based on network statistics and the linear-in-means peer-effects model. These simulations demonstrate that the estimates of regression coefficients and GMM estimators based on our imputed networks exhibit strong finite-sample performance. Finally, we illustrate the practical usefulness of the proposed method through an empirical application to the study of how social networks affect microfinance adoption (Banerjee et al., 2013).

Related literature We contribute to the literature on estimating model parameters with sampled network data by developing an imputation method that allows for a fully nonparametric network formation model. Chandrasekhar and Lewis (2011) show that models estimated using sampled networks are generally biased. They propose an analytical correction for certain special cases and, for more general settings, they estimate a parametric network formation model and conduct GMM using simulated moments. Boucher and Houndetoungan (2025) study the estimation of the linear-in-means peer-effects model when the network data are corrupted, taking egocentrically sampled networks as a special case. Their approach relies on obtaining a consistent estimator of the distribution of the network, which is typically available only under parametric specifications. Herstad (2023) also examine peer-effects estimation with missing links, focusing on settings with large networks and relying on the network formation

model of [Graham \(2017\)](#). A key distinction relative to this line of work is that our method does not rely on parametric specifications of the network formation process and imposes no restrictions on unobserved heterogeneity. Therefore, parameter estimation based on our imputation method is more robust. [Hsieh et al. \(2024\)](#) propose weighted estimators to recover several network-level statistics without assuming any network formation model. In comparison, our approach accommodates more general sampling mechanisms and supports a broader class of downstream regression analyses. In addition, our approach to GMM estimation differs from that of [Chandrasekhar and Lewis \(2011\)](#), who construct moments by integrating over the missing network data. Instead, we directly plug the imputed network into the moment conditions to avoid heavy computational burden.

Another related line of research studies the recovery of missing links under partial network data or even in the absence of network data. [McCormick and Zheng \(2015\)](#) and [Breza et al. \(2020\)](#) demonstrate that aggregated relational data can help recover network structure without observing any links. [de Paula, Rasul and Souza \(2025\)](#) study how network ties can be identified from panel data. [Thirkettle \(2019\)](#) show that network statistics are set identified under partially sampled networks, although their analysis relies on a strategic network formation model, whereas ours is based on a dyadic formation framework. Other sampling designs have also been studied—for example, induced subgraphs in [Chatterjee \(2015\)](#) and censored networks in [Griffith \(2022\)](#).

Our work also relates to the growing literature on estimation with mismeasured network data, as empirical analysis based on imputed networks inevitably inherits the errors introduced in the imputation stage. For peer-effects models, [Lewbel, Qu and Tang \(2024\)](#) show that when the number of misclassified links grows at a rate slower than the network size, the 2SLS estimator of [Bramoullé, Djebbari and Fortin \(2009\)](#) remains consistent and standard inference methods remain valid. [Lewbel, Qu and Tang \(2025\)](#) propose an adjusted 2SLS estimator for peer-effects models under i.i.d. link misclassification. [Hardy et al. \(2024\)](#) also study link misclassification under i.i.d. errors and obtain consistent treatment effects using an expectation-maximization algorithm. [Cai \(2022\)](#) examine linear regression on centrality measures under link-level i.i.d. Bernoulli errors, and propose bias-correction and inference methods for OLS estimators. The setting we consider is technically more challenging than those in the existing literature. Our imputed networks contain errors that are dependent across links and may have bias of the same order as their standard deviation. Our paper contributes to the literature by characterizing the convergence rate of linear-in-means peer-effects estimators in settings that allow for far more general forms of network measurement error.

Methodologically, the studies most directly connected to ours are in the recent literature on nonlinear factor models in panel-data causal inference, particularly the methods of [Feng \(2020\)](#), [Feng \(2023\)](#), and [Deaner, Hsiang and Zelenev \(2025\)](#), which also allow for nonparametric and nonlinear factor structures. Our local two-way fixed-effects regression is analogous to a local linear regression and therefore achieves a second-order approximation error. In contrast, the estimator in [Deaner, Hsiang and Zelenev \(2025\)](#) corresponds to a local constant regression and attains only a first-order approximation error. The local PCA methods of [Feng \(2020\)](#) and [Feng \(2023\)](#) can be viewed as local polynomial regressions that fully exploit smoothness of nonlinear functions. Our approach differs from local PCA in that it does not require specifying the latent factor dimension and is more robust to the misspecification of smoothness assumptions. More broadly, unlike most methods in this strand of the literature, our approach flexibly incorporates observed covariates when imputing missing values.

Finally, our analysis is connected to several strands of the econometrics and statistics literature. The use of pseudo-distance as a measure of latent similarity was first developed by [Zhang, Levina and Zhu \(2017\)](#) and has since been applied in a range of contexts, including nonparametric graphon estimation ([Zhang, Levina and Zhu, 2017; Zelenev, 2020](#)), controlling for unobservables using network data ([Auerbach, 2022](#)), and nonlinear factor models ([Feng, 2020, 2023; Beyhum and Mugnier, 2024](#);

Mugnier, 2025; Deaner, Hsiang and Zeleneev, 2025). Our analysis is also related to the dyadic network formation literature (Graham, 2017; Dzemski, 2019; Chen, Fernández-Val and Weidner, 2021; Ma, Su and Zhang, 2022) and the graphon estimation literature (Gao, Lu and Zhou, 2015; Xu, 2018). Kitamura and Laage (2024) consider covariate-assisted graphon estimation similar to ours, but they focus on stochastic block models, whereas we focus on more general nonlinear link-formation process. Finally, extracting the component of the network explained by covariates relies on the dyadic nonparametric regression developed by Graham, Niu and Powell (2021).

The rest of the paper is organized as follows. Section 2 introduces the framework and estimation procedure. Section 3 provides statistical guarantees for the imputation. Section 4 presents theoretical properties of the GMM estimators based on imputed networks. Section 5 presents Monte Carlo results for both the imputation and the downstream empirical analysis. The empirical application to Banerjee et al. (2013) is given in Section 6. Section 7 concludes.

2 Method

2.1 Network Formation Model

Let $A \in \{0, 1\}^{N \times N}$ denote the adjacency matrix of an unweighted network, where $A_{ij} = 1$ if there is a link between i and j and $A_{ij} = 0$ otherwise. We impose $A_{ij} = A_{ji}$ for all i, j and $A_{ii} = 0$, so that the network is symmetric and has no self-loops. For each individual i , let $X_i \in \mathbb{R}^{d_x}$ denote the observed covariates (e.g., gender, education, and other demographic characteristics), and let $\xi_i \in \mathbb{R}^{d_\xi}$ denote the latent factors. The network A is generated according to the following process. Conditional on observable and unobservable characteristics $\{X_i, \xi_i\}_{i=1}^N$, links $\{A_{ij}\}_{1 \leq i < j \leq N}$ are generated independently with probability $\mathbb{P}(A_{ij} = 1 \mid \{X_i, \xi_i\}_{i=1}^N) = f(X_i, \xi_i, X_j, \xi_j)$, where $f : \mathbb{R}^{d_x + d_\xi} \times \mathbb{R}^{d_x + d_\xi} \rightarrow [0, 1]$ is a symmetric function, referred to as the *graphon*, whose functional form is unknown to econometricians. It is therefore convenient to write

$$A_{ij} = f(X_i, \xi_i, X_j, \xi_j) + \epsilon_{ij}, \quad \forall 1 \leq i < j \leq N \quad (2.1)$$

where ϵ_{ij} are realization errors, which is conditionally independent across pairs $\{(i, j) \mid 1 \leq i < j \leq N\}$ given $\{X_i, \xi_i\}_{i=1}^N$.

The network formation model in (2.1) is a conditionally independent dyadic model. Conditional on the observable and unobservable characteristics $\{X_i, \xi_i\}_{i=1}^N$, the graphon f does not depend on other individuals or on other links in the network. In other words, while A_{ij} and A_{ik} ($k \neq j$) may be correlated unconditionally, they are independent given $\{X_i, \xi_i\}_{i=1}^N$. This framework is flexible and encompasses a wide range of commonly used network formation models. Below, we provide several examples.

Example 1 (Stochastic block model) *A widely used example consistent with our network formation framework is the stochastic block model (SBM). The SBM assumes that each individual belongs to one of G groups, indexed by $g \in \{1, \dots, G\}$, and that the probability of link formation depends solely on group membership. In particular, the probability of a link between two individuals is determined entirely by the groups to which they belong, implying that individuals within the same group share identical connection probabilities with other individuals. The stochastic block model can be viewed as a special case of our network formation process in which the support of (X_i, ξ_i) is finite and consists of only G distinct values. Under this restriction, the graphon f reduces to a piecewise constant function defined over the G^2 possible group pairs.*

Example 2 (Network formation with transferable utility) Suppose that individuals i and j form a link if the total surplus from doing so is positive:

$$A_{ij} = \mathbf{1}(\omega(X_i, X_j)' \beta + g(\xi_i, \xi_j) - U_{ij} \geq 0) \quad (2.2)$$

The total surplus consists of three components:

- **(Homophily in observables)** $\omega(X_i, X_j)$ measures the distance between observable characteristics of i and j . For instance, one may define $\omega(X_i, X_j) = (|X_{i1} - X_{j1}|, \dots, |X_{id_X} - X_{jd_X}|)'$ for absolute differences, or $\omega(X_i, X_j) = ((X_{i1} - X_{j1})^2, \dots, (X_{id_X} - X_{jd_X})^2)'$ for squared differences. A negative coefficient $\beta < 0$ captures homophily, meaning that individuals with similar observables are more likely to form a link.
- **(Individual heterogeneity)** The function $g(\xi_i, \xi_j)$ captures how unobserved individual heterogeneity affects the probability of link formation. For example, consider the specification $g(\xi_i, \xi_j) = \xi_{i1} + \xi_{j1} - (\xi_{i2} - \xi_{j2})^2$. Here, ξ_{i1} represents degree heterogeneity, under which individuals with larger values of ξ_{i1} tend to form more connections. The quadratic term $-(\xi_{i2} - \xi_{j2})^2$ captures homophily in unobservables, with link probabilities decreasing in the distance between individuals' latent characteristics.
- **(Idiosyncratic shocks)** $\{U_{ij} \mid 1 \leq i < j \leq N\}$ captures the idiosyncratic preference shock that is independent of (X, ξ) . Let $F(\cdot)$ be the cumulative distribution of U_{ij} , then $\mathbb{P}(A_{ij} = 1 \mid \{X_i, \xi_i\}_{i=1}^N) = F(\omega(X_i, X_j)' \beta + g(\xi_i, \xi_j))$. Specifically, when U_{ij} follows type-I extreme value distribution, we have


$$f(X_i, \xi_i, X_j, \xi_j) = \frac{\exp(\omega(X_i, X_j)' \beta + g(\xi_i, \xi_j))}{1 + \exp(\omega(X_i, X_j)' \beta + g(\xi_i, \xi_j))}$$

When $g(\xi_i, \xi_j)$ includes only the additive component $\xi_i + \xi_j$, the model reduces to the model of [Graham \(2017\)](#), which has been widely used in empirical applications.


Beyond these examples, our network formation framework in (2.1) can flexibly accommodate richer forms of heterogeneity. For example, it allows for heterogeneous slope coefficients when the homophily effect on observables, captured by β in (2.2), depends on the latent factors (ξ_i, ξ_j) . Similarly, the unobserved heterogeneity term $g(\xi_i, \xi_j)$ can capture more complex interaction structures, such as interactive fixed effects of the form $\xi_i' \xi_j$. However, our dyadic model excludes strategic network formation, where the formation of one link may depend on the presence of other links (see [de Paula \(2020\)](#)).

2.2 Data

Economists observe individual covariates X_i (e.g., gender, age) for *all* i in the network but do not observe the full network A ; instead, only a sampled network is available. In this paper, we focus on egocentric sampling, which is one of the most commonly used sampling designs in empirical work. Specifically, we randomly sample $n < N$ nodes in the network and ask all their friends not only in the sample but in the whole network. The sampling probabilities can depend on both covariates X_i and unobserved characteristics ξ_i . Let $\mathcal{S} \subset \{1, 2, \dots, N\}$ denote the set of sampled individuals with $|\mathcal{S}| = n$. Then a link A_{ij} is observed if $i \in \mathcal{S}$ or $j \in \mathcal{S}$, whereas links between two non-sampled nodes (i.e., $i \notin \mathcal{S}$ and $j \notin \mathcal{S}$) are unobserved. A simple example follows to illustrate the setup.



(a) Full network



(b) Sampled network by sampling $\{1, 2\}$, red nodes are not in the sample, dashed edges are not observed

$$\begin{pmatrix} 0 & 0 & 1 & 1 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 1 & 1 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 & 1 & 0 \end{pmatrix} \Rightarrow \begin{pmatrix} 0 & 0 & 1 & 1 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ \hline 1 & 1 & \times & \times & \times & \times \\ 1 & 0 & \times & \times & \times & \times \\ 1 & 0 & \times & \times & \times & \times \\ 0 & 0 & \times & \times & \times & \times \end{pmatrix}$$

(c) From full adjacency matrix to partially observed adjacency matrix, $\times$ denotes missing entry

Figure 1: An example of how egocentric sampling works

Since it causes no harm to relabel the agents, we can do so such that $\mathcal{S} = \{1, 2, \dots, n\}$, in which case the observed adjacency matrix A^{obs} admits the following block structure:

$$A^{\text{obs}} = \begin{pmatrix} A_{\mathcal{S}\mathcal{S}} & A_{\mathcal{S}\mathcal{S}^c} \\ A_{\mathcal{S}^c\mathcal{S}} & \textcolor{red}{A_{\mathcal{S}^c\mathcal{S}^c}} \end{pmatrix} \quad (2.3)$$

Here, $A_{\mathcal{S}\mathcal{S}}$ consists of links between two sampled individuals, $A_{\mathcal{S}^c\mathcal{S}} = A'_{\mathcal{S}\mathcal{S}^c}$ contains links between sampled and unsampled individuals, and the bottom right block $A_{\mathcal{S}^c\mathcal{S}^c}$ is unobserved because both individuals in each dyad are unsampled; we highlight this block in red.

2.3 Imputation

To illustrate the main idea of our imputation approach, let us start with a simple case in which both X_i and ξ_i are observable for every individual in the network. In this case, imputing missing links would be straightforward: we could estimate $f(\cdot, \cdot)$ nonparametrically from the observed adjacency matrix to obtain estimates $\hat{f}(\cdot, \cdot)$, and then use these estimates to impute the missing links in the submatrix $A_{\mathcal{S}^c\mathcal{S}^c}$.

The main difficulty, however, is that the latent factors ξ are unobserved for all individuals. To make this challenge more transparent, let us first abstract from the observed covariates X and focus on the case where link formation depends solely on the latent factors. In this setting, the imputation problem reduces to recovering information about the latent factors ξ from the observed subnetwork and using them to predict the missing entries of the adjacency matrix.

Our approach is motivated by the following idea. When network formation is dyadic, as in (2.1), individuals with similar ξ are expected to have similar rows in the (denoised) adjacency matrix. This observation provides a useful heuristic: similarity in observed connections can indicate similarity in latent

positions. Building on this idea, for any unsampled individual i . The similarity is inferred by comparing their connection patterns in the observed subnetwork. We then use the observed connections of these sampled individuals, weighted by their similarity to i , to impute the missing links for i .

Because the information about ξ is estimated rather than directly observed, it is inappropriate to ignore the observed covariates or absorb them into the latent factors, as doing so would discard the noiseless information contained in X . Therefore, when X is available, our method first extracts the component of link formation explained by the observed covariates and then applies the latent-factor-based procedure described above to the residuals. These two components are subsequently combined to obtain the final imputed network.

2.3.1 Extract information explained by X

For notational simplicity, we write $f_{ij} = f(X_i, \xi_i, X_j, \xi_j)$. As discussed previously, when individual covariates X are available, it is appropriate to first extract the component that can be explained by X . Consider the decomposition:

$$f_{ij} = \underbrace{\mathbb{E}(f_{ij} \mid X_i, X_j)}_{\Pi_{ij}} + \underbrace{f_{ij} - \mathbb{E}(f_{ij} \mid X_i, X_j)}_{\Pi_{ij}^\perp}$$

where Π_{ij} represents the part of the graphon that can be explained by the observed covariates, and $\Pi_{ij}^\perp$ captures the residual component of the graphon that cannot be explained solely by the observed covariates. Note that the observed link A_{ij} can then be written as $A_{ij} = \Pi_{ij} + (\Pi_{ij}^\perp + \epsilon_{ij})$, and the composite error term satisfies $\mathbb{E}(\Pi_{ij}^\perp + \epsilon_{ij} \mid X_i, X_j) = 0$, which implies that dyadic nonparametric regression of A_{ij} on (X_i, X_j) consistently estimates Π_{ij} . [Graham, Niu and Powell \(2021\)](#) derive the pointwise and uniform minimax risks for estimating such conditional expectations and show that the Nadaraya-Watson (NW) estimator attains the optimal convergence rate when using higher-order kernels². In addition, dyadic nonparametric regression can be implemented using observed subset of adjacency matrix, and since the covariates X are observed for every individual in the network, the nonparametric estimator $\hat{\Pi}_{ij}$ can be computed even when both i and j are unsampled. The implementation details are discussed in the Appendix.

In the next two steps, we focus on estimating this residual component $\Pi^\perp$ using pseudo-distance and two-way fixed-effect imputation procedure described below.

2.3.2 Calculate similarity

We now describe how to infer similarity between latent factors using observed network links.

For notational simplicity, let $\zeta_i := (X_i, \xi_i)$ for each i , and let $f_{ij} = f(\zeta_i, \zeta_j)$ for all $1 \leq i, j \leq N$. The adjacency matrix A can be expressed as $A = P + E$. Here, P collects the conditional probabilities of link formation, i.e., $P_{ij} = f_{ij}$ when $i \neq j$, and $P_{ii} = 0$ (because self-loops are ruled out), and E collects error terms ϵ_{ij} ³. Our approach relies on the following idea. For any unsampled individual i and sampled individual i' , if their denoised rows in the observed network (or equivalently, the corresponding rows in P) appear sufficiently similar, then under suitable regularity conditions, ζ_i and $\zeta_{i'}$ are also similar in the latent space.

²In practice, however, we implement local polynomial regression rather than higher-order kernels, as it is the standard approach in local regression.

³Likewise, we impose $\epsilon_{ii} = 0$ for each i .

Following [Zhang, Levina and Zhu \(2017\)](#), we define the pseudo-distance between i and i' as

$$\hat{d}_{ii'} = \max_{\substack{k=1,\dots,N, \\ k \neq i, i'}} \left| \frac{1}{n} \sum_{\ell=1}^n A_{k\ell} (A_{i\ell} - A_{i'\ell}) \right|, \quad i, i' = 1, 2, \dots, N$$

The definition above relies only on the friendships of sampled individuals and can therefore be computed from the observed subnetwork. We will show that the pseudo-distance $\hat{d}$ effectively de-noises the adjacency matrix A , so that the distances computed from A and from the noise-free matrix F are asymptotically equivalent. In addition, under regularity conditions discussed later, a small value of $\hat{d}_{ii'}$ implies that the corresponding latent characteristics ζ_i and $\zeta_{i'}$ are sufficiently close in the latent space.

It is important to note that the discussion here concerns the composite characteristic ζ_i , rather than the observable covariate X_i alone. This is because (i) X_i and ξ_i enter the network formation process in a non-separable way, making it impossible to isolate their effects without additional assumptions; and (ii) the information in X_i that can be directly explained has already been extracted in the dyadic nonparametric regression, so there is no need to distinguish between X_i and ξ_i in this step.

2.3.3 Local two-way fixed-effects imputation

For the projection residuals, we employ the pseudo-distance $\hat{d}$ obtained in the previous step to construct a local estimator for $\Pi_{ij}^\perp$, referred to as the two-way fixed-effect imputation. Specifically, for unsampled individuals i and j , we estimate the local two-way fixed-effects regression to obtain $\hat{a}_i$ and $\hat{b}_j$:

$$(\hat{\mathbf{a}}, \hat{\mathbf{b}}) \in \underset{\mathbf{a}, \mathbf{b} \in \mathbb{R}^{n+1}}{\operatorname{argmin}} \sum_{\substack{i' \in \mathcal{S} \cup \{i\}, j' \in \mathcal{S} \cup \{j\} \\ (i', j') \neq (i, j)}} \left(A_{i'j'} - \hat{\Pi}_{i'j'} - a_{i'} - b_{j'} \right)^2 K_h(\hat{d}_{ii'}) K_h(\hat{d}_{jj'}) \quad (2.4)$$

where $K_h(\cdot) := h^{-1}K(\cdot/h)$ is a kernel function with bandwidth h , and $\hat{d}$ denotes the pseudo-distance. Then the imputed $\Pi_{ij}^\perp$ is obtained via $\hat{\Pi}_{ij}^\perp = \hat{a}_i + \hat{b}_j$, and the imputed network link is given by:

$$\hat{A}_{ij} = \hat{\Pi}_{ij} + \hat{\Pi}_{ij}^\perp = \hat{\Pi}_{ij} + \hat{a}_i + \hat{b}_j$$

The imputation is “local” in the sense that, by using a kernel estimator, we rely only on information from individuals i' and j' whose pseudo-distances $\hat{d}_{ii'}$ and $\hat{d}_{jj'}$ (or, equivalently, the distances between their latent positions) are sufficiently small when estimating fixed effects. Note that, although our notations omit the indices (i, j) for simplicity, the estimator $(\hat{\mathbf{a}}, \hat{\mathbf{b}})$ in (2.4) is pair-specific and therefore must be re-estimated for each (i, j) .

The procedure is referred to as a “two-way fixed-effects imputation” because (2.4) takes the form of the two-way fixed-effects regression rather than a local constant regression. This is analogous to replacing a local constant estimator with a local polynomial estimator in nonparametric regression: incorporating the two-way fixed-effects provides a higher-order local approximation and thus improves estimation accuracy. We will formally justify this result later using a Taylor expansion.

It is important to note that $\hat{A}_{ij}$ can only approximate the conditional link-formation probability f_{ij} rather than the missing link A_{ij} . This is because the missing link contains an unpredictable idiosyncratic error ϵ_{ij} , which cannot be predicted from the available data.

We impute all missing links based on the previous steps and merge these imputed entries with the observed ones to obtain a complete imputed matrix $\hat{A}$. To facilitate understanding of the main idea of our algorithm, we summarize below a simplified version of our method that excludes cross-fitting and cross-validation.

Algorithm 1 Imputation (Basic Idea)

Input: Kernel $K(\cdot)$, bandwidths h , incomplete network A^{obs} , and covariate matrix X .

Output: Imputed network $\hat{A}$.

Step 1: Extract information explained by X

Implement dyadic nonparametric regression to obtain $\hat{\Pi}(\cdot, \cdot)$.

Step 2: Calculate similarity

For each $i \in \mathcal{S}^c$ and $i' \in \mathcal{S}$, calculate the pseudo-distance using observed adjacency matrix:

$$\hat{d}_{ii'} = \max_{\substack{k=1, \dots, N, \\ k \neq i, i'}} \left| \frac{1}{n} \sum_{\ell=1}^n A_{k\ell} (A_{i\ell} - A_{i'\ell}) \right| \quad (2.5)$$

Step 3: Local imputation

For each missing entry (i, j) , i.e., $i \in \mathcal{S}^c$ and $j \in \mathcal{S}^c$, estimate the local two-way fixed-effects regression:

$$(\hat{\mathbf{a}}, \hat{\mathbf{b}}) \in \underset{\mathbf{a}, \mathbf{b} \in \mathbb{R}^{n+1}}{\text{argmin}} \sum_{\substack{i' \in \mathcal{S} \cup \{i\}, j' \in \mathcal{S} \cup \{j\}, \\ (i', j') \neq (i, j)}} \left(A_{i'j'} - \hat{\Pi}_{i'j'} - a_{i'} - b_{j'} \right)^2 K_h(\hat{d}_{ii'}) K_h(\hat{d}_{jj'}) \quad (2.6)$$

to obtain $\tilde{A}_{ij} = \hat{a}_i + \hat{b}_j + \hat{\Pi}_{ij}$. By combining the imputed entries with the observed entries of the adjacency matrix, we obtain the (preliminary) imputed adjacency matrix $\tilde{A}$. Truncate $\tilde{A}_{ij}$ to get

$$\hat{A}_{ij} = \begin{cases} \tilde{A}_{ij}, & \text{if } \tilde{A}_{ij} \in [0, 1] \\ 1 & \text{if } \tilde{A}_{ij} > 1 \\ 0 & \text{if } \tilde{A}_{ij} < 0 \end{cases}$$

and zero the diagonal of $\hat{A}$ such that $\hat{A}_{ii} = 0$ for all i .

2.3.4 Cross-validation and sample-splitting

Although the proposed algorithm is straightforward to implement, deriving the asymptotic properties of $\hat{A}_{ij}$ can be technically challenging because of the potential dependence between the estimated pseudo-distance $\hat{d}$ and the error terms ϵ in the two-way fixed-effects regression.

A standard approach to eliminate this dependence is to employ cross-fitting, in which the data are partitioned into folds so that the first-step and second-step estimations are conducted on separate subsets of the data. Specifically, we split the sample $\mathcal{S}$ into two parts, $\mathcal{S}_1$ and $\mathcal{S}_2$, of equal size. We then use only the friendships of $\mathcal{S}_1$ to calculate the similarity $\hat{d}(\cdot, \cdot)$:

$$\hat{d}_{ii'} = \max_{\substack{k=1, \dots, N, \\ k \neq i, i'}} \left| \frac{1}{|\mathcal{S}_1|} \sum_{\ell \in \mathcal{S}_1} A_{k\ell} (A_{i\ell} - A_{i'\ell}) \right|, \quad i \in \mathcal{S}^c, i' \in \mathcal{S}_2$$

In the imputation stage, we use no link information from $\mathcal{S}_1$, and all calculations rely on the friendships of $\mathcal{S}_2$:

$$(\hat{\mathbf{a}}, \hat{\mathbf{b}}) \in \underset{\mathbf{a}, \mathbf{b} \in \mathbb{R}^{|\mathcal{S}_2|+1}}{\text{argmin}} \sum_{\substack{i' \in \mathcal{S}_2 \cup \{i\}, j' \in \mathcal{S}_2 \cup \{j\}, \\ (i', j') \neq (i, j)}} \left(A_{i'j'} - \hat{\Pi}_{i'j'} - a_{i'} - b_{j'} \right)^2 K_h(\hat{d}_{ii'}) K_h(\hat{d}_{jj'})$$
$$\hat{A}_{ij} = \hat{a}_i + \hat{b}_j + \hat{\Pi}_{ij}$$

In addition, a practical concern is the choice of the tuning parameter h . We recommend selecting h

using a leave-one-out cross-validation procedure, and the details of cross-validation, together with the sample-splitting, are provided in the Appendix.

3 Theory for Imputation

Our method accommodates both low-rank graphon and high-rank graphon, where the support of ζ is a continuum and f is nonlinear. In what follows, we focus on the second case, while noting that the same arguments apply, with minor modifications, to the low-rank graphon. We now introduce the regularity conditions.

Assumption 1 (Regularity Conditions) *Suppose that*

- (i) **(Asymptotics)** *Large- N , large- n asymptotics with $N, n \rightarrow \infty$ and $\frac{\log N}{n} \rightarrow 0$.*
- (ii) **(Independence)** *$\{(X_i, \xi_i)\}_{i=1}^N$ are i.i.d., and conditional on $\{(X_i, \xi_i)\}_{i=1}^N$, $\{A_{ij}\}_{1 \leq i < j \leq N}$ are generated independently according to (2.1).*
- (iii) **(Random sampling)** *Let $D_i \in \{0, 1\}$ indicate whether i is sampled. Assume $D_i \perp (A, X, \xi)$ for all i , and $\{D_i\}_{i=1}^N$ are independent across i .*
- (iv) **(Compact support)** *Let $\zeta_i = (X'_i, \xi'_i)'$ and $d_\zeta = d_X + d_\xi$, assume that the support of ζ_i is compact on $\mathbb{R}^{d_\zeta}$. In addition, there exist constants $0 < \underline{c} < \bar{c} < \infty$ such that for any $\zeta \in \text{supp}(\zeta)$ and any $h > 0$, $\underline{c}h^{d_\zeta} \leq \mathbb{P}(\|\zeta_i - \zeta\| \leq h) \leq \bar{c}h^{d_\zeta}$.*
- (v) **(Smoothness)** *For every fixed $\zeta_2 \in \text{supp}(\zeta)$, the mapping $\zeta \mapsto f(\zeta, \zeta_2)$ is twice continuously differentiable, and its second derivatives are uniformly bounded for all $\zeta, \zeta_2 \in \text{supp}(\zeta)$.*

Assumption 1(i) indicates that our method is designed for large-network settings in which both the network size N and the number of sampled individuals n grow large. Our analysis does not rely on n and N growing at the same rate and continues to hold when n grows more slowly, e.g., $n = O(N^\alpha)$ for any $\alpha \in (0, 1)$.

Assumption 1(ii) imposes that $\{(X_i, \xi_i)\}_{i=1}^N$ are i.i.d., and the data-generating process of A follows a conditionally independent dyadic model (2.1). This model also follows from the Aldous-Hoover Theorem (Aldous, 1981; Hoover, 1979), which shows that any infinitely exchangeable random graph admits such a representation. As discussed earlier, although (2.1) is flexible enough to encompass a wide range of network formation models commonly used in economics and statistics, it rules out strategic network formation, in which individuals linking decisions depend on those of others; see de Paula (2020) for a review.

Assumption 1(iii) is standard in egocentric sampling designs, which assume that individuals are randomly drawn from the population. However, our method also applies to more general sampling schemes in which individuals are randomly sampled with heterogeneous probabilities that depend on both observed and unobserved characteristics.

The first part of Assumption 1(iv) imposes the boundedness of ζ , and the second part introduces a standard bounded-density condition. Intuitively, it ensures that the distribution of ζ_i is neither locally too sparse nor too concentrated by requiring that, for any point ζ and a small radius $h > 0$, the probability that ζ_i falls within a ball of radius h centered at ζ is of the same order as the volume of that ball. This requirement is automatically satisfied when the support is convex and the probability density function of ζ is uniformly bounded above and bounded away from zero. This assumption is common for establishing the asymptotic properties of nonparametric kernel estimators.

Assumption 1(v) requires the graphon f to be twice continuously differentiable with uniformly bounded second derivatives. This smoothness condition is needed because our two-way fixed-effects imputation method relies on a second-order Taylor expansion of $\Pi_{ij}^\perp$ within a local neighborhood to eliminate the first-order approximation bias, thereby improving accuracy. Methods that rely on higher-order Taylor expansions, such as the local PCA approach proposed by Feng (2023), correspondingly require the existence of higher-order derivatives. We discuss this point in detail later.

It should be noted that the requirement of second-order differentiability of f is not a mild condition. For instance, consider the network formation model with degree heterogeneity in Example 2. If the distance function $\omega(\cdot, \cdot)$ is defined by the absolute distance, i.e., $\omega(X_i, X_j) = |X_i - X_j|$, which is common in applied research, then the corresponding graphon f is at most Lipschitz continuous rather than twice differentiable. Nevertheless, our method remains valid and can consistently impute the conditional probability of link formation, although the approximation error no longer attains the second-order rate.

3.1 Asymptotic analysis of pseudo-distance

We now develop the asymptotic properties of the pseudo-distance $\hat{d}_{ii'}$ and show that a sufficiently small $\hat{d}_{ii'}$ implies that ζ_i and $\zeta_{i'}$ are close in the latent space. In the following lemma, we establish the convergence of $\hat{d}_{ii'}$ to its population counterpart $d_{ii'}$, defined as

$$d_{ii'} = \sup_{\zeta_k \in \text{supp}(\zeta)} \left| \int_{\zeta_\ell} g(\zeta_k, \zeta_\ell) (g(\zeta_i, \zeta_\ell) - g(\zeta_{i'}, \zeta_\ell)) d\mathbb{P}_\xi(\zeta_\ell) \right|$$

Lemma 1 *Under Assumption 1, there exist constants $\gamma_1, \gamma_2 > 0$ such that*

$$\mathbb{P} \left(\max_{i \in S^c, i' \in S_2} |\hat{d}_{ii'} - d_{ii'}| \geq \gamma_1 \frac{\log N}{N^{1/d_\zeta}} + \gamma_2 \frac{\log N}{\sqrt{n}} \right) \leq 1 - n^{-1/2}$$

Lemma 1 shows that the approximation error of $\hat{d}_{ii'}$ relative to its population counterpart $d_{ii'}$ can be decomposed into two components. The first term, $\gamma_1 \frac{\log N}{N^{1/d_\zeta}}$, captures the matching discrepancy in the latent characteristics ζ_i . Its rate deteriorates with the dimension of the latent space d_ζ , which is consistent with the well-known “curse of dimensionality” in the matching literature. The second term, $\gamma_2 \frac{\log N}{\sqrt{n}}$, reflects the effect of the idiosyncratic error ϵ_{ij} . It shows that although the true graphon is unobservable, the pseudo-distance can asymptotically denoise the observed network. The matching discrepancy error dominates the effect of idiosyncratic errors when $N \sim n$ and $d_\zeta \geq 3$, which is the common case in practice. Finally, for notational simplicity, let

$$\delta_{N,n} := \gamma_1 \frac{\log N}{N^{1/d_\zeta}} + \gamma_2 \frac{\log N}{\sqrt{n}}$$

and henceforth focus on the event under which $|\hat{d}_{ii'} - d_{ii'}|$ is uniformly bounded by $\delta_{N,n}$, as implied by Lemma 1.

The following assumption is critical and requires that the population pseudo-distance $d_{ii'}$ be sufficiently informative to reveal the latent distance $\|\zeta_i - \zeta_{i'}\|$.

Assumption 2 (Informativeness) *For all $\zeta_i, \zeta_{i'} \in \text{supp}(\zeta)$, there exists a constant $\lambda > 0$ such that $d_{ii'} \geq \lambda \|\zeta_i - \zeta_{i'}\|$.*

This assumption ensures that when the population pseudo-distance $d_{ii'}$ is sufficiently small, the latent positions ζ_i and $\zeta_{i'}$ are close to each other. Hence, $d_{ii'}$ can be used to identify individuals with similar latent characteristics.

Example 1 (Continued) When each individual belongs to one of G groups, $g \in \{1, \dots, G\}$, Assumption 2 is guaranteed as long as there is heterogeneity across different groups, that is, no two groups share identical linking patterns. A formal illustration and the proof are provided in Lemma D.1 in the Appendix. When the graphon f takes a more general low-rank form, the informativeness condition still holds under similarly mild assumptions. We refer the reader to the discussions in Feng (2023) and Deaner, Hsiang and Zelenev (2025) for further details.

Example 2 (Continued) In Example 2, the latent type ζ may lie on a continuous support, and the graphon f is not necessarily low rank. Therefore, verifying Assumption 2 in this setting is generally challenging. Nevertheless, we show that the informativeness assumption holds in the following model which serves as a benchmark in empirical applications. Specifically, Assumption 2 holds when (i) U_{ij} in (2.2) follows a Type-I extreme value distribution, and (ii) the model allows for degree heterogeneity and homophily effects on both observable and unobservable characteristics, where homophily can be captured by either absolute or quadratic distance. This finding implies that the informativeness assumption is empirically plausible and unlikely to impose restrictive constraints in applied network analysis. A formal illustration and the proof are provided in Lemma D.2 in the Appendix.

Remark An alternative to the pseudo-distance is the Euclidean distance. More precisely, for any pair of nodes i and i' , define the squared Euclidean distance as $\hat{r}_{ii'}^2 = \frac{1}{|\mathcal{S}_1|} \sum_{\ell \in \mathcal{S}_1} (A_{i\ell} - A_{i'\ell})^2$. The main drawback of using the squared Euclidean distance is that it does not automatically denoise the observed network, because $\hat{r}_{ii'}^2$ includes the inner products of each error vector with itself, $\frac{1}{|\mathcal{S}_1|} \sum_{\ell \in \mathcal{S}_1} \epsilon_{i\ell}^2 + \frac{1}{|\mathcal{S}_1|} \sum_{\ell \in \mathcal{S}_1} \epsilon_{i'\ell}^2$, which do not vanish even as $n \rightarrow \infty$. In contrast, the pseudo-distance involves cross inner products of independent error vectors, so that the influence of noise decreases at the $1/\sqrt{n}$ rate, thereby asymptotically denoising the adjacency matrix.

Another alternative to the pseudo-distance is the difference in average degrees, defined as $\hat{\rho}_{ii'} = \frac{1}{|\mathcal{S}_1|} \sum_{\ell \in \mathcal{S}_1} A_{i\ell} - \frac{1}{|\mathcal{S}_1|} \sum_{\ell \in \mathcal{S}_1} A_{i'\ell}$. Although the average-degree distance can also denoise the adjacency matrix, it requires a stronger informativeness assumption than Assumption 2. In particular, using average-degree distance requires the population average degree,

$$\zeta \mapsto \int f(\zeta, \zeta') \mathbb{P}(\zeta')$$

be a strictly monotonic function of ζ . This monotonicity requirement corresponds to the injectivity condition in Bonhomme, Lamadon and Manresa (2022). However, the condition does not hold in either Example 1 or Example 2 and is stronger than the Assumption 2 in our paper.

3.2 Asymptotic analysis of imputation

Since the imputation procedure involves two parts, a nonparametric dyadic regression and a two-way fixed-effects regression, the imputation error can accordingly be decomposed into two parts. Our asymptotic analysis focuses on the second part, as the first-stage nonparametric estimator is generally much more accurate. The asymptotic properties of the first-stage estimator have been well established in the literature, see Graham, Niu and Powell (2021).

Assumption 3 The estimation error of $\hat{\Pi}_{ij}$ admits the following uniform bound:

$$\sup_{x_i, x_j \in \mathcal{X}} \mathbb{E} \left(\hat{\Pi}(x_i, x_j) - \Pi(x_i, x_j) \right) = O \left(n^{\frac{-2}{4+d_X}} \log(N) \right), \quad \sup_{x_i, x_j \in \mathcal{X}} \text{Var}^{1/2} \left(\hat{\Pi}(x_i, x_j) \right) = O \left(n^{\frac{-2}{4+d_X}} \log(N) \right)$$

Assumption 3 states that the first-stage nonparametric dyadic regression estimator admits uniform bounds on its bias and standard deviation. Up to logarithmic factors, both bounds attain the rate

$n^{-2/(4+d_X)}$, which is the optimal nonparametric mean-squared-error rate for a twice continuously differentiable target function with d_X -dimensional covariates. Under Assumption 1, [Graham, Niu and Powell \(2021\)](#) show that the Nadaraya-Watson estimator with a higher-order kernel satisfies Assumption 3.

Remark First, although the dyadic conditional expectation $\Pi(\cdot, \cdot)$ takes $2d_X$ arguments, the nonparametric convergence rate in Assumption 3 behaves as if Π depended on only d_X arguments, making it consistent with the conclusion in [Graham, Niu and Powell \(2021\)](#). Second, as noted by [Graham, Niu and Powell \(2021\)](#), when the entire network is observed, the regression uses $O(N^2)$ dyads, but the effective sample size is just $O(N)$ because the estimator is a U-statistic. In our settings, however, the effective sample size is of order n (not $\sqrt{Nn}$) as stated in Assumption 3. This argument can be verified via Hoeffding decomposition of dyadic nonparametric estimators.

Assumption 4 Suppose that the kernel $K : \mathbb{R} \mapsto \mathbb{R}_+$ is bounded and supported on $[-1, 1]$. In addition, $K(0) > 0$ and $K(\cdot)$ is Lipschitz continuous with constant $\bar{K} > 0$.

This assumption imposes regularity conditions on kernel K used in the second step. The requirement is standard in the literature, and many commonly used kernels, including the Epanechnikov kernel, satisfy Assumption 4.

We now briefly explain why $\hat{d}_{ii'}$ can be viewed as a noisy measure of $\|\zeta_i - \zeta_{i'}\|$ and can thus serve as its proxy in the kernel function. First, let h denote the bandwidth of the kernel function. When imputing $\Pi_{ij}^\perp$, Assumption 4 allows us to restrict attention to links (i', j') for which both $\hat{d}_{ii'} \leq h$ and $\hat{d}_{jj'} \leq h$. Second, by Lemma 1, we have $|\hat{d}_{ii'} - d_{ii'}| \leq \delta_{N,n}$ with high probability. When $\delta_{N,n}/h \rightarrow 0$, the approximation error in $\hat{d}_{ii'}$ is asymptotically negligible relative to the bandwidth h , so $d_{ii'}$ can be replaced by $\hat{d}_{ii'}$ without loss of first-order accuracy. Furthermore, under Assumption 2, $d_{ii'}$ is informative about the latent distance in the sense that $\|\zeta_i - \zeta_{i'}\| \leq \lambda^{-1}d_{ii'}$. Combining these results yields, with probability approaching one, $\|\zeta_i - \zeta_{i'}\| \leq \lambda^{-1}h$, which implies that using pseudo-distances in the kernel function selects pairs whose latent distances lie within a $\lambda^{-1}h$ -neighborhood when imputing missing links.

The following theorem provides the theoretical guarantee for our imputation method.

Theorem 2 (Imputation errors) Under Assumption 1, 2, 3, 4, if (i) $h \rightarrow 0$, (ii) $\delta_{N,n}/h \rightarrow 0$, and (iii) $nh^{d_\zeta}/\log N \rightarrow \infty$, then there exists constant $\gamma_3 > 0$ such that:

$$\mathbb{P}\left(\max_{i,j \in \mathcal{S}^c} |\hat{A}_{ij} - P_{ij}| \leq \gamma_3 \left(h^2 + \frac{\log N}{\sqrt{nh^{d_\zeta}}}\right)\right) \geq 1 - n^{-1/2} \quad (3.1)$$

Moreover, the imputation error admits the following bias-variance decomposition. Specifically, with probability at least $1 - n^{-1/2}$,

$$\max_{i,j \in \mathcal{S}^c} \mathbb{E}\left(\hat{A}_{ij} - P_{ij} \mid \{\zeta_i\}_{1 \leq i \leq N}\right) = O\left(h^2 + \frac{1}{nh^{d_\zeta}}\right), \quad \max_{i,j \in \mathcal{S}^c} \text{Var}(\hat{A}_{ij} \mid \{\zeta_i\}_{1 \leq i \leq N}) = O\left(\frac{\log N}{nh^{d_\zeta}} + \delta_{N,n}^2 h^2\right) \quad (3.2)$$

Theorem 2 establishes link-level error bounds, rather than a Frobenius-norm bound for a submatrix (e.g., [Gao, Lu and Zhou \(2015\)](#) and [Xu \(2018\)](#)). The convergence is uniform over all missing links, which implies that the imputation error is controlled at the link level and is particularly useful for applications that require accurate imputation of each link probability.

Following the classical nonparametric regression literature, we decompose the estimation error into bias and variance. The bias term is of order $(h^2 + 1/(nh^{d_\zeta}))$, where (i) h^2 arises because our local two-way fixed-effects regression removes the first-order bias, leaving only the second-order approximation error, and (ii) $1/(nh^{d_\zeta})$ reflects the contamination from diagonal entries of the matrix A (which are 0

instead of $f_{ii} + \epsilon_{ii}$ because we rule out self-loops). In most empirically relevant cases, the first term, h^2 , is dominant. In particular, under the MSE-optimal bandwidth choice, it can be directly verified that $1/(nh^{d_\zeta}) = o(h^2)$.

The variance consists of two terms. The first term, $\log N/(nh^{d_\zeta})$, matches the rate in the standard nonparametric literature. The second term, $\delta_{N,n}h$, is special to our estimator and arises from the use of the pseudo-distance as a substitute for the unobserved latent factors ζ_i and ζ_j . When the latent factors are observable, we have $\delta_{N,n} = 0$, and the second term disappears, reducing our result to cases [Tsybakov \(2008\)](#) and [Graham, Niu and Powell \(2021\)](#).

The bandwidth h governs the bias–variance trade-off. However, because the bias and variance each consist of two components, the trade-off deviates from the classical setting. As h decreases, h^2 and $h^2\delta_{N,n}^2$ diminish, while $1/(nh^{d_\zeta})$ increases. Therefore, the optimal choice of h depends jointly on the relative magnitudes of n and N and on d_ζ . Nevertheless, it is straightforward to verify that, when n and N are of the same order, setting $h \asymp n^{-1/(4+d_\zeta)}$ attains the minimum MSE.

Remark Our method employs two-way fixed-effects regression for imputation, which effectively removes the first-order approximation error. The intuition behind this approximation can be illustrated by a Taylor expansion. Heuristically, to impute A_{ij} , if we ignore the first-step nonparametric dyadic regression and consider the Taylor expansion of $f(\zeta_{i'}, \zeta_{j'})$ around the fixed point (ζ_i, ζ_j) :

$$\begin{aligned} f(\zeta_{i'}, \zeta_{j'}) &= \underbrace{f(\zeta_i, \zeta_j) + (\nabla_{\zeta_i} f(\zeta_i, \zeta_j))' (\zeta_{i'} - \zeta_i)}_{a_{i'}} + \underbrace{(\nabla_{\zeta_j} f(\zeta_i, \zeta_j))' (\zeta_{j'} - \zeta_j)}_{b_{j'}} + O(\|\zeta_{i'} - \zeta_i\|^2 + \|\zeta_{j'} - \zeta_j\|^2) \\ &= a_{i'} + b_{j'} + O(\|\zeta_{i'} - \zeta_i\|^2 + \|\zeta_{j'} - \zeta_j\|^2) \end{aligned}$$

Because we employ a kernel-weighted estimator based on the pseudo-distances $\hat{d}_{ii'}, \hat{d}_{jj'}$ with bandwidth h , and since, as discussed above, $\hat{d}_{ii'}$ and $\hat{d}_{jj'}$ differ from the latent-space distances $\|\zeta_{i'} - \zeta_i\|$ and $\|\zeta_{j'} - \zeta_j\|$ only up to a bounded factor, the contribution to imputation comes from links (i', j') with $\hat{d}_{ii'}, \hat{d}_{jj'} \leq h$. For those links, we obtain

$$f(\zeta_{i'}, \zeta_{j'}) = a_{i'} + b_{j'} + O(h^2)$$

Hence, the two-way fixed-effects regression absorbs all first-order approximation errors, leaving only the second-order remainder $O(h^2)$.

The idea of using two-way fixed-effects imputation is closely related to the approaches developed in [Freeman and Weidner \(2023\)](#) and [Beyhum and Mugnier \(2024\)](#). Our approach differs from theirs in two respects. First, their methods do not involve missing data and are primarily designed to estimate structural parameters in panel data models, whereas our method aims to predict the formation probability of missing links. Second, they discretize the data into groups using k-means clustering, while we employ a kernel estimator.

The methods most conceptually related to ours are those of [Feng \(2023\)](#) and [Deaner, Hsiang and Zelenev \(2025\)](#). [Deaner, Hsiang and Zelenev \(2025\)](#) propose a local-constant estimator to impute potential outcomes. Because they assume only that the potential outcome function is Lipschitz continuous, their approximation error is of order $O(h)$. Building on the ideas of [Freeman and Weidner \(2023\)](#) and [Beyhum and Mugnier \(2024\)](#), we assume that the graphon is twice continuously differentiable and, by using a two-way fixed-effects imputation, we reduce the approximation error to $O(h^2)$. In comparison, the local PCA method proposed by [Feng \(2023\)](#) leverages factor models together with higher-order Taylor expansions to achieve more accurate approximations. In particular, [Feng \(2023\)](#) show that the local PCA estimator attains Stone’s minimax-optimal rate for nonparametric regression. We refer interested

readers to that paper for further details.

We adopt the two-way fixed-effects model instead of local PCA approach for three reasons. First, local PCA method requires prior knowledge of the dimension d_ζ , which is typically unknown in practice and rarely supported by prior information. In contrast, our approach does not require prior knowledge of d_ζ . Second, the higher-order smoothness assumption is often violated in empirically relevant settings. As discussed earlier, in many applied network formation models, the homophily effect depends on the absolute distance between characteristics, implying that the graphon is only Lipschitz continuous. Our approach is robust to such lack of smoothness and can still achieve first-order approximation accuracy in this case. Third, networks in most economic applications are relatively small. For instance, in development economics, village networks typically contain around 200 households, and with a sampling rate of roughly 40%, the effective sample size available for imputing a specific missing link can be fewer than 20. Under such small-sample conditions, achieving theoretical higher-order accuracy is difficult. Our numerical simulations also show that the finite-sample performance of our method is better to that of the local PCA.

4 Theory for Downstream Estimation

We consider the application of our method to estimating parameters in downstream economic models. The data $\{(A_m, X_m, W_m, Y_m)\}_{m=1}^M$ consist of M large i.i.d. networks indexed by $m = 1, 2, \dots, M$, and each network has size N_m . Here, $A_m \in \{0, 1\}^{N_m \times N_m}$ is the adjacency matrix, $Y_m \in \mathbb{R}^{N_m}$ is the outcome variable, $X_m \in \mathbb{R}^{N_m \times d_X}$ denotes the covariates that directly enter the network formation model as in (2.1), whereas $W_m \in \mathbb{R}^{N_m \times d_W}$ contains variables that do not necessarily appear in the network formation model but affect the outcome Y_m . Note that W_m and X_m may overlap, i.e., $W_m \cap X_m \neq \emptyset$.

The adjacency matrix A_m is not fully observed because of egocentric sampling, but the outcome Y_m and the covariates (X_m, W_m) are observed for all individuals in the network. Let n_m be the number of sampled individuals in network m . We mainly focus on the asymptotics in which $M, N_m, n_m \rightarrow \infty$. The large-network asymptotics $N_m, n_m \rightarrow \infty$ are required for consistent imputation of the adjacency matrix, but letting $M \rightarrow \infty$ is not necessary; our analysis also applies to single large network.

We allow for heterogeneous network formation process f_m for each $m = 1, \dots, M$, so that information from other networks does not help in imputing a given network. Therefore, we implement imputation method separately for each incomplete network, obtain the imputed adjacency matrix $\hat{A}_m$, and use it, rather than the observed network A^{obs} , in the moment conditions to construct the GMM estimator.

4.1 Consistency

GMM We work with the moment function

$$\psi(A_m, X_m, W_m, Y_m, \alpha) : [0, 1]^{N_m \times N_m} \times \mathbb{R}^{N_m \times d_X} \times \mathbb{R}^{N_m \times d_W} \times \mathbb{R}^{d_\alpha} \mapsto \mathbb{R}^q$$

where q is the number of moment conditions and $\alpha \in \mathbb{R}^{d_\alpha}$ is the structural parameter of interest in the downstream analysis. We assume the following moment condition

$$\mathbb{E}(\psi(A_m, W_m, Y_m, \alpha_0)) = 0 \tag{4.1}$$

where α_0 is the true parameter. Since A_m is partially observed due to sampling, we impute each sampled network separately to obtain $\{\hat{A}_m \mid m = 1, \dots, M\}$, and then use the imputed network to solve:

$$\hat{\alpha} = \arg \min_{\alpha \in \text{supp}(\alpha)} \left(\frac{1}{M} \sum_{m=1}^M \psi(\hat{A}_m, W_m, Y_m, \alpha) \right)' \Sigma_M \left(\frac{1}{M} \sum_{m=1}^M \psi(\hat{A}_m, W_m, Y_m, \alpha) \right) \quad (4.2)$$

where $\Sigma_M \in \mathbb{R}^{q \times q}$ is the weight matrix.

Example 3 (Regression on centrality) A large literature studies the relationship between individuals' network positions (centrality) and their outcomes, for example, [Hochberg, Ljungqvist and Lu \(2007\)](#), [Cruz, Labonne and Querubin \(2017\)](#), and [Banerjee et al. \(2013\)](#). The regressions used in such studies can be written as:

$$Y_{mi} = \alpha_C + \alpha_1 \phi_{mi}(A_m) + e_{mi}, \quad i = 1, 2, \dots, N_m, \quad m = 1, 2, \dots, M \quad (4.3)$$

where $\alpha = (\alpha_C, \alpha_1)'$ is the parameter of interest, $\phi_{mi} : [0, 1]^{N_m \times N_m} \rightarrow \mathbb{R}$ computes the centrality of individuals i . For example:

- (i) **(Degree centrality)** For any matrix $\tilde{A}_m \in [0, 1]^{N_m \times N_m}$, if $\phi_{mi}(\tilde{A}_m) = \frac{1}{N_m} \sum_{j=1}^{N_m} \tilde{A}_{m,ij}$, then $\phi_{mi}(\cdot)$ represents the normalized degree of individual i when $\tilde{A}$ is an adjacency matrix.
- (ii) **(Eigenvector centrality)** For any symmetric matrix $\tilde{A}_m \in [0, 1]^{N_m \times N_m}$, let $\tilde{A}_m = U_m D_m U_m'$ be the eigenvalue decomposition of $\tilde{A}$, where $U_m \in \mathbb{R}^{N_m \times N_m}$ is an orthonormal matrix containing the eigenvectors of $\tilde{A}_m$ and $D_m \in \mathbb{R}^{N_m \times N_m}$ is a diagonal matrix with eigenvalues arranged in decreasing order. Define $\phi_{mi}(\tilde{A}_m) = \sqrt{N_m} U_{m,i1}$, where $U_{m,i1}$ denotes the i -th entry of the leading eigenvector. When $\tilde{A}_m$ is a symmetric adjacency matrix, $\phi_{mi}(\cdot)$ corresponds to the eigenvector centrality of individual i .

$\{e_{mi} \mid m = 1, \dots, M, i = 1, \dots, N_m\}$ contains the error terms that may be correlated within networks, so inference should be clustered at the network level. We use the moment function:

$$\psi(\hat{A}_m, W_m, Y_m, \alpha) := \frac{1}{N_m} \sum_{i=1}^{N_m} (1, \phi_{mi}(\hat{A}_m))' (Y_{mi} - \alpha_C - \alpha_1 \phi_{mi}(\hat{A}_m))$$

to obtain estimates of α . When the errors are independent within each network, our results extend to the case of a single large network (i.e., $M = 1$), and we still obtain a consistent estimator.

Example 4 (Linear-in-means peer-effects model) The linear-in-means peer-effects model is widely used to study peer influence in social networks. The model is:

$$Y_m = \alpha_C + \alpha_{\bar{Y}} G_m Y_m + \alpha_W W_m + \alpha_{\bar{W}} G_m W_m + e_m \quad (4.4)$$

where $\alpha = (\alpha_C, \alpha_{\bar{Y}}, \alpha'_W, \alpha'_{\bar{W}})'$, $\alpha_{\bar{Y}}$ captures the endogenous effect, and G_m is the row-normalized adjacency matrix⁴. $G_m Y_m$ is endogenous due to the reflection problem ([Manski \(1993\)](#)), the model must be estimated using instrumental variables. Under the identification conditions in [Bramoullé, Djebbari and](#)

⁴For any $i, j \in 1, \dots, N_m$, $G_{m,ij}$ is obtained through:

$$G_{m,ij} = \frac{A_{m,ij}}{\sum_{j'=1}^{N_m} A_{m,ij'}}$$

Fortin (2009), the following moment function can be constructed:

$$\psi(A_m, W_m, Y_m, \alpha) = \frac{1}{N_m} Z'_m (Y_m - \alpha_C - \alpha_{\bar{Y}} G_m Y_m - \alpha_W W_m - \alpha_{\bar{W}} G_m W_m)$$

where $Z_m = [1, W_m, G_m W_m, G_m^2 W_m]$ is the instrument. When the network A_m is incomplete, we compute $\hat{G}_m$ and $\hat{Z}_m$ using imputed network $\hat{A}$ ⁵, and use the moment function:

$$\psi(\hat{A}_m, W_m, Y_m, \alpha) = \frac{1}{N_m} \hat{Z}'_m (Y_m - \alpha_C - \alpha_{\bar{Y}} \hat{G}_m Y_m - \alpha_W W_m - \alpha_{\bar{W}} \hat{G}_m W_m)$$

to obtain α . Since $\{e_{mi} \mid m = 1, \dots, M, i = 1, \dots, N_m\}$ contains the error terms that may be correlated within networks, so inference should be clustered at the network level. However, when the errors are independent within each network, our results extend to the case of a single large network (i.e., $M = 1$), and we still obtain a consistent estimator.

Remark Our approach for GMM estimation differs from that of Chandrasekhar and Lewis (2011). Specifically, Chandrasekhar and Lewis (2011) construct moments by integrating over the missing network data, typically using simulation to approximate (conditional) expectations. As a consequence, their implementation is computationally demanding in GMM settings and relies on parametric assumptions about the distribution of the error terms (see the discussion in Chandrasekhar and Lewis (2011) and Boucher and Houndetoungan (2025)). By contrast, our approach directly plugs the imputed network into the moment function $\psi(\cdot)$, avoiding simulation and distributional assumptions on the errors. A key advantage of Chandrasekhar and Lewis (2011)'s approach is that it can accommodate complex network statistics, such as average path length, that cannot be computed using a simple plug-in strategy. However, even in this case, our imputed adjacency matrix can still serve as the basis for the simulation-based GMM procedure using the theoretical framework developed in Section 4.2 of Chandrasekhar and Lewis (2011).

Assumption 5 (GMM) Suppose that

- (i) (**Asymptotics**) Large- M asymptotics with $M \rightarrow \infty$.
- (ii) (**Independence**) $\{(A_m, X_m, W_m, Y_m)\}_{m=1}^M$ are i.i.d..
- (iii) (**Identification**) There is a unique α_0 such that $\mathbb{E}(\psi(A_m, W_m, Y_m, \alpha_0)) = 0$.
- (iv) (**Idiosyncrasy-robustness**) Let P_m be matrix of conditional probability of link formation, then as $N \rightarrow \infty$,

$$\sup_{1 \leq m \leq M} \sup_{\alpha} \|\psi(P_m, W_m, Y_m, \alpha) - \psi(A_m, W_m, Y_m, \alpha)\| = o_P(1)$$

- (v) $\Sigma_M \xrightarrow{P} \Sigma$ and Σ is positive definite.
- (vi) (**Compactness and integrable**) The support of α is compact, $\alpha \mapsto \psi(A_m, W_m, Y_m, \alpha)$ is continuous almost everywhere, and $\mathbb{E} \left(\sup_{\alpha \in \text{supp}(\alpha)} \|\psi(A_m, W_m, Y_m, \alpha)\|^2 \right) < \infty$.
- (vii) Let $\mathcal{S}_m$ be the set of sampled nodes in $\mathcal{A}$, then $\sup_{m \leq M} N_m^{-1} \|\hat{A}_{m, \mathcal{S}_m^c \mathcal{S}_m^c} - P_{m, \mathcal{S}_m^c \mathcal{S}_m^c}\|_F = o_P(1)$.

⁵For any $i, j \in 1, \dots, N_m$, $\hat{G}_{m,ij}$ is obtained through:

$$\hat{G}_{m,ij} = \frac{\hat{A}_{m,ij}}{\sum_{j'=1}^{N_m} \hat{A}_{m,ij'}}$$

and $\hat{Z}_m = [1, W_m, \hat{G}_m W_m, \hat{G}_m^2 W_m]$

(viii) (**Lipschitz**) For any $P_{m1}, P_{m2} \in [0, 1]^{N \times N}$, we have:

$$\sup_{\alpha \in \text{supp}(\alpha)} \|\psi(P_{m1}, W_m, Y_m, \alpha) - \psi(P_{m2}, W_m, Y_m, \alpha)\| \leq L(W_m, Y_m) N_m^{-1} \|\hat{P}_{m1} - P_{m2}\|_F$$

$$\text{and } \mathbb{E} L^2(W_m, Y_m) < \infty.$$

Assumption 5(i) and (ii) are standard to ensure the consistency of M-estimator. However, if we are willing to assume that the error terms are independent within networks, then these two assumptions are no longer needed, and our analysis can apply to studying single large network; see the discussion in Examples 3 and 4.

Assumption 5(iii) is a standard identification condition and ensures that the moment condition does not hold at $\alpha \neq \alpha_0$. However, in the large-network asymptotic regime, weak identification becomes a concern. For example, in Example 4, if W is i.i.d. within each network and independent of the network (e.g., random treatment assignment), the instrument Z becomes asymptotically collinear, leading to an inconsistent estimator (see Hayes and Levin (2024) for further discussion). In the following analysis, we therefore assume that W is correlated with X to avoid the weak-identification problem.

Assumption 5(iv) implies that the difference between probability matrix P and the adjacency matrix A in moment conditions becomes asymptotically negligible as the network size N grows. This assumption is key for establishing the consistency of our plug-in estimator: because our imputation can only approximate probability matrix P instead of missing links, our estimator is consistent only if replacing A with P leaves the moment condition unchanged asymptotically. Since A decomposes into probability matrix P , which is predictable from observed information, and an unpredictable error matrix, this assumption requires that the contribution of the unpredictable error matrix vanishes asymptotically. This condition is satisfied in many empirical applications using network data, such as the linear-in-means peer-effects model and linear regressions on network centralities, because network statistics aggregate a large number of links and naturally denoise link-level errors (see the discussion in the following text).

Assumption 5(v) and (vi) are standard conditions for establishing the consistency of M-estimators. Assumption 5(vii) states that, for each network m , the imputed network $\hat{A}_m$ is consistent for the probability matrix P_m on the missing subnetwork in the Frobenius norm. This condition is weaker than the conclusion in Theorem 4, as the latter establishes consistency in the sup norm for the missing submatrix.

Assumption 5(viii) imposes a Lipschitz condition requiring that perturbations in P_m lead to changes in ψ up to a factor $L(W_m, Y_m)$ with a finite second moment, uniformly over α . Since our parameter of interest is α and P_m serves as a nuisance parameter, this assumption can be interpreted as a smoothness condition on the nuisance parameters within the framework of semi-parametric M-estimation.

Example 3 (Continued) We discuss key conditions, Assumption 5(iv) and (viii), for linear regressions on degree centrality and eigenvector centrality, respectively. For degree centrality, verification is straightforward. Because degree centrality is simply the row average of the adjacency matrix, the difference between moments constructed from the observed adjacency matrix A_m and the underlying probability matrix P_m is proportional to the row average of error terms. Since the row average of errors converges at rate $O_p(1/\sqrt{N_m})$, Assumption 5 follows immediately. A similar argument, together with the Cauchy-Schwarz inequality, verifies Assumption 5(viii).

For eigenvector centrality, we impose a standard eigen-gap condition requiring that the largest eigenvalue of P_m is separated from the second-largest eigenvalue uniformly across networks. Such conditions are common in spectral analysis and ensure that the leading eigenspace is well separated. Under this assumption, the Davis-Kahan theorem Yu, Wang and Samworth (2015) implies that the perturbation in eigenvector centrality is proportional to the normalized spectral norm of the error matrix, i.e., $\|E\|_{\text{op}}/N_m$,

which is of order $1/\sqrt{N_m}$ by [Bandeira and Van Handel \(2016\)](#). This establishes Assumption 5(iv). Assumption 5(viii) follows by a similar argument.

Example 4 (Continued) For the linear-in-means peer-effects model, if each element in W_m is uniformly bounded and the strict exogeneity condition $\mathbb{E}(e_{m,i} \mid W_m) = 0$ holds for every i , then Assumptions 5(iv) and (viii) can be verified by arguments analogous to those used for the centrality regressions.

Theorem 3 (Consistency of estimator) Under Assumption 5, the estimator $\hat{\alpha}$ defined as in (4.2) is consistent for α_0 .

Theorem 3 is particularly relevant in empirical applications, where interest typically focuses on the consistency of parameter estimates constructed from the imputed network rather than on the accuracy of the imputed links themselves. This theorem shows that the plug-in estimator based on our imputed adjacency matrices is consistent. The key advantages of our approach are that it is robust under misspecification of (i) the network formation model, (ii) the dimension of latent heterogeneity, and (iii) the smoothness of the network formation process. Moreover, even when a researcher prefers to impose a specific parametric network formation model, for computational simplicity or due to strong prior knowledge, our nonparametric procedure still provides a useful benchmark for assessing how sensitive the parameter estimates are to the parametric assumptions and for diagnosing potential misspecification in the imposed network formation model.

4.2 Analysis of linear-in-means peer-effect model

Although Theorem 3 establishes the consistency of the plug-in estimator based on our imputed adjacency matrices, analyzing its convergence rate and inference properties is challenging. First, as discussed in [Cai \(2022\)](#), even when each link contains only i.i.d. classical measurement error, network statistics used in moment conditions typically aggregate errors across links, thereby inducing non-classical measurement error that complicates the analysis. Second, the measurement error generated by our imputation procedure is correlated across links, and because the imputation is nonparametric, its bias and variance may be of the same order, further complicating the analysis of convergence.

Since network links typically enter the moment conditions through network statistics, the asymptotic properties of imputed links established in Theorem 2 could be different from those of the GMM estimators that rely on the imputed network. Therefore, the bandwidth that is optimal for imputing individual links may not be optimal for downstream analysis using network statistics. In the following text, we focus on the linear-in-means peer-effects regression and study the asymptotic behavior of its parameter estimates. We also provide guidance for downstream regression analysis using our imputed networks.

Consider the linear-in-means peer-effects model (4.4) in Example 4, given the weight matrix Σ_M . Let $\hat{V}_m = [\mathbf{1}, \hat{G}_m Y_m, W_m, \hat{G}_m W_m]$, $\hat{Z}_m = [\mathbf{1}, W_m, \hat{G}_m W_m, \hat{G}_m^2 W_m]$, and $\alpha = (\alpha_C, \alpha_{\bar{Y}}, \alpha'_W, \alpha'_{\bar{W}})'$ the GMM estimator is:

$$\hat{\alpha} = \left(\frac{1}{M} \sum_{m=1}^M \hat{V}_m' \hat{Z}_m \Sigma_m \hat{Z}_m' \hat{V}_m \right)^{-1} \left(\frac{1}{M} \sum_{m=1}^M \hat{V}_m' \hat{Z}_m \Sigma_m \hat{Z}_m' Y_m \right) \quad (4.5)$$

Assumption 6 Suppose that

- (i) **(Sampling)** $\{(A_m, X_m, W_m, Y_m)\}_{m=1}^M$ are i.i.d., and the data is generated according to Example 4.
- (ii) **(Exogeneity)** The error vector e_m satisfies $\mathbb{E}(e_m \mid \{X_{m,i}, \xi_{m,i}\}_{i=1}^{N_m}, W_m) = 0$. In addition, we assume that $e_m \perp \epsilon_m \mid (\{X_{m,i}, \xi_{m,i}\}_{i=1}^{N_m}, W_m)$, where $\epsilon_m := \{\epsilon_{m,ij} \mid 1 \leq i < j \leq N_m\}$ are error terms in the network generating process (2.1).

(iii) (**Asymptotics**) The network sizes $\{N_m \mid m = 1, \dots, N\}$ are of the same order, and sample sizes $\{n_m \mid m = 1, \dots, M\}$ are of the same order as well, i.e.,

$$\sup_{M \rightarrow \infty} \inf_{1 \leq m_1, m_2 \leq M} (N_{m_1}/N_{m_2}) > 0, \quad \sup_{M \rightarrow \infty} \inf_{1 \leq m_1, m_2 \leq M} (n_{m_1}/n_{m_2}) > 0$$

(iv) (**Compactness**) The support of $W_{m,i}$ is uniformly bounded over i and m .

(v) (**Non-degeneracy**) Let $\lambda_{\min}(\cdot)$ denote the smallest singular value of a matrix, then,

$$\lambda_{\min} \left(\mathbb{E} \left(\frac{1}{N_m} Z'_m W_m \right)' \mathbb{E} \left(\frac{1}{N_m} Z'_m W_m \right) \right) > 0$$

Assumption 6(ii) imposes the strict exogeneity condition and rules out correlated effects (Bramoullé, Djebbari and Fortin, 2020). Note that Assumption 6(ii) is stronger than the standard assumption in the literature, i.e., $\mathbb{E}(e_m \mid G_m, W_m) = 0$. This is because the network data are incomplete in our setting, This is because the network data are incomplete in our setting, and we therefore require the error term to be orthogonal to the information used for network imputation. We further assume that the error terms in the linear-in-means peer-effects model, e_m , are conditionally independent of the errors in the network formation process, ϵ_m , which ensures that the imputation error is independent of Y_m .

Assumption 6(iii) requires that all networks have sizes of the same order and that the sample sizes within networks are also of the same order. In other words, no network or sample becomes disproportionately large or disproportionately small.

Assumption 6(v) imposes a non-degeneracy condition requiring that the smallest singular value of $\mathbb{E} \left(\frac{1}{N_m} Z'_m W_m \right)' \mathbb{E} \left(\frac{1}{N_m} Z'_m W_m \right)$ is bounded away from zero. This requirement is analogous to the relevance condition in instrumental-variable estimation, and is particularly important under large-network asymptotics. For example, if W_m is i.i.d. within each network and independent of the network (e.g., random treatment assignment), the instrument Z_m becomes asymptotically collinear, causing the smallest singular value of the above matrix to shrink toward zero and leading to an inconsistent IV estimator. In practice, one can let W_m to be correlated with X_m to satisfy this assumption (see Hayes and Levin (2024) for further discussion).

Theorem 4 Under Assumption 6 and conditions in Theorem 2, let $\bar{N} := \frac{1}{M} \sum_{m=1, \dots, M} N_m$ and $\bar{n} := \frac{1}{M} \sum_{m=1, \dots, M} n_m$, we have

$$\sqrt{M}(\hat{\alpha} - \alpha_0 - b) \xrightarrow{d} \mathcal{N}(0, \Omega + v)$$

where $b = O \left(h^2 + \frac{\log \bar{N}}{\bar{n}^2 h^{2d_\zeta}} \right)$, $v = O \left(\delta_{\bar{N}\bar{n}}^2 h^2 + \frac{\log \bar{N}}{\bar{n}^2 h^{2d_\zeta}} \right)$.

The theorem states that $\hat{\alpha} - \alpha_0$ is asymptotically normally distributed, but its limiting distribution is centered away from zero by an $O \left(h^2 + \frac{\log \bar{N}}{\bar{n}^2 h^{2d_\zeta}} \right)$ bias term. The first component, h^2 , arises from the bias of imputed links, while the second term is, in most cases, asymptotically negligible relative to h^2 in most empirically relevant cases⁶. The asymptotic variance can be decomposed into two components: Ω corresponds to the variance that would obtain if the network A were perfectly observed, and v captures the additional variability introduced by the nonparametric estimation of $\hat{A}$. Since the variance component

⁶To see this, note that the optimal bandwidth for minimizing the mean squared error of the imputed links is of order $h^* \sim n^{\frac{2}{4+d_\zeta}}$. For the second bias component to dominate the first, that is, $h^2 = o \left(\frac{1}{\bar{n}^2 h^{2d_\zeta}} \right)$, we would need $h = o \left(n^{-\frac{1}{1+d_\zeta}} \right)$, which is much smaller than $(h^*)^2$. Such undersmoothing is rare in practice.

induced by using the imputed network shrinks at rate $1/\sqrt{M}$, whereas the bias does not, this highlights a difference between the asymptotic behavior of the estimator and that of the link-level imputation error.

To illustrate the difference between the asymptotic behavior of the estimator and that of the link-level imputation error, consider the case $d_\zeta \geq 2$ and the optimal bandwidth for minimizing the MSE of the imputed links, i.e., $h \sim n^{-2/(4+d_\zeta)}$. The bias of $\hat{\alpha}$ is of order $O(h^2)$, which is of the same order as the bias of the link imputation error in most empirically relevant cases. In contrast, the additional variance induced by the imputed network decreases at the rate $1/M$ relative to the variance of the link imputation error.

These theoretical observations suggest that reducing the MSE of $\hat{\alpha}$ requires undersmoothing in the imputation step, that is, using a smaller bandwidth h to reduce the bias of the imputed links, rather than attempting to balance the bias and variance of the link-level imputation error. It is worth noting that the existing nonparametric inference literature typically employs undersmoothing for the purpose of constructing valid confidence intervals. In contrast, in our setting, undersmoothing is required even when the objective is to minimize the MSE of $\hat{\alpha}$. This requirement arises from the different asymptotic behaviors of network statistics and link imputation errors.

However, in practice, the asymptotic analysis above has two limitations. First, network sizes are relatively small in most economics applications, so reducing h leaves very few observations available for local imputation. Second, the number of networks M is also small and rarely exceeds 40 in most empirical settings. For these reasons, we recommend that applied researchers start from the optimal bandwidth h selected by cross-validation in the imputation step, then gradually decrease h and examine the corresponding estimates $\hat{\alpha}$. Once the estimates become unstable as h decreases, we should stop undersmoothing at that point.

5 Simulation Study

In this section, we examine the finite-sample properties of the proposed method using a series of numerical experiments. We compare its performance with several alternative imputation approaches, including the low-rank imputation methods proposed by [Bai and Ng \(2021\)](#) and [Li et al. \(2023\)](#), and the local PCA method developed by [Feng \(2023\)](#). We also use numerical experiments to illustrate (i) the importance of incorporating observed covariates for improving prediction accuracy, and (ii) the influence of sample splitting on finite-sample performance. We begin with introducing alternative method.

5.1 Alternative Methods

Covariate-only method This method is a dyadic nonparametric regression-based imputation approach that relies solely on the observed covariates X . When unobserved heterogeneity is present, the prediction error of this approach does not diminish as the sample size increases. We compare its numerical performance with that of our proposed method to highlight the importance of accounting for unobserved heterogeneity. We refer to this method as **X** in the following text.

Low-rank imputation When the observed matrix follows a strong-factor structure with an additive noise component and the missing entries exhibit a block-missing pattern⁷ (see (2.3)), [Bai and Ng \(2021\)](#) and [Li et al. \(2023\)](#) propose using low-rank estimation to impute the missing data. The key idea is to first recover the latent factors using the observed submatrices A_{SS} , A_{SS^c} , and A_{S^cS} , and then reconstruct the missing bottom-right submatrix $A_{S^cS^c}$ through the product of the estimated factors.

⁷[Bai and Ng \(2021\)](#) refer to this pattern as a tall-wide structure in the context of panel data, while [Li et al. \(2023\)](#) study an analogous setting in network data and refer to it as egocentric sampling.

When the graphon f is nonlinear as in Example 2, the graphon matrix is typically high-rank and thus violates the low-rank assumption required by Bai and Ng (2021) and Li et al. (2023). Although their methods are not theoretically valid in such settings, we include them in our numerical experiments for comparison, given their simplicity and computational efficiency, to assess how our proposed method performs in practice relative to these benchmarks. It is worth noting that both Bai and Ng (2021) and Li et al. (2023) assume that the true number of factors is known a priori. In practice, this information is rarely available. Hence, in our numerical experiments, we select the number of factors via cross-validation.

Local PCA The local PCA method proposed by Feng (2023) is suitable for high-rank graphon. Specifically, it first finds the neighbors of each node i based on the pseudo-distance using k -nearest neighbors (kNN), and then performs low-rank imputation on the submatrix formed by node i 's neighbors to impute the missing entries. As discussed earlier, when the underlying graphon is smoother than twice continuously differentiable, the local PCA method can, in theory, achieve a higher-order approximation error. We compare our method with local PCA to assess whether this theoretical advantage persists in finite samples.

A practical concern is that the local PCA method requires prior knowledge of the latent dimension d_ζ , since this determines the rank when performing PCA on each submatrix. However, d_ζ is typically unknown and rarely supported by prior knowledge. While Feng (2023) propose an ad hoc procedure in their numerical simulations to address this issue, we instead use cross-validation to jointly select both the number of neighbors k in the kNN step and the rank used in PCA.

It is worth noting that local PCA does not incorporate observed covariates⁸. In our numerical experiments, we therefore consider two implementations of their method: (1) **LPCA**, which ignores covariates X and applies local PCA directly to the adjacency matrix, and (2) **X-LPCA**, which first removes the variation explained by X and then applies local PCA to the residual matrix.

Local TWFE without covariates We also consider an alternative method that directly applies our local TWFE approach to the adjacency matrix without extracting the variation explained by X . As discussed earlier, because the information about ζ recovered from observed data is noisy, it is unwise to ignore the observed covariates and absorb them into the latent factors. We include the numerical simulation results of this method to evaluate the importance of incorporating covariate information. We refer to this method as **LTWFE** and to our proposed method as **X-LTWFE** in the following text.

Sample splitting While our asymptotic analysis relies on sample splitting for theoretical tractability, this requirement is primarily technical. In practice, sample splitting reduces the effective sample size, and since network sizes are relatively small in most economics applications, its finite-sample performance can be worse than that of the full-sample implementation. Accordingly, we implement all alternative methods without sample splitting.

For our proposed method, we consider both versions, with and without sample splitting, to examine how this choice affects finite-sample performance. We refer to our method with sample splitting as **X-LTWFE-SP** in the following text.

Summary We summarize all imputation methods used in the numerical experiments below:

- (i) **(X)** Covariate-only nonparametric method that imputes missing links using X alone.
- (ii) **(LR)** The low-rank imputation methods of Bai and Ng (2021) and Li et al. (2023).

⁸Feng (2023) briefly discuss how to include covariates in a linear form in their appendix.

- (iii) **(LPCA)** The local PCA method that ignores covariates.
- (iv) **(X-LPCA)** First removes the variation explained by X and then applies local PCA to the residual matrix.
- (v) **(LTWFE)** The local two-way fixed-effects method that ignores covariates.
- (vi) **(X-LTWFE)** Our proposed method. It first removes the variation explained by X and then applies the local two-way fixed-effects regression to the residual matrix.
- (vii) **(X-LTWFE-SP)** Our proposed method with sample-splitting.

5.2 Simulation for Imputation Accuracy

The network in the simulation is generated as

$$A_{ij} = \mathbf{1}(\omega(X_i, X_j)' \beta + g(\xi_i, \xi_j) - U_{ij} \geq 0) \quad (5.1)$$

$$\omega(X_i, X_j)' = ((X_{i1} - X_{j1})^2, (X_{i2} - X_{j2})^2), \quad g(\xi_i, \xi_j) = \xi_{i1} + \xi_{j1} - \frac{1}{8}(\xi_{i2} - \xi_{j2})^2$$

The dimensions of the observable and latent characteristics are $d_X = d_\xi = 2$, and $\{(X_i, \xi_i) \mid i = 1, \dots, N\}$ is i.i.d. across individuals. Covariates and latent factors are generated according to

$$X_{id} = \frac{1}{2}(\xi_{i1} + \xi_{i2}) + \epsilon_{X,id}, \quad \forall d = 1, 2$$

and

$$\xi_i \sim \mathcal{N}\left(0, \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}\right), \quad \epsilon_{X,i1}, \epsilon_{X,i2} \sim \text{Uniform}([-1, 1]), \quad \epsilon_{X,i1} \perp \epsilon_{X,i2}$$

The idiosyncratic shocks $\{U_{ij}\}_{1 \leq i < j \leq N}$ are independently drawn from a type-I extreme-value distribution. The coefficient vector β captures homophily effects on observables and takes values $\beta \in \{(-0.5, 0.5), (-2, -2)\}$. We vary the magnitude of β to control the sparsity of the network: for example, when $\beta = (-0.5, 0.5)$, the resulting network tends to be denser, with a larger number of links being formed.

The network size is $N = 200$, and the simulations consider sampling rates $\varphi := n/N \in \{0.2, 0.3, 0.4, 0.5, 0.6, 0.8\}$ to compare the performance of the proposed and alternative methods under different sampling scenarios. We conduct $S = 1,000$ Monte Carlo replications in the numerical experiment, and for each replication s , we construct the imputed network $\hat{A}_s$ and compute the mean squared error (MSE) between $\hat{A}_s$ and the underlying probability P_s on the missing subnetwork, namely $\frac{1}{(N-n)(N-n-1)} \sum_{i,j \in S^c} (\hat{A}_{s,ij} - P_{s,ij})^2$. The reported statistic is the root mean-squared error (RMSE):

$$\text{RMSE} = \sqrt{\frac{1}{S} \sum_{s=1}^S \frac{1}{(N-n)(N-n-1)} \sum_{i,j \in S^c} (\hat{A}_{s,ij} - P_{s,ij})^2} \quad (5.2)$$

Table 1: Simulation: Imputation

	X ($\times 0.01$)	LR ($\times 0.01$)	LPCA ($\times 0.01$)	LTWFE ($\times 0.01$)	X-LPCA ($\times 0.01$)	X-LTWFE ($\times 0.01$)	X-LTWFE-SP ($\times 0.01$)
$\beta = (-0.5, -0.5)$							
$\varphi = 0.2$	21.0	15.8	15.9	15.3	14.9	10.9	13.1
$\varphi = 0.3$	20.9	13.9	14.0	13.9	12.3	9.7	11.3
$\varphi = 0.4$	20.8	12.4	12.5	13.0	10.8	8.9	10.4
$\varphi = 0.5$	20.8	11.3	11.3	12.3	9.8	8.4	9.7
$\varphi = 0.6$	20.8	10.3	10.4	11.9	9.2	8.1	9.3
$\varphi = 0.8$	20.6	9.2	9.2	11.0	8.3	7.5	8.6
$\beta = (-2, -2)$							
$\varphi = 0.2$	13.8	16.3	16.4	15.3	12.7	10.2	11.3
$\varphi = 0.3$	13.7	14.8	14.9	14.0	11.5	9.4	10.3
$\varphi = 0.4$	13.7	13.3	13.5	13.1	10.5	8.9	9.7
$\varphi = 0.5$	13.6	11.7	11.8	12.5	9.8	8.6	9.3
$\varphi = 0.6$	13.6	10.5	10.6	11.9	9.3	8.3	9.0
$\varphi = 0.8$	13.5	9.2	9.2	11.0	8.4	7.8	8.5

Note: We conduct Monte Carlo simulations based on 1,000 repetitions of the network formation model (5.1). The network size is $N = 200$. We set $\beta \in \{(-0.5, -0.5), (-2, -2)\}$ and sampling rates $\varphi \in \{0.2, 0.3, 0.4, 0.5, 0.6, 0.8\}$. We report RMSE as in (5.2) for the following imputation methods: (1) covariate-only nonparametric imputation that imputes missing outcomes using X alone (**X**), (2) the low-rank imputation methods of Bai and Ng (2021) and Li et al. (2023) (**LR**), (3) the local PCA method that ignores covariates (**LPCA**), (4) the local PCA method using covariates (**X-LPCA**), (5) the local two-way fixed-effects regression that ignores covariates (**LTWFE**), (6) the local two-way fixed-effects regression using covariates (**X-LTWFE**), and (7) X-LTWFE method with sample-splitting (**X-LTWFE-SP**). It is important to note that LR, LPCA, LTWFE, X-LPCA, and X-LTWFE are implemented without sample splitting. RMSEs are presented in units of 0.01 for ease of reading.

We evaluate the performance of our imputation method and alternative approaches in both dense networks (e.g., $\beta = (-0.5, -0.5)$, for which the average probability of forming a link is about 34%) and sparse networks (e.g., $\beta = (-2, -2)$, for which the average probability of forming a link is about 10%). The simulation results are reported in Table 1, with RMSEs presented in units of 0.01 for ease of reading.

The first column (X) reports the RMSE of covariate-only method, which imputes missing links using X alone. Because this method ignores unobserved heterogeneity, its imputations are systematically biased, and the error does not vanish as the network size or sampling rate increases. Consistent with the theoretical prediction, the numerical results show that, as the sampling rate φ increases, the RMSE of the covariate-only method exhibits very limited improvement (from 0.210 to 0.206 in the dense scenario, and from 0.138 to 0.135 in the sparse scenario).

The second column (LR) reports the RMSE of the low-rank imputation, and the third column (LPCA) reports the RMSE of the local PCA. Since the underlying graphon in (5.1) is high-rank, local PCA should, in theory, outperform the low-rank approach. However, the numerical results indicate that the RMSEs of the two methods are nearly the same across all scenarios. We conjecture that this pattern is driven by two reasons: (1) Although the graphon is not exactly low-rank, it is approximately low-rank, so the low-rank approach still provides a good approximation to the graphon matrix, especially when performance is measured using Frobenius norm. (2) Local PCA relies only on the subnetwork formed by neighbors, which leads to substantial sample loss and limits its performance in finite samples. In line with our theory, the local PCA estimator outperforms the covariate-only (X) method in most cases, and

its prediction error decreases as the sampling rate increases.

The fourth column (LTWFE) reports the RMSE of the local two-way fixed effects imputation that ignores covariates. In line with the theory, its prediction error decreases as n increases. Table 1 shows that at low sampling rates, the LTWFE method performs better than local PCA, whereas once the sampling rate becomes large (e.g., when φ exceeds 40%), local PCA outperforms LTWFE. This suggests that although local PCA has better asymptotic properties, in the sense that it achieves higher-order approximation error, it requires a larger sample size to realize this advantage, whereas LTWFE is more robust in small samples. In addition, the LTWFE method is more robust to network sparsity: as shown in Table 1, its performance is nearly unaffected by changes in the sparsity of the adjacency matrix.

The fifth column (X-LPCA) and the sixth column (X-LTWFE) report the performance of LPCA and LTWFE when incorporating covariate information. These methods first extract the variation of the missing links that can be explained by X , and then apply LPCA and LTWFE to the residual matrix, respectively. Compared with LPCA and LTWFE, incorporating covariate information leads to substantial improvements in prediction accuracy, particularly for X-LTWFE. The gains are especially large when the sampling rate is low or when the network is sparse. X-LPCA exhibits a similar pattern. In addition, Table 1 shows that the proposed method outperforms all alternatives in our simulations, and its performance remains robust even under sparse networks and low sampling rates, which are common in empirical applications.

The last column (X-LTWFE-SP) reports the performance of our proposed estimator with sample splitting. As discussed earlier, although sample-splitting facilitates asymptotic analysis, it reduces the effective sample size, and this issue can be severe in small-sample settings. In addition, it is reasonable to expect (though we do not formally prove it) that the correlation between the pseudo-distance and the error terms diminishes, making sample splitting unnecessary in theory. Table 1 shows that performance of proposed imputation under sample splitting is always worse than the performance without splitting the sample, but the difference between the two methods becomes smaller as the sample size increases.

In summary, the simulations show that our proposed method outperforms the alternative methods and is more robust across different levels of network sparsity and sampling rates. In addition, for finite samples, we recommend using the version without sample splitting in order to achieve higher prediction accuracy. Further numerical results are provided in the Appendix.

5.3 Simulation for estimates of parameter

In this subsection, we evaluate the performance of parameter estimates in downstream analyses that rely on the imputed networks. We compare the performance of our method with the alternative approaches introduced previously. We focus on two widely used classes of empirical exercises in economics that are based on network data.

Regression on network statistics Suppose for each network $m = 1, \dots, M$, the outcome Y_{mi} is generated according to:

$$Y_{mi} = \alpha_0 + \alpha_1 \phi_{mi}(A_m) + u_m + e_{mi}, \quad u_m \sim \mathcal{N}(0, 1/4), \quad e_{mi} \sim \mathcal{N}(0, 1/4) \quad (5.3)$$

where $\phi_{mi} : [0, 1]^{N_m \times N_m} \rightarrow \mathbb{R}$ computes a network statistic for individual i in network M , $\{u_m\}_{1 \leq m \leq M}$ are i.i.d. network-level random effects independent of A_m , and e_{mi} is i.i.d. idiosyncratic error. We consider two specifically, normalized average degree and eigenvector centrality, as defined in Example 3. The networks $\{A_m\}_{1 \leq m \leq M}$ according to (5.1) with $\beta = (-0.5, -0.5)$, which yields relatively dense networks with a larger number of links.

We set the number of networks to $M = 40$ and fix network size at $N_m = 200$ for all m . The simulations consider sampling rates $\varphi := n/N \in \{0.2, 0.3, 0.4, 0.5\}$ to compare the performance of the proposed method and alternative methods under different levels of sampling. For reference, we also report the performance of regressions using the complete networks (**CD**). We set $\alpha_0 = 0$ and $\alpha_1 = 0.5$.

We conduct $S = 1,000$ Monte Carlo replications. For each replication s , we generate M independent network, impute each network, and estimate the linear regression. The bias and standard deviation of $\hat{\alpha}_1$ are reported in Table 2.

Linear-in-mean peer-effects model Suppose for each network $m = 1, \dots, M$, the outcome Y_{mi} is generated according to:

$$\begin{aligned} Y_m &= \alpha_C + \alpha_{\bar{Y}} G_m Y_m + \alpha_W W_m + \alpha_{\bar{W}} G_m W_m u_m + e_m \\ W_{mi,d} &= \xi_{mi,d} + \frac{1}{2} X_{mi,d} \xi_{mi,d}, \quad d = 1, 2 \\ u_m &\sim \mathcal{N}(0, 1/25), \quad e_{mi} \sim \mathcal{N}(0, 1) \end{aligned} \tag{5.4}$$

where $\{u_m\}_{1 \leq m \leq M}$ are i.i.d. network-level random effects independent of A_m , and e_{mi} is i.i.d. idiosyncratic error. The networks $\{A_m\}_{1 \leq m \leq M}$ according to (5.1) with $\beta = (-0.5, -0.5)$, which yields relatively dense networks with a larger number of links.

We set the number of networks to $M = 40$ and fix network size at $N_m = 200$ for all m . The simulations consider sampling rates $\varphi := n/N \in \{0.2, 0.3, 0.4, 0.5\}$ to compare the performance of the proposed method and alternative methods under different levels of sampling. For reference, we also report the performance of regressions using the complete networks (**CD**). We set $\alpha_C = 0$, $\alpha_{\bar{Y}} = 0.5$, $\alpha_W = \alpha_{\bar{W}} = (1, 1)$. We conduct $S = 1,000$ Monte Carlo replications. For each replication s , we generate M independent network, impute each network, and estimate the GMM estimator using (4.5) with $\Sigma_M = \mathbf{I}_q$. The bias and standard deviation of estimate of endogenous effect $\hat{\alpha}_{\bar{Y}}$ are reported in Table 3.

Table 2: Simulation: Linear regression on centrality

	CD ($\times 0.01$)	X ($\times 0.01$)	LR ($\times 0.01$)	LPCA ($\times 0.01$)	LTWFE ($\times 0.01$)	X-LPCA ($\times 0.01$)	X-LTWFE ($\times 0.01$)	X-LTWFE-SP ($\times 0.01$)
Panel A: Degree centrality								
$\varphi = 0.2$								
BIAS	-0.05	9.62	-3.54	-1.51	-1.49	3.69	-0.69	-5.11
STD	4.07	12.65	4.51	4.49	4.67	7.31	4.68	4.66
$\varphi = 0.3$								
BIAS	-0.05	10.48	-1.6	-1.74	-0.65	2.51	-0.03	-2.16
STD	4.07	9.03	4.24	4.20	4.31	5.03	4.36	4.29
$\varphi = 0.4$								
BIAS	-0.05	9.00	-0.74	-0.81	-0.33	1.53	0.14	-0.91
STD	4.07	7.08	4.16	4.15	4.22	4.50	4.27	4.23
$\varphi = 0.5$								
BIAS	-0.05	6.75	-0.31	-0.34	-0.18	0.91	0.15	-0.32
STD	4.07	5.90	4.12	4.11	4.14	4.30	4.18	4.17
Panel B: Eigenvector centrality								
$\varphi = 0.2$								
BIAS	-0.01	7.02	-3.94	-4.19	-0.40	2.27	0.39	-0.61
STD	0.61	1.00	0.66	0.67	0.68	0.86	0.67	0.70
$\varphi = 0.3$								
BIAS	-0.01	6.36	-1.93	-2.05	-0.07	1.10	0.54	0.43
STD	0.61	0.84	0.62	0.62	0.64	0.68	0.64	0.66
$\varphi = 0.4$								
BIAS	-0.01	4.83	-0.99	-1.05	-0.01	0.67	0.43	0.60
STD	0.61	0.75	0.62	0.62	0.63	0.64	0.63	0.64
$\varphi = 0.5$								
BIAS	-0.01	3.38	-0.50	-0.54	0.01	0.42	0.30	0.50
STD	0.61	0.67	0.61	0.61	0.61	0.62	0.62	0.62

Note: We conduct Monte Carlo simulations based on 1,000 repetitions of the network formation model (5.1). The network size is $N = 200$. The number of networks is $M = 40$. We set $\beta = (-0.5, -0.5)$ and sampling rates $\varphi \in \{0.2, 0.3, 0.4, 0.5\}$. We report bias and standard deviation $\hat{\alpha}_1$ for the following imputation methods: (1) estimates based on complete network data (**CD**) (2) covariate-only nonparametric imputation that imputes missing outcomes using X alone (**X**), (3) the low-rank imputation methods of Bai and Ng (2021) and Li et al. (2023) (**LR**), (4) the local PCA method that ignores covariates (**LPCA**), (5) the local PCA method using covariates (**X-LPCA**), (6) the local two-way fixed-effects regression that ignores covariates (**LTWFE**), (7) the local two-way fixed-effects regression using covariates (**X-LTWFE**), and (8) X-LTWFE method with sample-splitting (**X-LTWFE-SP**). It is important to note that LR, LPCA, LTWFE, X-LPCA, and X-LTWFE are implemented without sample splitting. Bias and standard deviation are presented in units of 0.01 for ease of reading.

Table 2 displays patterns similar to those in Table 1. As the sampling rate increases, both the bias and the standard deviation of the estimator decline in most cases. This improvement is not only due to more accurate imputation at higher sampling rates (Theorem 2), but also because the aggregate statistics themselves are computed more precisely when fewer observations are missing. The latter explains why the bias and standard deviation in the **X** column also decrease. In most cases, the X-LTWFE estimator delivers the best performance and remains stable even when the sampling rate is small. The only

exception arises when eigenvector centrality is used and the sampling rate is high. In this case, LTWFE yields an extremely small bias—almost identical to that obtained using the complete data. This pattern appears to be driven by features of our numerical design.

It is also worth noting that the bias-std ratio of the linear regression estimator is not close to zero in these simulations. This arises from the use of nonparametric imputation and is particularly evident when eigenvector centrality is employed. Even if researchers choose to rely on a parametric model for imputation, the robustness of our estimator implies that our method remains valuable as a robustness check in applied work.

Table 3: Simulation: Linear-in-means peer-effects model

	CD ($\times 0.01$)	X ($\times 0.01$)	LTWFE ($\times 0.01$)	X-LPCA ($\times 0.01$)	X-LTWFE ($\times 0.01$)	X-LTWFE-SP ($\times 0.01$)
Panel A: Estimates $\alpha_{\bar{Y}}$						
$\varphi = 0.2$						
BIAS	0.70	28.32	16.57	11.59	10.16	13.22
STD	7.50	13.49	13.31	12.63	9.23	11.38
$\varphi = 0.3$						
BIAS	0.70	23.85	11.48	8.50	10.82	11.83
STD	7.50	11.11	9.70	9.53	8.03	9.01
$\varphi = 0.4$						
BIAS	0.70	18.07	6.89	5.66	9.73	9.69
STD	7.50	9.18	8.73	8.40	7.65	8.12
$\varphi = 0.5$						
BIAS	0.70	12.53	3.57	3.18	7.34	7.75
STD	7.50	8.12	8.44	8.07	7.54	7.78
Panel B: Estimates $\alpha_{W,1}$						
$\varphi = 0.2$						
BIAS	0.03	-11.14	18.28	-3.30	3.03	5.66
STD	1.49	1.63	2.11	1.73	1.58	1.79
$\varphi = 0.3$						
BIAS	0.03	9.84	11.57	-1.84	2.04	4.06
STD	1.49	1.52	1.87	1.57	1.51	1.63
$\varphi = 0.4$						
BIAS	0.03	-8.55	7.02	-1.29	1.19	2.68
STD	1.49	1.47	1.73	1.49	1.50	1.56
$\varphi = 0.5$						
BIAS	0.03	-7.13	4.03	-0.98	0.59	1.58
STD	1.49	1.45	1.64	1.48	1.49	1.51

Note: We conduct Monte Carlo simulations based on 1,000 repetitions of the network formation model (5.1). The network size is $N = 200$. The number of networks is $M = 40$. We set $\beta = (-0.5, -0.5)$ and sampling rates $\varphi \in \{0.2, 0.3, 0.4, 0.5\}$. We report bias and standard deviation of $\hat{\alpha}_{\bar{Y}}$ and $\hat{\alpha}_{W,1}$ for the following imputation methods: (1) estimates based on complete network data (**CD**) (2) covariate-only nonparametric imputation that imputes missing outcomes using X alone (**X**), (3) the local two-way fixed-effects regression that ignores covariates (**LTWFE**), (4) the local PCA method using covariates (**X-LPCA**), (5) the local two-way fixed-effects regression using covariates (**X-LTWFE**), and (6) X-LTWFE method with sample-splitting (**X-LTWFE-SP**). It is important to note that LTWFE, X-LPCA, and X-LTWFE are implemented without sample splitting. Bias and standard deviation are presented in units of 0.01 for ease of reading.

Table 3 reports the bias and standard deviation of the estimates $\alpha_{\bar{Y}}$ and $\alpha_{W,1}$ obtained using X, LTWFE, X-PCA, X-LTWFE, and sample-splitting X-LTWFE. The full set of simulation results are presented in Table 6 and Table 7 in Appendix. Similar to those in Table 2, as the sampling rate increases, both the bias and the standard deviation of the estimator decline. In most cases, the estimator based on our proposed method (X-LTWFE) performs very well, especially when the sampling rate is small. We also observe that X-LPCA delivers strong performance: X-LPCA attains a smaller bias in

estimating the endogenous effect α_Y than our proposed estimator when the sampling rate exceeds 30%. In addition, when LTWFE is applied without incorporating covariate information (see the LTWFE column), it delivers a smaller bias in estimating the endogenous effect than X-LTWFE. However, its performance deteriorates substantially when the sampling rate becomes small ($\varphi \leq 30\%$).

The numerical patterns align with the implications of Theorem 4: the bias-to-std ratio of the estimators remains bounded away from zero. This phenomenon is inevitable when the graphon is a non-parametric function of latent factors and poses challenges for inference. Nevertheless, the robustness properties of our estimator suggest that it is valuable as a robustness check in empirical applications.

6 Empirical Application

This section presents an empirical application using the network data of 43 villages in rural India from Banerjee et al. (2013). For each village, the researchers informed certain households (injection nodes or leaders) about the information of microfinance program and asked them to encourage other villagers to join. The researchers want study how households' microfinance take-up decisions are influenced by network characteristics. The data consist of egocentrically sampled social network information and demographic variables (for both sampled and unsampled households). There are $M = 43$ villages, with an average of approximately $N \approx 200$ households per village, and a sampling rate of about $\varphi = 45\%$.

We need to impute the missing links in the adjacency matrices due to egocentric sampling. Since the dataset includes rich demographic information with $d_X > 10$ and the number of households within each village is relatively small, fully nonparametric regression becomes infeasible. Thus, we replace the first-step nonparametric regression with a linear projection on X (another option is to select a subset of important variables and then conduct a nonparametric regression, but this would rely heavily on researchers' discretion). We then apply the local two-way fixed-effects regression to the residuals from this projection.

We focus on the *household-level* regressions in Banerjee et al. (2013). The outcome variable is each household's microfinance take-up decision ($y_i = 1$ if the household decides to take part in the program). We consider three specifications: (i) how individuals' normalized average degrees (defined as in Example 3) affect their take-up decisions; (ii) how individuals' eigenvector centralities (defined as in Example 3) affect their take-up decisions; and (iii) how peers' decisions affect individual take-up through a linear-in-means peer-effects model. Here, W is a dummy variable, indicating whether a household contains injection nodes (See Example 4).

Table 4: Empirical Application: Linear regression on Centrality

	Panel A: Degree centrality		Panel B: Eigenvector centrality	
	Sampled	Imputed	Sampled	Imputed
α_C	0.16	0.13	0.18	0.18
α_1	0.58	0.83	-0.015	-0.011

Note: This table provides estimates of regressions of households' microfinance take-up decisions on households' network statistics, using data from Banerjee et al. (2013). We report results for both sampled (incomplete) networks and imputed networks using our proposed method. The estimates using degree centralities are reported in Panel A. and the estimates using eigenvector centralities are reported in Panel B. α_C represents the intercept and α_1 is the coefficient before the centrality measures.

Table 4 reports estimates of household-level regressions in which the dependent variable is a household's microfinance take-up decision and the regressors are household-level network characteristics. We

provide regression results for both sampled (incomplete) networks and imputed network using our proposed method. Panel A shows that, when using the imputed network, a household’s take-up probability increases by approximately 8.3% when its normalized average degree increases by 10% (equivalently, when the household has roughly 20 more network connections within the village). The estimated effect is weaker when using sampled networks, with the coefficient α_1 decreasing from 0.83 to 0.58. However, as shown in Panel B, we do not find a positive correlation between a household’s take-up probability and its eigenvector centrality, either when using sampled networks or when using imputed networks.

Table 5: Empirical Application: Linear-in-means Peer-effects Model

	α_C	$\alpha_{\bar{Y}}$	α_X	$\alpha_{\bar{X}}$
Sampled	-0.026	0.187	-0.003	0.086
Imputed	-1.376	0.450	0.001	0.106

Note: This table provides estimates of linear-in-means peer-effects model (defined as in Example 4) using data from Banerjee et al. (2013). We report results for both sampled (incomplete) networks and imputed networks using our proposed method. Here, W is a dummy variable, indicating whether a household contains injection nodes. The column α_C reports the estimates of intercept, $\alpha_{\bar{Y}}$ reports the estimates of endogenous effects, and α_X and $\alpha_{\bar{X}}$ report the coefficients on W_i and $(GW)_i$, respectively.

Table 5 reports estimates of household-level linear-in-means peer-effects model in which the dependent variable is a household’s microfinance take-up decision and the regressor W is a dummy variable indicating whether a household contains injection nodes. We provide regression results for both sampled (incomplete) networks and imputed network using our proposed method. $\alpha_{\bar{Y}}$ represents the endogenous peer effect, capturing the influence of neighbors’ decisions to participate in microfinance. Using the imputed network, the estimate of $\alpha_{\bar{Y}}$ is 0.45, implying that a household’s take-up probability increases by 4.5% when the fraction of its friends who join the program increases by 10% (corresponding to roughly two additional friends joining the program, on average). The estimated endogenous effect is weaker when using sampled networks, with the coefficient decreasing from 0.45 to 0.187. In addition, Table 5 shows a negative direct effect ($\alpha_X = -0.003$) when using sampled networks, whereas the direct effect is positive ($\alpha_X = 0.001$) when using imputed networks. For the contextual peer effect, both sampled and imputed networks yield positive estimates, with the estimate based on the imputed network being larger.

7 Conclusion

Sampled network data are common in empirical research because collecting full network information is costly, but using sampled networks can lead to biased estimates. We propose a flexible imputation method for sampled networks and show that downstream empirical analysis using the imputed network yields consistent parameter estimates.

Our method separately imputes the part of each missing link that is explained by covariates and the remaining variation not captured by them: the former via projection onto observed covariates, and the latter using local two-way fixed effects. The imputation method avoids parametric assumptions, does not rely on low-rank restrictions, and flexibly incorporates observed covariates and unobserved heterogeneity. We establish entrywise convergence rates for the imputed matrix and prove consistency of GMM estimators based on the imputed network. We further derive the convergence rate of the corresponding estimator in the linear-in-mean peer effects model. Simulations show strong performance of our method both in terms of imputation accuracy and in downstream empirical analysis.

The robustness of our imputation to misspecification of network formation models comes at the cost of making downstream inference more challenging. In the paper, we discuss how undersmoothing can

help improve the downstream estimator, and other approaches, such as bias correction (Calonico, Cattaneo and Farrell, 2018) and bias-aware inference (Armstrong and Kolesár, 2020), are also potentially applicable. Developing inference procedures for downstream estimators that remain robust to misspecification in network formation is an important topic for future research. In addition, our method is only applicable to dyadic network formation. Extending current imputation methods to strategically formed networks (for example, Mele (2017)) would also be valuable.

References

- Aldous, David J. 1981. “Representations for partially exchangeable arrays of random variables.” *Journal of Multivariate Analysis*, 11(4): 581–598.
- Arcones, Miguel A. 1995. “A Bernstein-type inequality for U-statistics and U-processes.” *Statistics & probability letters*, 22(3): 239–247.
- Armstrong, Timothy B, and Michal Kolesár. 2020. “Simple and honest confidence intervals in nonparametric regression.” *Quantitative Economics*, 11(1): 1–39.
- Auerbach, Eric. 2022. “Identification and estimation of a partially linear regression model using network data.” *Econometrica*, 90(1): 347–365.
- Bai, Jushan, and Serena Ng. 2021. “Matrix completion, counterfactuals, and factor analysis of missing data.” *Journal of the American Statistical Association*, 116(536): 1746–1763.
- Bandeira, Afonso S, and Ramon Van Handel. 2016. “Sharp nonasymptotic bounds on the norm of random matrices with independent entries.”
- Banerjee, Abhijit, Arun G Chandrasekhar, Esther Duflo, and Matthew O Jackson. 2013. “The diffusion of microfinance.” *Science*, 341(6144): 1236–1248.
- Beaman, Lori, and Andrew Dillon. 2018. “Diffusion of agricultural information within social networks: Evidence on gender inequalities from Mali.” *Journal of Development Economics*, 133: 147–161.
- Beyhum, Jad, and Martin Mugnier. 2024. “Inference after discretizing time-varying unobserved heterogeneity.” *arXiv preprint arXiv:2412.07352*.
- Bonhomme, Stéphane, Thibaut Lamadon, and Elena Manresa. 2022. “Discretizing unobserved heterogeneity.” *Econometrica*, 90(2): 625–643.
- Boucher, Vincent, and Aristide Houndetoungan. 2025. “Estimating peer effects using partial network data.” *Review of Economics and Statistics*, 1–48.
- Bramoullé, Yann, Habiba Djebbari, and Bernard Fortin. 2009. “Identification of peer effects through social networks.” *Journal of econometrics*, 150(1): 41–55.
- Bramoullé, Yann, Habiba Djebbari, and Bernard Fortin. 2020. “Peer effects in networks: A survey.” *Annual Review of Economics*, 12(1): 603–629.
- Breza, Emily, Arun G Chandrasekhar, Tyler H McCormick, and Mengjie Pan. 2020. “Using aggregated relational data to feasibly identify network structure without network data.” *American Economic Review*, 110(8): 2454–2484.
- Cai, Jing, Alain De Janvry, and Elisabeth Sadoulet. 2015. “Social networks and the decision to insure.” *American Economic Journal: Applied Economics*, 7(2): 81–108.

- Cai, Tianxi, T Tony Cai, and Anru Zhang.** 2016. “Structured matrix completion with applications to genomic data integration.” *Journal of the American Statistical Association*, 111(514): 621–633.
- Cai, Yong.** 2022. “Linear regression with centrality measures.” *arXiv preprint arXiv:2210.10024*.
- Calonico, Sebastian, Matias D Cattaneo, and Max H Farrell.** 2018. “On the effect of bias estimation on coverage accuracy in nonparametric inference.” *Journal of the American Statistical Association*, 113(522): 767–779.
- Chandrasekhar, Arun, and Randall Lewis.** 2011. “Econometrics of sampled networks.” *Unpublished manuscript, MIT.[422]*.
- Chatterjee, Sourav.** 2015. “Matrix estimation by universal singular value thresholding.” *The Annals of Statistics*.
- Chen, Mingli, Iván Fernández-Val, and Martin Weidner.** 2021. “Nonlinear factor models for network and panel data.” *Journal of Econometrics*, 220(2): 296–324.
- Conley, Timothy G, and Christopher R Udry.** 2010. “Learning about a new technology: Pineapple in Ghana.” *American economic review*, 100(1): 35–69.
- Cruz, Cesi, Julien Labonne, and Pablo Querubin.** 2017. “Politician family networks and electoral outcomes: Evidence from the Philippines.” *American Economic Review*, 107(10): 3006–3037.
- Deaner, Ben, Chen-Wei Hsiang, and Andrei Zeleneev.** 2025. “Inferring Treatment Effects in Large Panels by Uncovering Latent Similarities.” *arXiv preprint arXiv:2503.20769*.
- de Paula, Aureo.** 2020. “Econometric models of network formation.” *Annual Review of Economics*, 12(1): 775–799.
- de Paula, Aureo, Imran Rasul, and Pedro CL Souza.** 2025. “Identifying network ties from panel data: Theory and an application to tax competition.” *Review of Economic Studies*, 92(4): 2691–2729.
- Dzanski, Andreas.** 2019. “An empirical model of dyadic link formation in a network with unobserved heterogeneity.” *Review of Economics and Statistics*, 101(5): 763–776.
- Feng, Yingjie.** 2020. “Causal inference in possibly nonlinear factor models.” *arXiv preprint arXiv:2008.13651*.
- Feng, Yingjie.** 2023. “Optimal estimation of large-dimensional nonlinear factor models.” *arXiv preprint arXiv:2311.07243*.
- Freeman, Hugo, and Martin Weidner.** 2023. “Linear panel regressions with two-way unobserved heterogeneity.” *Journal of Econometrics*, 237(1): 105498.
- Gao, Chao, Yu Lu, and Harrison H Zhou.** 2015. “Rate-optimal graphon estimation.”
- Graham, Bryan S.** 2017. “An econometric model of network formation with degree heterogeneity.” *Econometrica*, 85(4): 1033–1063.
- Graham, Bryan S, Fengshi Niu, and James L Powell.** 2021. “Minimax risk and uniform convergence rates for nonparametric dyadic regression.” National Bureau of Economic Research.
- Griffith, Alan.** 2022. “Name your friends, but only five? the importance of censoring in peer effects estimates using social network data.” *Journal of Labor Economics*, 40(4): 779–805.

- Hardy, Morgan, Rachel M Heath, Wesley Lee, and Tyler H McCormick.** 2024. “Estimating spillovers using imprecisely measured networks.” *arXiv preprint arXiv:1904.00136*.
- Hayes, Alex, and Keith Levin.** 2024. “Peer effects in the linear-in-means model may be inestimable even when identified.” *arXiv preprint arXiv:2410.10772*.
- Herstad, E.** 2023. “Estimating peer effects and network formation models with missing network links.” Working Paper.
- Hochberg, Yael V, Alexander Ljungqvist, and Yang Lu.** 2007. “Whom you know matters: Venture capital networks and investment performance.” *The journal of finance*, 62(1): 251–301.
- Hoover, Douglas N.** 1979. “Relations on Probability Spaces and Arrays of Random Variables.” *Preprint, Institute for Advanced Study*.
- Hsieh, Chih-Sheng, Yu-Chin Hsu, Stanley IM Ko, Jaromír Kovářík, and Trevon D Logan.** 2024. “Non-representative sampled networks: Estimation of network structural properties by weighting.” *Journal of Econometrics*, 240(1): 105689.
- Kitamura, Yuichi, and Louise Laage.** 2024. “Estimating stochastic block models in the presence of covariates.” *arXiv preprint arXiv:2402.16322*.
- Lewbel, Arthur, Xi Qu, and Xun Tang.** 2024. “Ignoring measurement errors in social networks.” *The Econometrics Journal*, 27(2): 171–187.
- Lewbel, Arthur, Xi Qu, and Xun Tang.** 2025. “Estimating Social Network Models with Link Misclassification.” *arXiv preprint arXiv:2509.07343*.
- Li, Tianxi, Yun-Jhong Wu, Elizaveta Levina, and Ji Zhu.** 2023. “Link prediction for egocentrically sampled networks.” *Journal of Computational and Graphical Statistics*, 1–24.
- Manski, Charles F.** 1993. “Identification of endogenous social effects: The reflection problem.” *The review of economic studies*, 60(3): 531–542.
- Ma, Shujie, Liangjun Su, and Yichong Zhang.** 2022. “Detecting latent communities in network formation models.” *The Journal of Machine Learning Research*, 23(1): 13971–14031.
- McCormick, Tyler H, and Tian Zheng.** 2015. “Latent surface models for networks using aggregated relational data.” *Journal of the American Statistical Association*, 110(512): 1684–1695.
- Mele, Angelo.** 2017. “A structural model of dense network formation.” *Econometrica*, 85(3): 825–850.
- Mugnier, Martin.** 2025. “A simple and computationally trivial estimator for grouped fixed effects models.” *Journal of Econometrics*, 250: 106011.
- Newey, Whitney K, and Daniel McFadden.** 1994. “Large sample estimation and hypothesis testing.” *Handbook of econometrics*, 4: 2111–2245.
- Sacerdote, Bruce.** 2001. “Peer effects with random assignment: Results for Dartmouth roommates.” *The Quarterly journal of economics*, 116(2): 681–704.
- Thirkettle, Matthew.** 2019. “Identification and estimation of network statistics with missing link data.” *Unpublished manuscript*.
- Tsybakov, Alexandre B.** 2008. *Introduction to Nonparametric Estimation*. Springer.

- Wainwright, Martin J.** 2019. *High-dimensional statistics: A non-asymptotic viewpoint*. Vol. 48, Cambridge university press.
- Xu, Jiaming.** 2018. “Rates of convergence of spectral methods for graphon estimation.” 5433–5442, PMLR.
- Yu, Yi, Tengyao Wang, and Richard J Samworth.** 2015. “A useful variant of the Davis–Kahan theorem for statisticians.” *Biometrika*, 102(2): 315–323.
- Zeleneev, Andrei.** 2020. “Identification and estimation of network models with nonparametric unobserved heterogeneity.” *Department of Economics, Princeton University*.
- Zhang, Yuan, Elizaveta Levina, and Ji Zhu.** 2017. “Estimating network edge probabilities by neighbourhood smoothing.” *Biometrika*, 104(4): 771–783.

A Implementation and Additional simulation results

A.1 Dyadic nonparametric regression

The extraction of the component of link formation that can be explained by covariates X builds on [Graham, Niu and Powell \(2021\)](#), who studies the asymptotic properties of dyadic nonparametric regression and shows that the Nadaraya-Watson estimator with higher-order kernels achieves the minimax rate. However, our implementation differs from their work in two important respects.

First, instead of using the Nadaraya-Watson estimator, we employ local linear regression. Local linear regression is more commonly used in the local smoothing literature because it delivers a second-order approximation error without requiring higher-order kernel functions. While one could, in principle, use higher-order local polynomial regression to obtain higher-order approximation errors, this is not desirable in our setting for two reasons: (i) sample sizes in applied dyadic settings are typically small; and (ii) the estimation error of the dyadic nonparametric regression step is asymptotically negligible relative to that of the local two-way fixed-effects regression (see Assumption 3 and Theorem 2). We therefore restrict attention to local linear regression in our implementation.

Second, we use $\omega(X_i, X_j)$ (defined as in Example 2) as the regressor rather than (X_i, X_j) . This choice captures homophily in observables in a parsimonious manner and is computationally more efficient. Of course, alternative specifications are possible, and researchers may adopt different formulations depending on the application.

A.2 Algorithm

In the main text, we present a simplified version of our procedure, without sample splitting or cross-validation, in Algorithm 1 to facilitate understanding of the main idea behind our approach. In this section, we introduce Algorithm 2 and 3 for cross-validation and sample-splitting. These two algorithms are employed in the numerical experiments (see Section 5), and we recommend their use in practice.

Algorithm 2 incorporates *leave-one-out* cross-validation for selecting the tuning parameter h but does not use sample splitting. This algorithm is recommended for empirical applications because, similar to other nonparametric procedures, our method relies critically on the choice of the tuning parameter. However, since Algorithm 2 does not employ sample splitting, developing a theoretical analysis based on it is challenging.

Algorithm 3 incorporates both cross-validation and sample splitting and serves as the basis of our theoretical analysis. We randomly split sampled individuals $\mathcal{S}$ into two subsets, $\mathcal{S}_1$ and $\mathcal{S}_2$, of equal sizes. We construct the pseudo-distance using only the network information from $\mathcal{S}_1$. Specifically, for any $i \in \mathcal{S}_2 \cup \mathcal{S}^c$ and $i' \in \mathcal{S}_2$,

$$\hat{d}_{ii'} = \max_{k \in \{1, \dots, N\} \setminus \{i, i'\}} \left| \frac{1}{|\mathcal{S}_1|} \sum_{\ell \in \mathcal{S}_1} A_{k\ell} (A_{i\ell} - A_{i'\ell}) \right| \quad (\text{A.1})$$

We then implement two-way fixed-effects imputation using only the network information from $\mathcal{S}_2$ (excluding links with individuals in $\mathcal{S}_1$). That is, for any $i, j \in \mathcal{S}^c$,

$$\begin{aligned} (\hat{a}, \hat{b}) \in \operatorname{argmin}_{\mathbf{a}, \mathbf{b} \in \mathbb{R}^{|\mathcal{S}_2|+1}} \sum_{\substack{i' \in \mathcal{S}_2 \cup \{i\}, j' \in \mathcal{S}_2 \cup \{j\}, \\ (i', j') \neq (i, j)}} \left(A_{i'j'} - \hat{\Pi}_{i'j'} - a_{i'} - b_{j'} \right)^2 K_h(\hat{d}_{ii'}) K_h(\hat{d}_{jj'}) \\ \hat{A}_{ij} = \hat{a}_i + \hat{b}_j + \hat{\Pi}_{ij} \end{aligned} \quad (\text{A.2})$$

However, since sample splitting reduces the effective sample size, and because network sizes are

relatively small in most economic applications, Algorithm 3's finite-sample performance can be worse than Algorithm 2 in finite samples (see Section 5). The details of Algorithm 2 and Algorithm 3 are provided below.

Algorithm 2 Imputation with Cross Validation

Input: Kernel $K(\cdot)$, a list of bandwidths $\{h_q\}_{q=1,\dots,Q}$, incomplete network A^{obs} , covariate matrix X .

Output: Imputed network $\hat{A}$.

Step 1: Extract information explained by X

Run dyadic nonparametric regression to obtain $\hat{\Pi}(\cdot, \cdot)$.

Step 2: Calculate similarity

For each $i \in 1, \dots, N$ and $i' \in \mathcal{S}$, calculate the pseudo distance using observed adjacency matrix:

$$\hat{d}_{ii'} = \max_{k \in \{1, \dots, N\} \setminus \{i, i'\}} \left| \frac{1}{n} \sum_{\ell=1}^n A_{k\ell} (A_{i\ell} - A_{i'\ell}) \right|$$

Step 3: Leave-one-out cross validation

For each $h_q \in \{h_1, \dots, h_Q\}$, and for each $i \in \mathcal{S}$ and $j \in \mathcal{S}^c$, estimate the local (weighted) two-way fixed-effects model:

$$(\hat{\mathbf{a}}(h_q), \hat{\mathbf{b}}(h_q)) \in \underset{\mathbf{a}, \mathbf{b} \in \mathbb{R}^n}{\operatorname{argmin}} \sum_{i' \in \mathcal{S}, j' \in \mathcal{S} \cup \{j\} \setminus \{i\}} \left(A_{i'j'} - \hat{\Pi}_{i'j'} - a_{i'} - b_{j'} \right)^2 K_{h_q}(\hat{d}_{ii'}) K_{h_q}(\hat{d}_{jj'})$$

to obtain $\bar{A}_{ij}(h_q) = \hat{a}_i(h_q) + \hat{b}_j(h_q) + \hat{\Pi}_{ij}$. Then the optimal h^* solves

$$h^* = \underset{h_q \in \{h_1, \dots, h_Q\}}{\operatorname{argmin}} \sum_{i \in \mathcal{S}, j \in \mathcal{S}^c} \left(A_{ij} - \bar{A}_{ij}(h_q) \right)^2$$

Step 4: Local imputation

For each $i \in \mathcal{S}^c$ and $j \in \mathcal{S}^c$, estimate the local (weighted) two-way fixed-effects model:

$$(\hat{\mathbf{a}}, \hat{\mathbf{b}}) \in \underset{\mathbf{a}, \mathbf{b} \in \mathbb{R}^{n+1}}{\operatorname{argmin}} \sum_{\substack{i' \in \mathcal{S} \cup \{i\}, j' \in \mathcal{S} \cup \{j\}, \\ (i', j') \neq (i, j)}} \left(A_{i'j'} - \hat{\Pi}_{i'j'} - a_{i'} - b_{j'} \right)^2 K_{h^*}(\hat{d}_{ii'}) K_{h^*}(\hat{d}_{jj'})$$

to obtain $\tilde{A}_{ij} = \hat{a}_i + \hat{b}_j + \hat{\Pi}_{ij}$. By combining the imputed entries with the observed entries of the adjacency matrix, we obtain the (preliminary) imputed adjacency matrix $\tilde{A}$. Truncate $\tilde{A}_{ij}$ to get

$$\hat{A}_{ij} = \begin{cases} \tilde{A}_{ij}, & \text{if } \tilde{A}_{ij} \in [0, 1] \\ 1 & \text{if } \tilde{A}_{ij} > 1 \\ 0 & \text{if } \tilde{A}_{ij} < 0 \end{cases}$$

and zero the diagonal of $\hat{A}$ such that $\hat{A}_{ii} = 0$ for all i .

Algorithm 3 Imputation with Cross Validation and Sample Splitting

Input: Kernel $K(\cdot)$, a list of bandwidths $\{h_q\}_{q=1,\dots,Q}$, incomplete network A^{obs} , covariate matrix X .

Randomly split $\mathcal{S}$ into two parts, $\mathcal{S}_1, \mathcal{S}_2$, with equal sizes.

Output: Imputed network $\hat{A}$.

Step 1: Extract information explained by X

Run dyadic nonparametric regression to obtain $\hat{\Pi}(\cdot, \cdot)$.

Step 2: Calculate similarity

For each $i \in \mathcal{S}_2 \cup \mathcal{S}^c$, and $i' \in \mathcal{S}_2$, calculate the pseudo distance using observed adjacency matrix:

$$\hat{d}_{ii'} = \max_{k \in \{1, \dots, N\} \setminus \{i, i'\}} \left| \frac{1}{|\mathcal{S}_1|} \sum_{\ell \in \mathcal{S}_1} A_{k\ell} (A_{i\ell} - A_{i'\ell}) \right|$$

Step 3: Leave-one-out cross validation

For each $h_q \in \{h_1, \dots, h_Q\}$, and for each $i \in \mathcal{S}_2$ and $j \in \mathcal{S}^c$, run the local (weighted) two-way fixed effect regression

$$(\hat{\mathbf{a}}(h_q), \hat{\mathbf{b}}(h_q)) \in \underset{\mathbf{a}, \mathbf{b} \in \mathbb{R}^{|\mathcal{S}_2|}}{\text{argmin}} \sum_{i' \in \mathcal{S}_2, j' \in \mathcal{S}_2 \cup \{j\} \setminus \{i\}} \left(A_{i'j'} - \hat{\Pi}_{i'j'} - a_{i'} - b_{j'} \right)^2 K_{h_q}(\hat{d}_{ii'}) K_{h_q}(\hat{d}_{jj'})$$

to obtain $\bar{A}_{ij}(h_q) = \hat{\mathbf{a}}_i(h_q) + \hat{\mathbf{b}}_j(h_q) + \hat{\Pi}_{X,ij}$. Then the optimal h^* solves

$$h^* = \underset{h_q \in \{h_1, \dots, h_Q\}}{\text{argmin}} \sum_{i \in \mathcal{S}_2, j \in \mathcal{S}^c} (A_{ij} - \bar{A}_{ij}(h_q))^2$$

Step 4: Local imputation

For each $i \in \mathcal{S}^c$ and $j \in \mathcal{S}^c$, estimate the local (weighted) two-way fixed-effects regression

$$(\hat{\mathbf{a}}, \hat{\mathbf{b}}) \in \underset{\mathbf{a}, \mathbf{b} \in \mathbb{R}^{|\mathcal{S}_2|+1}}{\text{argmin}} \sum_{\substack{i' \in \mathcal{S}_2 \cup \{i\}, j' \in \mathcal{S}_2 \cup \{j\}, \\ (i', j') \neq (i, j)}} \left(A_{i'j'} - \hat{\Pi}_{i'j'} - a_{i'} - b_{j'} \right)^2 K_{h^*}(\hat{d}_{ii'}) K_{h^*}(\hat{d}_{jj'})$$

to obtain $\tilde{A}_{ij} = \hat{\mathbf{a}}_i + \hat{\mathbf{b}}_j + \hat{\Pi}_{ij}$. By combining the imputed entries with the observed entries of the adjacency matrix, we obtain the (preliminary) imputed adjacency matrix $\tilde{A}$. Truncate $\tilde{A}_{ij}$ to get

$$\hat{A}_{ij} = \begin{cases} \tilde{A}_{ij}, & \text{if } \tilde{A}_{ij} \in [0, 1] \\ 1 & \text{if } \tilde{A}_{ij} > 1 \\ 0 & \text{if } \tilde{A}_{ij} < 0 \end{cases}$$

and zero the diagonal of $\hat{A}$ such that $\hat{A}_{ii} = 0$ for all i .

Table 6: Simulation: Linear-in-means peer-effects model (Supplementary results I)

	CD ($\times 0.01$)	X ($\times 0.01$)	LTWFE ($\times 0.01$)	X-LPCA ($\times 0.01$)	X-LTWFE ($\times 0.01$)	X-LTWFE-SP ($\times 0.01$)
Panel A: Estimates α_C						
$\varphi = 0.2$						
BIAS	-1.02	-2.45	-22.81	20.32	-13.24	-10.64
STD	14.39	24.71	23.96	23.69	17.81	22.13
$\varphi = 0.3$						
BIAS	-1.02	1.87	-22.64	13.21	-15.8	-16.09
STD	14.39	20.46	18.12	18.11	15.60	17.69
$\varphi = 0.4$						
BIAS	-1.02	7.24	-17.02	8.38	-14.66	-16.43
STD	14.39	17.16	16.61	16.12	14.81	15.85
$\varphi = 0.5$						
BIAS	-1.02	10.64	-11.16	5.80	-12.11	-14.16
STD	14.39	15.40	16.21	15.56	14.56	15.11
Panel A: Estimates $\alpha_{W,2}$						
$\varphi = 0.2$						
BIAS	0.10	13.48	43.41	19.51	19.01	20.39
STD	1.57	1.96	2.24	1.87	1.77	1.92
$\varphi = 0.3$						
BIAS	0.10	10.8	30.29	15.41	14.23	14.85
STD	1.57	1.87	2.03	1.81	1.73	1.77
$\varphi = 0.4$						
BIAS	0.10	8.24	20.3	11.4	10.23	10.47
STD	1.57	1.79	1.90	1.75	1.69	1.70
$\varphi = 0.5$						
BIAS	0.10	5.91	12.78	7.74	6.88	6.97
STD	1.57	1.73	1.77	1.69	1.66	1.65

Note: We conduct Monte Carlo simulations based on 1,000 repetitions of the network formation model (5.1). The network size is $N = 200$. The number of networks is $M = 40$. We set $\beta = (-0.5, -0.5)$ and sampling rates $\varphi \in \{0.2, 0.3, 0.4, 0.5\}$. We report bias and standard deviation of $\hat{\alpha}_C$ and $\hat{\alpha}_{W,2}$ for the following imputation methods: (1) estimates based on complete network data (**CD**) (2) covariate-only nonparametric imputation that imputes missing outcomes using X alone (**X**), (3) the local two-way fixed-effects regression that ignores covariates (**LTWFE**), (4) the local PCA method using covariates (**X-LPCA**), (5) the local two-way fixed-effects regression using covariates (**X-LTWFE**), and (6) X-LTWFE method with sample-splitting (**X-LTWFE-SP**). It is important to note that LTWFE, X-LPCA, and X-LTWFE are implemented without sample splitting. Bias and standard deviation are presented in units of 0.01 for ease of reading.

Table 7: Simulation: Linear-in-means peer-effects model (Supplementary results II)

	CD ($\times 0.01$)	X ($\times 0.01$)	LTWFE ($\times 0.01$)	X-LPCA ($\times 0.01$)	X-LTWFE ($\times 0.01$)	X-LTWFE-SP ($\times 0.01$)
Panel A: Estimates $\alpha_{\bar{W},1}$						
$\varphi = 0.2$						
BIAS	-1.54	-101.43	-66.82	-84.64	-47.86	-47.89
STD	14.29	22.62	24.97	20.31	17.02	20.28
$\varphi = 0.3$						
BIAS	-1.54	-92.36	-34.83	-58.39	-35.21	-36.9
STD	14.29	18.70	18.08	16.21	15.23	16.69
$\varphi = 0.4$						
BIAS	-1.54	-77.54	-16.13	-37.15	-25.24	-27.06
STD	14.29	15.44	16.26	14.89	14.67	15.39
$\varphi = 0.5$						
BIAS	-1.54	-50.38	-5.88	-21.92	-17.05	-18.76
STD	14.29	13.79	15.59	14.66	14.47	14.86
Panel A: Estimates $\alpha_{\bar{W},2}$						
$\varphi = 0.2$						
BIAS	-1.69	-19.57	-50.39	13.96	-13.15	3.66
STD	16.83	35.25	33.62	34.3	24.16	29.19
$\varphi = 0.3$						
BIAS	-1.69	-12.86	-32.09	3.75	-16.02	-5.44
STD	16.83	28.67	23.96	25.42	20.40	22.46
$\varphi = 0.4$						
BIAS	-1.69	-5.04	-18.38	-0.28	-14.64	-8.44
STD	16.83	23.61	20.78	21.52	18.75	19.68
$\varphi = 0.5$						
BIAS	-1.69	0.78	-8.47	-0.07	-11.43	-8.04
STD	16.83	20.63	19.78	19.95	18.09	18.54

Note: We conduct Monte Carlo simulations based on 1,000 repetitions of the network formation model (5.1). The network size is $N = 200$. The number of networks is $M = 40$. We set $\beta = (-0.5, -0.5)$ and sampling rates $\varphi \in \{0.2, 0.3, 0.4, 0.5\}$. We report bias and standard deviation of $\hat{\alpha}_{\bar{W},1}$ and $\hat{\alpha}_{\bar{W},2}$ for the following imputation methods: (1) estimates based on complete network data (**CD**) (2) covariate-only nonparametric imputation that imputes missing outcomes using X alone (**X**), (3) the local two-way fixed-effects regression that ignores covariates (**LTWFE**), (4) the local PCA method using covariates (**X-LPCA**), (5) the local two-way fixed-effects regression using covariates (**X-LTWFE**), and (6) X-LTWFE method with sample-splitting (**X-LTWFE-SP**). It is important to note that LTWFE, X-LPCA, and X-LTWFE are implemented without sample splitting. Bias and standard deviation are presented in units of 0.01 for ease of reading.

B Consistency of Imputation

Proof of Lemma 1. In the proof, we aim to establish

$$\mathbb{P} \left(\max_{i \in \mathcal{S}^c, i' \in \mathcal{S}_2} \left| \hat{d}_{ii'} - d_{ii'} \right| \geq \gamma_1 \frac{\log N}{N^{1/d_\zeta}} + \gamma_2 \frac{\log N}{\sqrt{n}} \right) \leq 1 - n^{-\frac{1}{2}}$$

Actually, we can show that for any $\delta > 0$, there exist corresponding constants $\gamma_1, \gamma_2 > 0$, such that

$$\mathbb{P} \left(\max_{i \in \mathcal{S}^c, i' \in \mathcal{S}_2} \left| \hat{d}_{ii'} - d_{ii'} \right| \geq \gamma_1 \frac{\log N}{N^{1/d_\zeta}} + \gamma_2 \frac{\log N}{\sqrt{n}} \right) \leq 1 - \delta n^{-\frac{1}{2}}$$

In the following proof, we use δ to denote a generic positive constant that may vary from line to line and absorb multiplicative or additive constants.

Recall that for any $i, i' \in \{1, \dots, N\}$,

$$\hat{d}_{ii'} = \max_{k \in \{i, i'\}} \left| \frac{1}{|\mathcal{S}_1|} \sum_{\ell \in \mathcal{S}_1} A_{k\ell} (A_{i\ell} - A_{i'\ell}) \right|$$

The population counterpart of $\hat{d}_{ii'}$ is

$$d_{ii'} := \sup_{\zeta \in \text{supp}(\zeta)} \left| \int_{\zeta_\ell} f(\zeta, \zeta_\ell) (f(\zeta_i, \zeta_\ell) - f(\zeta_{i'}, \zeta_\ell)) d\mathbb{P}_\zeta(\zeta_\ell) \right|$$

We define

$$\tilde{d}_{ii'} := \max_{k \notin \{i, i'\}} \left| \frac{1}{|\mathcal{S}_1|} \sum_{\ell \in \mathcal{S}_1 \setminus \{k\}} f(\zeta_k, \zeta_\ell) (f(\zeta_i, \zeta_\ell) - f(\zeta_{i'}, \zeta_\ell)) \right|$$

Then, by the triangle inequality, it follows that

$$\max_{i \in \mathcal{S}^c, i' \in \mathcal{S}_2} \left| \hat{d}_{ii'} - d_{ii'} \right| \leq \max_{i \in \mathcal{S}^c, i' \in \mathcal{S}_2} \left| \hat{d}_{ii'} - \tilde{d}_{ii'} \right| + \max_{i \in \mathcal{S}^c, i' \in \mathcal{S}_2} \left| \tilde{d}_{ii'} - d_{ii'} \right|$$

Bound on $\max_{i \in \mathcal{S}^c, i' \in \mathcal{S}_2} \left| \hat{d}_{ii'} - \tilde{d}_{ii'} \right|$ For any i, i' and any $k \in \{1, \dots, N\} \setminus \{i, i'\}$, we have

$$\begin{aligned} & \left| \left| \frac{1}{|\mathcal{S}_1|} \sum_{\ell \in \mathcal{S}_1} A_{k\ell} (A_{i\ell} - A_{i'\ell}) \right| - \left| \frac{1}{|\mathcal{S}_1|} \sum_{\ell \in \mathcal{S}_1 \setminus \{k\}} f(\zeta_k, \zeta_\ell) (f(\zeta_i, \zeta_\ell) - f(\zeta_{i'}, \zeta_\ell)) \right| \right| \\ & \leq \left| \frac{1}{|\mathcal{S}_1|} \sum_{\ell \in \mathcal{S}_1 \setminus \{k\}} \epsilon_{k\ell} (f(\zeta_i, \zeta_\ell) + f(\zeta_{i'}, \zeta_\ell)) \right| + \left| \frac{1}{|\mathcal{S}_1|} \sum_{\ell \in \mathcal{S}_1 \setminus \{k\}} f(\zeta_k, \zeta_\ell) (\epsilon_{i\ell} - \epsilon_{i'\ell}) \right| + \left| \frac{1}{|\mathcal{S}_1|} \sum_{\ell \in \mathcal{S}_1 \setminus \{k\}} \epsilon_{k\ell} (\epsilon_{i\ell} - \epsilon_{i'\ell}) \right| \end{aligned}$$

By the triangle inequality, it follows that

$$\begin{aligned}
\max_{i \in \mathcal{S}^c, i' \in \mathcal{S}_2} |\hat{d}_{ij} - \tilde{d}_{ij}| &\leq \underbrace{\max_{i \in \mathcal{S}^c, i' \in \mathcal{S}_2} \max_{k \notin \{i, i'\}} \left| \frac{1}{|\mathcal{S}_1|} \sum_{\ell \in \mathcal{S}_1 \setminus \{k\}} \epsilon_{k\ell} (f(\zeta_i, \zeta_\ell) + f(\zeta_{i'}, \zeta_\ell)) \right|}_{B_1} \\
&\quad + \underbrace{\max_{i \in \mathcal{S}^c, i' \in \mathcal{S}_2} \max_{k \notin \{i, i'\}} \left| \frac{1}{|\mathcal{S}_1|} \sum_{\ell \in \mathcal{S}_1 \setminus \{k\}} f(\zeta_k, \zeta_\ell) (\epsilon_{i\ell} - \epsilon_{i'\ell}) \right|}_{B_2} \\
&\quad + \underbrace{\max_{i \in \mathcal{S}^c, i' \in \mathcal{S}_2} \max_{k \notin \{i, i'\}} \left| \frac{1}{|\mathcal{S}_1|} \sum_{\ell \in \mathcal{S}_1 \setminus \{k\}} \epsilon_{k\ell} (\epsilon_{i\ell} - \epsilon_{i'\ell}) \right|}_{B_3}
\end{aligned}$$

Since the support of ζ is compact (Assumption 1(iv)) and the graphon f is smooth (Assumption 1(v)), there exists a constant ρ_f such that $\sup_f \|f\|_\infty \leq \rho_f$. Since the random variables $\{\zeta_i\}_{i=1}^N$ and $\{\epsilon_{ij}\}_{1 \leq i < j \leq N}$ are mutually independent (Assumption 1(ii)), by Bernstein's inequality (Lemma D.3), we obtain that for any $t > 0$,

$$\begin{aligned}
\mathbb{P}(B_1 \geq t) &\leq \mathbb{P} \left(\bigcup_{i \in \mathcal{S}^c, i' \in \mathcal{S}_2} \bigcup_{k \notin \{i, i'\}} \left\{ \left| \frac{1}{|\mathcal{S}_1|} \sum_{\ell \in \mathcal{S}_1 \setminus \{k\}} \epsilon_{k\ell} (f(\zeta_i, \zeta_\ell) + f(\zeta_{i'}, \zeta_\ell)) \right| \geq t \right\} \right) \\
&\leq \sum_{i \in \mathcal{S}^c, i' \in \mathcal{S}_2} \sum_{k \notin \{i, i'\}} \mathbb{P} \left(\left| \frac{1}{|\mathcal{S}_1|} \sum_{\ell \in \mathcal{S}_1 \setminus \{k\}} \epsilon_{k\ell} (f(\zeta_i, \zeta_\ell) + f(\zeta_{i'}, \zeta_\ell)) \right| \geq t \right) \\
&\leq N^3 \mathbb{P} \left(\left| \frac{1}{|\mathcal{S}_1|} \sum_{\ell \in \mathcal{S}_1 \setminus \{k\}} \epsilon_{k\ell} (f(\zeta_i, \zeta_\ell) + f(\zeta_{i'}, \zeta_\ell)) \right| \geq t \right) \\
&\leq 2N^3 \exp \left(-\frac{nt^2}{16\sigma^2\rho_f^2 + 8\sigma\rho_ft} \right)
\end{aligned}$$

In addition, since the random variables $\{\zeta_i\}_{i=1}^N$ and $\{\epsilon_{ij}\}_{1 \leq i < j \leq N}$ are mutually independent (Assumption 1(ii)), by Bernstein's inequality (Lemma D.3), we obtain that for any $t > 0$,

$$\begin{aligned}
\mathbb{P}(B_2 \geq t) &\leq \mathbb{P} \left(\bigcup_{i \in \mathcal{S}^c, i' \in \mathcal{S}_2} \bigcup_{k \notin \{i, i'\}} \left\{ \left| \frac{1}{|\mathcal{S}_1|} \sum_{\ell \in \mathcal{S}_1 \setminus \{k\}} f(\zeta_k, \zeta_\ell) (\epsilon_{i\ell} - \epsilon_{i'\ell}) \right| \geq t \right\} \right) \\
&\leq \sum_{i \in \mathcal{S}^c, i' \in \mathcal{S}_2} \sum_{k \notin \{i, i'\}} \mathbb{P} \left(\left| \frac{1}{|\mathcal{S}_1|} \sum_{\ell \in \mathcal{S}_1 \setminus \{k\}} f(\zeta_k, \zeta_\ell) (\epsilon_{i\ell} - \epsilon_{i'\ell}) \right| \geq t \right) \\
&\leq N^3 \mathbb{P} \left(\left| \frac{1}{|\mathcal{S}_1|} \sum_{\ell \in \mathcal{S}_1 \setminus \{k\}} f(\zeta_k, \zeta_\ell) (\epsilon_{i\ell} - \epsilon_{i'\ell}) \right| \geq t \right) \\
&\leq 2N^3 \exp \left(-\frac{nt^2}{16\sigma^2\rho_f^2 + 8\sigma\rho_ft} \right)
\end{aligned}$$

Based on the similar argument, we have that for any $t > 0$,

$$\begin{aligned}
\mathbb{P}(B_3 \geq t) &\leq \mathbb{P}\left(\bigcup_{i \in \mathcal{S}^c, i' \in \mathcal{S}_2} \bigcup_{k \notin \{i, i'\}} \left\{ \left| \frac{1}{|\mathcal{S}_1|} \sum_{\ell \in \mathcal{S}_1 \setminus \{k\}} \epsilon_{k\ell} (\epsilon_{i\ell} - \epsilon_{i'\ell}) \right| \geq t \right\}\right) \\
&\leq \sum_{i \in \mathcal{S}^c, i' \in \mathcal{S}_2} \sum_{k \notin \{i, i'\}} \mathbb{P}\left(\left| \frac{1}{|\mathcal{S}_1|} \sum_{\ell \in \mathcal{S}_1 \setminus \{k\}} \epsilon_{k\ell} (\epsilon_{i\ell} - \epsilon_{i'\ell}) \right| \geq t\right) \\
&\leq N^3 \mathbb{P}\left(\left| \frac{1}{|\mathcal{S}_1|} \sum_{\ell \in \mathcal{S}_1 \setminus \{k\}} \epsilon_{k\ell} (\epsilon_{i\ell} - \epsilon_{i'\ell}) \right| \geq t\right) \\
&\leq 2N^3 \exp\left(-\frac{nt^2}{16\sigma^4 + 8\sigma^2 t}\right)
\end{aligned}$$

Combining the probability bounds for B_1, B_2, B_3 , we conclude that there exists a sufficiently large constant $C_1 > 0$ such that when $t = C_1 \frac{\log N}{\sqrt{n}}$,

$$\mathbb{P}\left(\max_{i \in \mathcal{S}^c, i' \in \mathcal{S}_2} |\hat{d}_{ii'} - \tilde{d}_{ii'}| \geq C_1 \frac{\log N}{\sqrt{n}}\right) \leq \delta n^{-1/2}$$

Bound on $\max_{i \in \mathcal{S}^c, i' \in \mathcal{S}_2} |\tilde{d}_{ii'} - d_{ii'}|$ For any $i \in \mathcal{S}^c$ and $i' \in \mathcal{S}_2$,

$$\begin{aligned}
|\tilde{d}_{ij} - d_{ij}| &\leq \max_{k \notin \{i, i'\}} \left| \frac{1}{|\mathcal{S}_1|} \sum_{\ell \in \mathcal{S}_1 \setminus \{k\}} f(\zeta_k, \zeta_\ell) (f(\zeta_i, \zeta_\ell) - f(\zeta_{i'}, \zeta_\ell)) - \int_{\zeta_\ell} f(\zeta_k, \zeta_\ell) (f(\zeta_i, \zeta_\ell) - f(\zeta_{i'}, \zeta_\ell)) d\mathbb{P}_\zeta(\zeta_\ell) \right| \\
&\quad + \sup_{\zeta \in \text{supp}(\zeta)} \left| \int_{\zeta_\ell} f(\zeta, \zeta_\ell) (f(\zeta_i, \zeta_\ell) - f(\zeta_{i'}, \zeta_\ell)) d\mathbb{P}_\zeta(\zeta_\ell) \right| \\
&\quad - \max_{k \notin \{i, i'\}} \left| \int_{\zeta_\ell} f(\zeta_k, \zeta_\ell) (f(\zeta_i, \zeta_\ell) - f(\zeta_{i'}, \zeta_\ell)) d\mathbb{P}_\zeta(\zeta_\ell) \right|
\end{aligned}$$

It follows that $\max_{i \in \mathcal{S}^c, i' \in \mathcal{S}_2} |\tilde{d}_{ii'} - d_{ii'}|$ is bounded by

$$\begin{aligned}
&\underbrace{\max_{i \in \mathcal{S}^c, i' \in \mathcal{S}_2} \max_{k \notin \{i, i'\}} \left| \frac{1}{|\mathcal{S}_1|} \sum_{\ell \in \mathcal{S}_1 \setminus \{k\}} f(\zeta_k, \zeta_\ell) (f(\zeta_i, \zeta_\ell) - f(\zeta_{i'}, \zeta_\ell)) - \int_{\zeta_\ell} f(\zeta_k, \zeta_\ell) (f(\zeta_i, \zeta_\ell) - f(\zeta_{i'}, \zeta_\ell)) d\mathbb{P}_\zeta(\zeta_\ell) \right| + \frac{4\rho_f^2}{n}}_{Q_1} \\
&+ \underbrace{\max_{i \in \mathcal{S}^c, i' \in \mathcal{S}_2} \left(\sup_{\zeta \in \text{supp}(\zeta)} \left| \int_{\zeta_\ell} f(\zeta, \zeta_\ell) (f(\zeta_i, \zeta_\ell) - f(\zeta_{i'}, \zeta_\ell)) d\mathbb{P}_\zeta(\zeta_\ell) \right| - \max_{k \notin \{i, i'\}} \left| \int_{\zeta_\ell} f(\zeta_k, \zeta_\ell) (f(\zeta_i, \zeta_\ell) - f(\zeta_{i'}, \zeta_\ell)) d\mathbb{P}_\zeta(\zeta_\ell) \right| \right)}_{Q_2}
\end{aligned}$$

Since the random variables $\{\zeta_i\}_{i=1}^N$ are independent (Assumption 1(ii)), by Bernstein's inequality (Lemma D.3), we have that for any $t > 0$,

$$\mathbb{P}(Q_1 \geq t) \leq 2N^3 \exp\left(-\frac{nt^2}{16\rho_f^4 + 8\sigma\rho_f^4 t}\right).$$

Then, there exists a sufficiently large constant $C_2 > 0$ such that when $t = C_2 \frac{\log N}{\sqrt{n}}$, we have

$$\mathbb{P}\left(Q_1 \geq C_2 \frac{\log N}{\sqrt{n}}\right) \leq \delta n^{-1/2}$$

To establish a bound for Q_2 , let ζ^* be a maximizer of $d_{ii'}$ such that

$$d_{ii'} = \left| \int_{\zeta_\ell} f(\zeta^*, \zeta_\ell) (f(\zeta_i, \zeta_\ell) - f(\zeta_{i'}, \zeta_\ell)) d\mathbb{P}_\zeta(\zeta_\ell) \right|.$$

Then, we have:

$$\begin{aligned} Q_2 &\leq \max_{i \in \mathcal{S}^c, i' \in \mathcal{S}_2} \left(\left| \int_{\zeta_\ell} f(\zeta^*, \zeta_\ell) (f(\zeta_i, \zeta_\ell) - f(\zeta_{i'}, \zeta_\ell)) d\mathbb{P}_\zeta(\zeta_\ell) \right| - \min_{k \notin \{i, i'\}} \left| \int_{\zeta_\ell} f(\zeta_k, \zeta_\ell) (f(\zeta_i, \zeta_\ell) - f(\zeta_{i'}, \zeta_\ell)) d\mathbb{P}_\zeta(\zeta_\ell) \right| \right) \\ &\stackrel{(i)}{\leq} \max_{i \in \mathcal{S}^c, i' \in \mathcal{S}_2} \min_{k \notin \{i, i'\}} 2\rho_f L d_\zeta^{1/2} \|\zeta^* - \zeta_k\| \\ &\leq 2\rho_f L d_\zeta^{1/2} \max_{i \in \mathcal{S}^c, i' \in \mathcal{S}_2} \sup_{\zeta} \min_{k \notin \{i, i'\}} \|\zeta - \zeta_k\|. \end{aligned}$$

Here, inequality (i) follows from Assumption 1(v) that f is a Lipschitz function with constant $L > 0$. By Assumption 1(iv), there exists a t -covering of the support of ζ consisting of $\mathcal{J} \leq 4^{d_\zeta} d_\zeta^{d_\zeta/2} t^{-d_\zeta}$ open balls with radius t , denoted as $\{\bar{\zeta}_j\}_{j=1}^{\mathcal{J}}$. Therefore,

$$\sup_{\zeta \in \text{supp}(\zeta)} \min_{k \notin \{i, i'\}} \|\zeta - \zeta_k\| \leq \max_{j=1, \dots, \mathcal{J}} \min_{k \notin \{i, i'\}} \|\bar{\zeta}_j - \zeta_k\| + t.$$

It follows that for any $t \leq (2\underline{c})^{1/d_\zeta}$,

$$\begin{aligned} \mathbb{P} \left(\max_{i \in \mathcal{S}^c, i' \in \mathcal{S}_2} \sup_{\zeta \in \text{supp}(\zeta)} \min_{k \notin \{i, i'\}} \|\zeta - \zeta_k\| \geq 2t \right) &\leq \mathbb{P} \left(\bigcup_{i \in \mathcal{S}^c, i' \in \mathcal{S}_2} \left\{ \sup_{\zeta \in \text{supp}(\zeta)} \min_{k \notin \{i, i'\}} \|\zeta - \zeta_k\| \geq 2t \right\} \right) \\ &\leq \sum_{i \in \mathcal{S}^c, i' \in \mathcal{S}_2} \mathbb{P} \left(\sup_{\zeta \in \text{supp}(\zeta)} \min_{k \notin \{i, i'\}} \|\zeta - \zeta_k\| \geq 2t \right) \\ &\leq N^2 \mathbb{P} \left(\sup_{\zeta \in \text{supp}(\zeta)} \min_{k \notin \{i, i'\}} \|\zeta - \zeta_k\| \geq 2t \right) \\ &\leq N^2 \mathbb{P} \left(\max_{j=1, \dots, \mathcal{J}} \min_{k \notin \{i, i'\}} \|\bar{\zeta}_j - \zeta_k\| \geq t \right) \\ &\leq N^2 \mathcal{J} \mathbb{P} \left(\min_{k \notin \{i, i'\}} \|\bar{\zeta}_j - \zeta_k\| \geq t \right) \\ &\stackrel{(i)}{=} N^2 \mathcal{J} (\mathbb{P}(\|\bar{\zeta}_j - \zeta\| \geq t))^{N-2} \\ &\stackrel{(ii)}{\leq} 4N^2 \mathcal{J} (1 - \underline{c}t^{d_\zeta})^N \\ &\leq 4^{d_\zeta+1} d_\zeta^{d_\zeta/2} N^2 t^{-d_\zeta} (1 - \underline{c}t^{d_\zeta})^N \end{aligned}$$

where equality (i) follows from the independence of $\{\zeta_i \mid i = 1, \dots, N\}$ (Assumption 1(ii)), and inequality (ii) follows from Assumption 1(iv). Therefore, there exists a constant $C_3 > 0$ such that when $t = C_3 \frac{\log N}{N^{1/d_\zeta}}$,

$$\mathbb{P} \left(Q_2 \geq C_3 \frac{\log N}{N^{1/d_\zeta}} \right) \leq \delta n^{-1/2}.$$

Combining the above probability bounds and letting $\gamma_1 = C_3$ as $\gamma_2 = C_1 + C_2$, we obtain

$$\mathbb{P} \left(\max_{i \in \mathcal{S}^c, i' \in \mathcal{S}_2} \left| \hat{d}_{ij} - d_{ij} \right| \geq \gamma_1 \frac{\log N}{N^{1/d_\zeta}} + \gamma_2 \frac{\log N}{\sqrt{n}} + \frac{4\rho_f^2}{n} \right) \leq \delta n^{-1/2}$$

Since δ is a generic positive constant, we set $\delta = 1$ and prove Lemma 1.

Finally, since $\frac{4\rho_f^2}{n}$ is asymptotically negligible relative to $\gamma_2 \frac{\log N}{\sqrt{n}}$, it is sufficient to focus on the event

$$\mathcal{A} := \left\{ \max_{i \in \mathcal{S}^c, i' \in \mathcal{S}_2} |\hat{d}_{ii'} - d_{ii'}| \geq \gamma_1 \frac{\log N}{N^{1/d_\zeta}} + \gamma_2 \frac{\log N}{\sqrt{n}} \right\} = \left\{ \max_{i \in \mathcal{S}^c, i' \in \mathcal{S}_2} |\hat{d}_{ii'} - d_{ii'}| \geq \delta_{N,n} \right\} \quad (\text{B.1})$$

■

Lemma B.1 *Conditional on the event $\mathcal{A}$ defined in (B.1), for any $h > 0$ such that $\delta_{N,n}/h \rightarrow 0$, and for all $i \in \mathcal{S}^c, i' \in \mathcal{S}_2$, if $\hat{d}_{ii'} \leq h$, then there exists a constant $\eta > 0$ such that*

$$\|\zeta_i - \zeta_{i'}\| \leq \eta h$$

Proof of Lemma B.1. Note that for any $i \in \mathcal{S}^c$ and $i' \in \mathcal{S}_2$, since $\delta_{N,n}/h \rightarrow 0$, there exists a constant $C_1 > 1$ such that

$$\{\hat{d}_{ii'} \leq h\} \subseteq \{d_{ii'} \leq h + \delta_{N,n}\} \subseteq \{d_{ii'} \leq C_1 h\}.$$

Also, by Assumption 2, we have

$$\{d_{ii'} \leq C_1 h\} \subseteq \{\|\zeta_i - \zeta_{i'}\| \leq \lambda^{-1} C_1 h\}$$

Let $\eta = \lambda^{-1} C_1$. This completes the proof. ■

Proof of Theorem 2.

In the proof, we establish that (3.2) holds on the event $\mathcal{A}$ (defined in (B.1), with $\mathbb{P}(\mathcal{A}^c) \leq n^{-1/2}$ by Lemma 1). The probability bound (3.1) then follows directly from the same proof strategy. Since, under Assumption 3, the bias and variance of the dyadic nonparametric estimator $\hat{\Pi}(\cdot, \cdot)$ achieve faster convergence rates than those characterized in Theorem 2, we ignore the first step for analytical convenience and always focus on

$$\begin{aligned} (\hat{\mathbf{a}}, \hat{\mathbf{b}}) \in \underset{\mathbf{a}, \mathbf{b} \in \mathbb{R}^{|\mathcal{S}_2|+1}}{\text{argmin}} \sum_{\substack{i' \in \mathcal{S}_2 \cup \{i\}, j' \in \mathcal{S}_2 \cup \{j\}, \\ (i', j') \neq (i, j)}} (A_{i'j'} - a_{i'} - b_{j'})^2 K_h(\hat{d}_{ii'}) K_h(\hat{d}_{jj'}), \\ \hat{A}_{ij} = \hat{a}_i + \hat{b}_j \end{aligned} \quad (\text{B.2})$$

in the following proof. It is easy to see that these two estimators, (A.2) and (B.2), share the same asymptotic properties.

Fix $i, j \in \mathcal{S}^c$, for any $i' \in \mathcal{S}_2 \cup \{i\}$ and $j' \in \mathcal{S}_2 \cup \{j\}$, for notational simplicity, define

$$\begin{aligned} K_{h,i'}^{(i)} &:= K(d_{ii'}/h), \quad K_{h,j'}^{(j)} := K(d_{jj'}/h), \quad K_{h,i'j'}^{(ij)} := K_{h,i'}^{(i)} K_{h,j'}^{(j)}, \\ \hat{K}_{h,i'}^{(i)} &:= K(\hat{d}_{ii'}/h), \quad \hat{K}_{h,j'}^{(j)} := K(\hat{d}_{jj'}/h), \quad \hat{K}_{h,i'j'}^{(ij)} := \hat{K}_{h,i'}^{(i)} \hat{K}_{h,j'}^{(j)}. \end{aligned}$$

By Lemma B.2, the two-way fixed-effects estimator (B.2) admits the following decomposition:

$$\begin{aligned}
\hat{A}_{ij} &= \frac{\sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'}^{(i)} A_{i'j}}{\sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'}^{(i)}} + \frac{\sum_{j' \in \mathcal{S}_2} \hat{K}_{h,j'}^{(j)} A_{ij'}}{\sum_{j' \in \mathcal{S}_2} \hat{K}_{h,j'}^{(j)}} - \frac{\sum_{i',j' \in \mathcal{S}_2} \hat{K}_{h,i'j'}^{(ij)} A_{i'j'}}{\sum_{i',j' \in \mathcal{S}_2} \hat{K}_{h,i'j'}^{(ij)}} \\
&= \underbrace{\frac{\sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'}^{(i)} f_{i'j}}{\sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'}^{(i)}} + \frac{\sum_{j' \in \mathcal{S}_2} \hat{K}_{h,j'}^{(j)} f_{ij'}}{\sum_{j' \in \mathcal{S}_2} \hat{K}_{h,j'}^{(j)}} - \frac{\sum_{i',j' \in \mathcal{S}_2} \hat{K}_{h,i'j'}^{(ij)} f_{i'j'}}{\sum_{i',j' \in \mathcal{S}_2} \hat{K}_{h,i'j'}^{(ij)}}}_{Q_1^{(ij)}} \\
&\quad + \underbrace{\frac{\sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'}^{(i)} \epsilon_{i'j}}{\sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'}^{(i)}} + \frac{\sum_{j' \in \mathcal{S}_2} \hat{K}_{h,j'}^{(j)} \epsilon_{ij'}}{\sum_{j' \in \mathcal{S}_2} \hat{K}_{h,j'}^{(j)}} - \frac{\sum_{i',j' \in \mathcal{S}_2} \hat{K}_{h,i'j'}^{(ij)} \epsilon_{i'j'}}{\sum_{i',j' \in \mathcal{S}_2} \hat{K}_{h,i'j'}^{(ij)}}}_{Q_2^{(ij)}} - \underbrace{\frac{\sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'j'}^{(ij)} f_{i'i'}}{\sum_{i',j' \in \mathcal{S}_2} \hat{K}_{h,i'j'}^{(ij)}}}_{Q_3^{(ij)}}
\end{aligned} \tag{B.3}$$

Here, Q_3 arises from the fact that the diagonal elements of A are zero. In addition, by Assumption 1(v), the Taylor expansion of $f_{i'j'}$ around (ζ_i, ζ_j) is given by

$$\begin{aligned}
f_{i'j'} &= f(\zeta_i, \zeta_j) + \nabla'_{\zeta_i} f(\zeta_i, \zeta_j)(\zeta_{i'} - \zeta_i) + \nabla'_{\zeta_j} f(\zeta_i, \zeta_j)(\zeta_{j'} - \zeta_j) \\
&\quad + (\zeta_{i'} - \zeta_i, \zeta_{j'} - \zeta_j)' \nabla^2 f(\zeta_i + \eta'_{1,i'j'}(\zeta_{i'} - \zeta_i), \zeta_j + \eta'_{2,i'j'}(\zeta_{j'} - \zeta_j))(\zeta_{i'} - \zeta_i, \zeta_{j'} - \zeta_j)
\end{aligned}$$

where $\eta_{1,i'j'}, \eta_{2,i'j'} \in [0, 1]^{d_\zeta}$. We apply the Taylor expansion on each term in Q_1 to obtain

$$\begin{aligned}
\frac{\sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'}^{(i)} f_{i'j}}{\sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'}^{(i)}} &= f(\zeta_i, \zeta_j) + \frac{\sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'}^{(i)} \nabla'_{\zeta_i} f(\zeta_i, \zeta_j)(\zeta_{i'} - \zeta_i)}{\sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'}^{(i)}} + \hat{R}_1^{(ij)}, \\
\frac{\sum_{j' \in \mathcal{S}_2} \hat{K}_{h,j'}^{(j)} f_{ij'}}{\sum_{j' \in \mathcal{S}_2} \hat{K}_{h,j'}^{(j)}} &= f(\zeta_i, \zeta_j) + \frac{\sum_{j' \in \mathcal{S}_2} \hat{K}_{h,j'}^{(j)} \nabla'_{\zeta_j} f(\zeta_i, \zeta_j)(\zeta_{j'} - \zeta_j)}{\sum_{j' \in \mathcal{S}_2} \hat{K}_{h,j'}^{(j)}} + \hat{R}_2^{(ij)}
\end{aligned}$$

and

$$\frac{\sum_{i' \in \mathcal{S}_2, j' \in \mathcal{S}_2} \hat{K}_{h,i'j'}^{(ij)} f_{i'j'}}{\sum_{i' \in \mathcal{S}_2, j' \in \mathcal{S}_2} \hat{K}_{h,i'j'}^{(ij)}} = f(\zeta_i, \zeta_j) + \frac{\sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'}^{(i)} \nabla'_{\zeta_i} f(\zeta_i, \zeta_j)(\zeta_{i'} - \zeta_i)}{\sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'}^{(i)}} + \frac{\sum_{j' \in \mathcal{S}_2} \hat{K}_{h,j'}^{(j)} \nabla'_{\zeta_j} f(\zeta_i, \zeta_j)(\zeta_{j'} - \zeta_j)}{\sum_{j' \in \mathcal{S}_2} \hat{K}_{h,j'}^{(j)}} + \hat{R}_3^{(ij)}$$

Here, $\hat{R}_1^{(ij)}, \hat{R}_2^{(ij)}, \hat{R}_3^{(ij)}$ are quadratic remaining terms:

$$\begin{aligned}
\hat{R}_1^{(ij)} &= \frac{\sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'}^{(i)} (\zeta_i - \zeta_{i'})' H_{i'}^{(ij)} (\zeta_i - \zeta_{i'})}{\sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'}^{(i)}}, \quad \hat{R}_2^{(ij)} = \frac{\sum_{j' \in \mathcal{S}_2} \hat{K}_{h,j'}^{(j)} (\zeta_j - \zeta_{j'})' H_{j'}^{(ij)} (\zeta_j - \zeta_{j'})}{\sum_{j' \in \mathcal{S}_2} \hat{K}_{h,j'}^{(j)}}, \\
\hat{R}_3^{(ij)} &= \frac{\sum_{i',j' \in \mathcal{S}_2} \hat{K}_{h,i'j'}^{(ij)} (\zeta_{i'} - \zeta_i, \zeta_{j'} - \zeta_j)' H_{i'j'}^{(ij)} (\zeta_{i'} - \zeta_i, \zeta_{j'} - \zeta_j)'}{\sum_{i',j' \in \mathcal{S}_2} \hat{K}_{h,i'j'}^{(ij)}}
\end{aligned} \tag{B.4}$$

where $H_{i'}^{(ij)} \in \mathbb{R}^{d_\zeta \times d_\zeta}$, $H_{j'}^{(ij)} \in \mathbb{R}^{d_\zeta \times d_\zeta}$, and $H_{i'j'}^{(ij)} \in \mathbb{R}^{2d_\zeta \times 2d_\zeta}$ are Hessians defined as

$$\begin{aligned}
H_{i'}^{(ij)} &= \nabla_{\zeta_i, \zeta_i}^2 f(\zeta_i + \eta'_{1,i'j'}(\zeta_{i'} - \zeta_i), \zeta_j), \quad H_{j'}^{(ij)} = \nabla_{\zeta_j, \zeta_j}^2 f(\zeta_i, \zeta_j + \eta'_{2,i'j'}(\zeta_{j'} - \zeta_j)), \\
H_{i'j'}^{(ij)} &= \nabla^2 f(\zeta_i + \eta'_{1,i'j'}(\zeta_{i'} - \zeta_i), \zeta_j + \eta'_{2,i'j'}(\zeta_{j'} - \zeta_j))
\end{aligned} \tag{B.5}$$

Then it is straightforward for verify that

$$\hat{A}_{ij} = \underbrace{f_{ij} + \hat{R}_1^{(ij)} + \hat{R}_2^{(ij)} + \hat{R}_3^{(ij)}}_{Q_1^{(ij)}} + Q_2^{(ij)} - Q_3^{(ij)}.$$

Bias Note that

$$\mathbb{E}(\hat{A}_{ij} - f_{ij} \mid \{\zeta_i\}_{i=1}^N) = \mathbb{E}(\hat{R}_1^{(ij)} + \hat{R}_2^{(ij)} + \hat{R}_3^{(ij)} \mid \{\zeta_i\}_{i=1}^N) + \mathbb{E}(Q_2^{(ij)} \mid \{\zeta_i\}_{i=1}^N) - \mathbb{E}(Q_3^{(ij)} \mid \{\zeta_i\}_{i=1}^N)$$

Let $\rho_{f''} := \max_{\zeta_1, \zeta_2 \in \text{supp}(\zeta)} \lambda_{\max}(\nabla_{\zeta_1, \zeta_2}^2 f(\zeta_1, \zeta_2))$, where $\lambda_{\max}(\cdot)$ denotes the largest singular value of a matrix. Under boundedness condition of graphon's second-order derivative (Assumption 1(v)), $\rho_{f''} < \infty$. In addition, by Lemma B.1 and the compactness of kernel's support (Assumption 4), when conditional on $\mathcal{A}$, or equivalently, with probability at least $1 - \delta n^{-1/2}$ (see Lemma 1)⁹,

$$|\hat{R}_1^{(ij)}| = \left| \frac{\sum_{i' \in \mathcal{S}_2} \mathbf{1}(\|\zeta_i - \zeta_{i'}\| \leq \eta h) \hat{K}_{h,i'}^{(i)}(\zeta_i - \zeta_{i'})' H_{i'}^{(i)}(\zeta_i - \zeta_{i'})}{\sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'}^{(i)}} \right| \leq \rho_{f''} \eta^2 h^2$$

Here and in the follows proof, we use δ to denote a generic positive constant that may vary from line to line and absorb multiplicative or additive constants. By the same argument, we obtain that, with probability at least $1 - \delta n^{-1/2}$, $\hat{R}_2^{(ij)} \leq \rho_{f''} \eta^2 h^2$ and $\hat{R}_3^{(ij)} \leq 4\rho_{f''} \eta^2 h^2$. Thus, with probability at least $1 - \delta n^{-1/2}$,

$$\left| \mathbb{E}(\hat{R}_1^{(ij)} + \hat{R}_2^{(ij)} + \hat{R}_3^{(ij)} \mid \{\zeta_i\}_{i=1}^N) \right| \leq 6\rho_{f''} \eta^2 h^2$$

Since conditional $\mathcal{A}$, the inequality above holds for any $\zeta_i, \zeta_j \in \{\zeta_i\}_{i \in \mathcal{S}^c}$, it follows that, with probability at least $1 - \delta n^{-1/2}$,

$$\max_{i,j \in \mathcal{S}^c} \left| \mathbb{E}(\hat{R}_1^{(ij)} + \hat{R}_2^{(ij)} + \hat{R}_3^{(ij)} \mid \{\zeta_i\}_{i=1}^N) \right| \leq 6\rho_{f''} \eta^2 h^2 \quad (\text{B.6})$$

In addition, Assumption 1(ii) and sample-splitting (see Algorithm 3) guarantee that $\{\epsilon_{i'j'} \mid i', j' \in \mathcal{S}_2\}$ are independent of $\{\hat{K}_{i'}^{(i)}, \hat{K}_{j'}^{(j)} \mid i', j' \in \mathcal{S}_2\}$. Then, we have, for any $\zeta_i, \zeta_j \in \{\zeta_i\}_{i \in \mathcal{S}^c}$,

$$\max_{i,j \in \mathcal{S}^c} \mathbb{E}(Q_2^{(ij)} \mid \{\zeta_i\}_{i=1}^N) = 0 \quad (\text{B.7})$$

To bound $\mathbb{E}(Q_3^{(ij)} \mid \{\zeta_i\}_{i=1}^N) = 0$, let ρ_K denote $\|K(\cdot)\|_{\max}$. Then, with probability at least $1 - \delta n^{-1/2}$,

$$\begin{aligned} \max_{i,j \in \mathcal{S}^c} \left| \frac{\sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'i'}^{(ij)} f_{i'i'}}{\sum_{i', j' \in \mathcal{S}_2} \hat{K}_{h,i'i'}^{(ij)}} \right| &\stackrel{(i)}{\leq} \frac{1}{D_2 n^2 h^{2d_\zeta}} \max_{i,j \in \mathcal{S}^c} \left| \sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'i'}^{(ij)} f_{i'i'} \right| \\ &\leq \frac{\rho_K \rho_f}{D_2 n^2 h^{2d_\zeta}} \max_{i,j \in \mathcal{S}^c} \left| \sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'}^{(i)} + \sum_{j' \in \mathcal{S}_2} \hat{K}_{h,i'}^{(j)} \right| \\ &\stackrel{(ii)}{\leq} \frac{2\rho_K \rho_f}{D_2 n^2 h^{2d_\zeta}} \bar{C}_2 n h^{d_\zeta} \\ &= \frac{2\bar{C}_2 \rho_K \rho_f}{D_2 n h^{d_\zeta}} \end{aligned} \quad (\text{B.8})$$

Here, (i) holds with probability at least $1 - \delta n^{-1/2}$ because of (B.27) in Lemma B.26, (ii) holds with probability at least $1 - \delta n^{-1/2}$ because of (B.20) in Lemma B.19. It follows that, with probability at least $1 - \delta n^{-1/2}$,

$$\max_{i,j \in \mathcal{S}^c} \left| \mathbb{E}(Q_3^{(ij)} \mid \{\zeta_i\}_{i=1}^N) \right| \leq \frac{2\bar{C}_2 \rho_K \rho_f}{D_2 n h^{d_\zeta}} \quad (\text{B.9})$$

⁹Although the Lemma 1 states the bound $\mathbb{P}(\mathcal{A}) \geq 1 - n^{-1/2}$, it follows easily from the proof that for any $\delta > 0$, there exist constants γ_1 and γ_2 such that $\mathbb{P}(\mathcal{A}) \geq 1 - \delta n^{-1/2}$.

Therefore, combining (B.6), (B.7), and (B.9), we conclude that, with probability at least $1 - \delta n^{-1/2}$,

$$\max_{i,j \in \mathcal{S}^c} \mathbb{E}(\hat{A}_{ij} - f_{ij} \mid \{\zeta_i\}_{i=1}^N) = O_p \left(h^2 + \frac{1}{nh^{d_\zeta}} \right)$$

Variance For notational simplicity, define the counterparts of $(\hat{R}_1^{(ij)}, \hat{R}_2^{(ij)}, \hat{R}_3^{(ij)})$ evaluated at the population pseudo distance $d_{ii'}$ as follows

$$\begin{aligned} R_1^{(ij)} &= \frac{\sum_{i' \in \mathcal{S}_2} K_{h,i'}^{(i)} (\zeta_i - \zeta_{i'})' H_{i'}^{(i)} (\zeta_i - \zeta_{i'})}{\sum_{i' \in \mathcal{S}_2} K_{h,i'}^{(i)}}, \quad R_2^{(ij)} = \frac{\sum_{j' \in \mathcal{S}_2} K_{h,j'}^{(j)} (\zeta_j - \zeta_{j'})' H_{j'}^{(j)} (\zeta_j - \zeta_{j'})}{\sum_{j' \in \mathcal{S}_2} K_{h,j'}^{(j)}}, \\ R_3^{(ij)} &= \frac{\sum_{i',j' \in \mathcal{S}_2} K_{h,i'j'}^{(ij)} (\zeta_{i'} - \zeta_{i'}, \zeta_{j'} - \zeta_{j'})' H_{i'j'}^{(ij)} (\zeta_{i'} - \zeta_{i'}, \zeta_{j'} - \zeta_{j'})'}{\sum_{i',j' \in \mathcal{S}_2} K_{h,i'j'}^{(ij)}} \end{aligned} \quad (\text{B.10})$$

Then,

$$\hat{A}_{ij} = f_{ij} + \sum_{k=1,2,3} R_k^{(ij)} + \sum_{k=1,2,3} (\hat{R}_k^{(ij)} - R_k^{(ij)}) + Q_2^{(ij)} - Q_3^{(ij)},$$

and, by Cauchy-Schwarz inequality, the variance of the imputation estimator is bounded by:

$$\text{Var}(\hat{A}_{ij} \mid \{\zeta_i\}_{i=1}^N) = 3\text{Var}\left(\sum_{k=1,2,3} (\hat{R}_k^{(ij)} - R_k^{(ij)}) \mid \{\zeta_i\}_{i=1}^N\right) + 3\text{Var}(Q_2^{(ij)} \mid \{\zeta_i\}_{i=1}^N) + 3\text{Var}(Q_3^{(ij)} \mid \{\zeta_i\}_{i=1}^N) \quad (\text{B.11})$$

Step 1: Bound on $\max_{i,j \in \mathcal{S}^c} \text{Var}(\sum_{k=1,2,3} (\hat{R}_k^{(ij)} - R_k^{(ij)}) \mid \{\zeta_i\}_{i=1}^N)$ Note that for any positive constant δ , there exist constants $\underline{C}_1, \underline{C}_2, M > 0$ such that with probability at least $1 - \delta n^{-1/2}$,

$$\begin{aligned} &\max_{i \in \mathcal{S}^c} \text{Var}(\hat{R}_1^{(ij)} - R_1^{(ij)} \mid \{\zeta_i\}_{i=1}^N) \\ &= \max_{i \in \mathcal{S}^c} \text{Var}\left(\frac{\sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'}^{(i)} (\zeta_i - \zeta_{i'})' H_{i'}^{(i)} (\zeta_i - \zeta_{i'})}{\sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'}^{(i)}} - \frac{\sum_{i' \in \mathcal{S}_2} K_{h,i'}^{(i)} (\zeta_i - \zeta_{i'})' H_{i'}^{(i)} (\zeta_i - \zeta_{i'})}{\sum_{i' \in \mathcal{S}_2} K_{h,i'}^{(i)}} \mid \{\zeta_i\}_{i=1}^N\right) \\ &\stackrel{(i)}{\leq} \frac{1}{\min\{\underline{C}_1, \underline{C}_2\} n^2 h^{2d_\zeta}} \max_{i \in \mathcal{S}^c} \mathbb{E}\left(\sum_{i' \in \mathcal{S}_2} \mathbf{1}(d_{ii'} \leq h \vee \hat{d}_{ii'} \leq h) (\zeta_i - \zeta_{i'})' H_{i'}^{(i)} (\zeta_i - \zeta_{i'}) (\hat{K}_{h,i'}^{(i)} - K_{h,i'}^{(i)}) \mid \{\zeta_i\}_{i=1}^N\right)^2 \\ &\stackrel{(ii)}{\leq} \frac{\bar{K}^2 (\delta_{N,n}/h)^2}{\min\{\underline{C}_1, \underline{C}_2\} n^2 h^{2d_\zeta}} \max_{i \in \mathcal{S}^c} \mathbb{E}\left(\sum_{i' \in \mathcal{S}_2} \mathbf{1}(d_{ii'} \leq h \vee \hat{d}_{ii'} \leq h) (\zeta_i - \zeta_{i'})' H_{i'}^{(i)} (\zeta_i - \zeta_{i'}) \mid \{\zeta_i\}_{i=1}^N\right)^2 \\ &\stackrel{(iii)}{\leq} \frac{\bar{K}^2 (\delta_{N,n}/h)^2 \rho_{f''}^2 h^4 \max\{\eta^4, \lambda^{-4}\}}{\min\{\underline{C}_1, \underline{C}_2\} n^2 h^{2d_\zeta}} \max_{i \in \mathcal{S}^c} \mathbb{E}\left(\sum_{i' \in \mathcal{S}_2} \mathbf{1}(d_{ii'} \leq h \vee \hat{d}_{ii'} \leq h) \mid \{\zeta_i\}_{i=1}^N\right)^2 \\ &\leq \frac{2\bar{K}^2 (\delta_{N,n}/h)^2 \rho_{f''}^2 h^4 \max\{\eta^4, \lambda^{-4}\}}{\min\{\underline{C}_1, \underline{C}_2\} n^2 h^{2d_\zeta}} \max_{i \in \mathcal{S}^c} \left(\mathbb{E}\left(\sum_{i' \in \mathcal{S}_2} \mathbf{1}(d_{ii'} \leq h) \mid \{\zeta_i\}_{i=1}^N\right)^2 + \mathbb{E}\left(\sum_{i' \in \mathcal{S}_2} \mathbf{1}(\hat{d}_{ii'} \leq h) \mid \{\zeta_i\}_{i=1}^N\right)^2\right) \\ &\stackrel{(iv)}{\leq} \frac{2\bar{K}^2 (\delta_{N,n}/h)^2 \rho_{f''}^2 h^4 \max\{\eta^4, \lambda^{-4}\}}{\min\{\underline{C}_1, \underline{C}_2\} n^2 h^{2d_\zeta}} M n^2 h^{2d_\zeta} \\ &= O(\delta_{NT}^2 h^2) \end{aligned} \quad (\text{B.12})$$

Here, inequality (i) is from (B.19) and (B.20) in Lemma B.3, inequality (ii) holds because Assumption 2 and Lemma 1, inequality (iii) follows from compact support of kernel function (Assumption 4) and Lemma B.1. Inequality (iv) follows from (B.19) and (B.20) in Lemma B.3, by replacing K with uniform kernel on $[-1, 1]$. By the same argument, we can also show that, with probability at least $1 -$

$\delta n^{-1/2}$, $\max_{i \in \mathcal{S}^c} \text{Var} \left(\hat{R}_2^{(ij)} - R_2^{(ij)} \mid \{\zeta_i\}_{i=1}^N \right) = O(\delta_{N,n}^2 h^2)$ and $\max_{i \in \mathcal{S}^c} \text{Var} \left(\hat{R}_3^{(ij)} - R_3^{(ij)} \mid \{\zeta_i\}_{i=1}^N \right) = O(\delta_{N,n}^2 h^2)$. Therefore, we have, with probability at least $1 - \delta n^{-1/2}$,

$$\max_{i,j \in \mathcal{S}^c} \text{Var} \left(\sum_{k=1,2,3} (\hat{R}_k^{(ij)} - R_k^{(ij)}) \mid \{\zeta_i\}_{i=1}^N \right) = O(\delta_{N,n}^2 h^2). \quad (\text{B.13})$$

Step 2: Bound on $\max_{i,j \in \mathcal{S}^c} \text{Var}(Q_2^{(ij)} \mid \{\zeta_i\}_{i=1}^N)$ Note that for $i, j \in \mathcal{S}^c$, (i) $\{\epsilon_{i'j'} \mid i' \in \mathcal{S}_2, j' \in \mathcal{S}_2\}$ is independent of $\{\hat{d}_{ii'} \mid i = 1, \dots, N, i' \in \mathcal{S}_2\}$ conditional on $\{\zeta_i\}_{i=1}^N$ (Assumption 1(ii)), (ii) $\hat{K}_{ij}^{(ij)}$ is uniformly bounded (Assumption 4), and (iii) conditional $\{\zeta_i\}_{i=1}^N, \{\epsilon_{i'j'} \mid i', j' \in \mathcal{S}_2\}$ is a set of independent sub-Gaussian random variables with $\sigma = 1$, (iv) fixing ζ_i, ζ_j , where $i, j \in \mathcal{S}^c$, $\text{Var}(\hat{K}_{h,i'}^{(i)} \epsilon_{i'j} \mid \{\zeta_i\}_{i \in \mathcal{S}^c}) = O(h^{d_\zeta})$, $\text{Var}(\hat{K}_{h,j'}^{(j)} \epsilon_{ij'} \mid \{\zeta_i\}_{i \in \mathcal{S}^c}) = O(h^{d_\zeta})$, $\text{Var}(\hat{K}_{h,i'j'}^{(ij)} \epsilon_{i'j'} \mid \{\zeta_i\}_{i \in \mathcal{S}^c}) = O(h^{2d_\zeta})$. Then, we employ Bernstein inequality (Lemma D.3) to show there exists constant $\bar{M}$ such that for any $t_1, t_2 > 0$,

$$\begin{aligned} \mathbb{P} \left(\left| \sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'}^{(i)} \epsilon_{i'j} \right| \geq t \mid \{\zeta_i\}_{i \in \mathcal{S}^c} \right) &\leq 2 \exp \left(\frac{-t^2}{\bar{M}^2 n h^{d_\zeta} + \frac{1}{3} \bar{M} t} \right) \\ \mathbb{P} \left(\left| \sum_{j' \in \mathcal{S}_2} \hat{K}_{h,j'}^{(j)} \epsilon_{ij'} \right| \geq t \mid \{\zeta_i\}_{i \in \mathcal{S}^c} \right) &\leq 2 \exp \left(\frac{-t^2}{\bar{M}^2 n h^{d_\zeta} + \frac{1}{3} \bar{M} t} \right) \\ \mathbb{P} \left(\left| \sum_{i',j' \in \mathcal{S}_2} \hat{K}_{h,i'j'}^{(ij)} \epsilon_{i'j'} \right| \geq t \mid \{\zeta_i\}_{i \in \mathcal{S}^c} \right) &\leq 2 \exp \left(\frac{-t^2}{\bar{M}^2 n^2 h^{2d_\zeta} + \frac{1}{3} \bar{M} t} \right) \end{aligned}$$

and

$$\begin{aligned} \max_{i \in \mathcal{S}^c} \mathbb{P} \left(\left| \sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'}^{(i)} \epsilon_{i'j} \right| \geq t \mid \{\zeta_i\}_{i \in \mathcal{S}^c} \right) &\leq 2N \exp \left(\frac{-t^2}{\bar{M}^2 n h^{d_\zeta} + \frac{1}{3} \bar{M} t} \right) \\ \max_{i \in \mathcal{S}^c} \mathbb{P} \left(\left| \sum_{j' \in \mathcal{S}_2} \hat{K}_{h,j'}^{(j)} \epsilon_{ij'} \right| \geq t \mid \{\zeta_i\}_{i \in \mathcal{S}^c} \right) &\leq 2N \exp \left(\frac{-t^2}{\bar{M}^2 n h^{d_\zeta} + \frac{1}{3} \bar{M} t} \right) \\ \max_{i,j \in \mathcal{S}^c} \mathbb{P} \left(\left| \sum_{i',j' \in \mathcal{S}_2} \hat{K}_{h,i'j'}^{(ij)} \epsilon_{i'j'} \right| \geq t \mid \{\zeta_i\}_{i \in \mathcal{S}^c} \right) &\leq 2N \exp \left(\frac{-t^2}{\bar{M}^2 n^2 h^{2d_\zeta} + \frac{1}{3} \bar{M} t} \right) \end{aligned}$$

Then there exists $M_1, M_2 > 0$ such that when $t = M_1 \sqrt{n h^{d_\zeta} \log N}, M_2 n h^{d_\zeta} \sqrt{\log N}$,

$$\begin{aligned} \mathbb{P} \left(\max_{i \in \mathcal{S}^c} \left| \sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'}^{(i)} \epsilon_{i'j} \right| \geq M_1 \sqrt{n h^{d_\zeta} \log N} \mid \{\zeta_i\}_{i \in \mathcal{S}^c} \right) &\leq \delta n^{-1/2} \\ \mathbb{P} \left(\max_{i \in \mathcal{S}^c} \left| \sum_{j' \in \mathcal{S}_2} \hat{K}_{h,j'}^{(j)} \epsilon_{ij'} \right| \geq M_1 \sqrt{n h^{d_\zeta} \log N} \mid \{\zeta_i\}_{i \in \mathcal{S}^c} \right) &\leq \delta n^{-1/2} \\ \mathbb{P} \left(\max_{i,j \in \mathcal{S}^c} \left| \sum_{i',j' \in \mathcal{S}_2} \hat{K}_{h,i'j'}^{(ij)} \epsilon_{i'j'} \right| \geq M_2 n h^{d_\zeta} \sqrt{\log N} \mid \{\zeta_i\}_{i \in \mathcal{S}^c} \right) &\leq \delta n^{-1/2} \end{aligned} \quad (\text{B.14})$$

We show that with probability at least $1 - \delta n^{-1/2}$,

$$\begin{aligned} \max_{i,j \in \mathcal{S}^c} \text{Var} \left(\frac{\sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'}^{(i)} \epsilon_{i'j}}{\sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'}^{(i)}} \mid \{\zeta_i\}_{i=1}^N \right) &\stackrel{(i)}{\leq} \frac{1}{\underline{C}_2^2 n^2 h^{2d_\zeta}} \mathbb{E} \left(\sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'}^{(i)} \epsilon_{i'j} \mid \{\zeta_i\}_{i=1}^N \right)^2 \\ &\stackrel{(ii)}{\leq} \frac{M_1^2 n h^{d_\zeta} \log N}{\underline{C}_2^2 n^2 h^{2d_\zeta}} \\ &= \frac{M_1^2 \log N}{\underline{C}_2^2 n h^{d_\zeta}} \end{aligned} \quad (\text{B.15})$$

Here, (i) holds with probability at least $1 - \delta n^{-1/2}$ because of (B.20) in Lemma B.3, and (ii), holding probability at least $1 - \delta n^{-1/2}$, follows from (B.14).¹⁰ Using the same argument, we also show that $\max_{i,j \in \mathcal{S}^c} \text{Var} \left(\frac{\sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'}^{(i)} \epsilon_{i'j}}{\sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'}^{(i)}} \mid \{\zeta_i\}_{i=1}^N \right) \leq \frac{M_2^2 \log N}{\underline{C}_2^2 n h^{d_\zeta}}$ with probability at least $1 - \delta n^{-1/2}$.

In addition, with probability at least $1 - \delta n^{-1/2}$:

$$\begin{aligned} \max_{i,j \in \mathcal{S}^c} \text{Var} \left(\frac{\sum_{i',j' \in \mathcal{S}_2} \hat{K}_{h,i'j'}^{(ij)} \epsilon_{i'j'}}{\sum_{i',j' \in \mathcal{S}_2} \hat{K}_{h,i'j'}^{(ij)}} \mid \{\zeta_i\}_{i=1}^N \right) &\stackrel{(i)}{\leq} \frac{1}{\underline{D}_2^2 n^4 h^{4d_\zeta}} \mathbb{E} \left(\sum_{i',j' \in \mathcal{S}_2} \hat{K}_{h,i'j'}^{(ij)} \epsilon_{i'j'} \mid \{\zeta_i\}_{i=1}^N \right)^2 \\ &\stackrel{(ii)}{\leq} \frac{M_2^2 n^2 h^{2d_\zeta} \log N}{\underline{D}_2^2 n^4 h^{4d_\zeta}} \\ &= \frac{M_2^2 \log N}{\underline{D}_2^2 n^2 h^{2d_\zeta}} \end{aligned} \quad (\text{B.16})$$

Here, (i) holds with probability at least $1 - \delta n^{-1/2}$ because of (B.27) in Lemma B.3, and (ii), holding probability at least $1 - \delta n^{-1/2}$, follows from (B.14). Therefore, with probability at least $1 - \delta n^{-1/2}$,

$$\max_{i,j \in \mathcal{S}^c} \text{Var}(Q_2^{(ij)} \mid \{\zeta_i\}_{i=1}^N) = O \left(\frac{\log N}{n h^{d_\zeta}} + \frac{\log N}{n^2 h^{2d_\zeta}} \right) = O \left(\frac{\log N}{n h^{d_\zeta}} \right) \quad (\text{B.17})$$

Step 3: Bound on $\max_{i,j \in \mathcal{S}^c} \text{Var}(Q_3^{(ij)} \mid \{\zeta_i\}_{i=1}^N)$ Note that, with probability at least $1 - \delta n^{-1/2}$,

$$\max_{i,j \in \mathcal{S}^c} \text{Var}(Q_3^{(ij)} \mid \{\zeta_i\}_{i=1}^N) \leq \max_{i,j \in \mathcal{S}^c} (\mathbb{E}(Q_3^{(ij)} \mid \{\zeta_i\}_{i=1}^N))^2 \stackrel{(i)}{\leq} \frac{4\bar{C}_2^2 \rho_K^2 \rho_f^2}{\underline{D}_2^2 n^2 h^{2d_\zeta}} \quad (\text{B.18})$$

where (i) follows from (B.9).

Therefore, combining (B.11), (B.13), (B.17), and (B.18) yields that, with probability at least $1 - \delta n^{-1/2}$,

$$\max_{i,j \in \mathcal{S}^c} \text{Var}(\hat{A}_{ij} - f_{ij} \mid \{\zeta_i\}_{i=1}^N) = O \left(\delta_{N,n}^2 h^2 + \frac{\log N}{n h^{d_\zeta}} + \frac{1}{n^2 h^{2d_\zeta}} \right) = O \left(\delta_{N,n}^2 h^2 + \frac{\log N}{n h^{d_\zeta}} \right)$$

We complete the proof. $\blacksquare$

Lemma B.2 *The solution of (B.2) is (B.3).*

Proof of Lemma B.2. Without loss of generality, we impose the normalization $\sum_{j \in \mathcal{S}_2} \hat{K}_{h,j'}^{(j)} \hat{b}_{j'} = 0$. For any $i, j \in \mathcal{S}^c$, the first-order conditions with respect to a_i and b_j are given by

$$\begin{aligned} \sum_{j' \in \mathcal{S}_2} \hat{K}_{h,j'}^{(j)} A_{ij'} - \sum_{j' \in \mathcal{S}_2} \hat{K}_{h,j'}^{(j)} \hat{a}_i - \underbrace{\sum_{j' \in \mathcal{S}_2} \hat{K}_{h,j'}^{(j)} \hat{b}_{j'}}_{=0} &= 0, \\ \sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'}^{(i)} A_{i'j} - \sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'}^{(i)} \hat{b}_j - \sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'}^{(i)} \hat{a}_{i'} &= 0. \end{aligned}$$

Therefore, we have

$$\hat{a}_i = \frac{\sum_{j' \in \mathcal{S}_2} \hat{K}_{h,j'}^{(j)} A_{ij'}}{\sum_{j' \in \mathcal{S}_2} \hat{K}_{h,j'}^{(j)}}, \quad \hat{b}_j = \frac{\sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'}^{(i)} A_{i'j} - \sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'}^{(i)} \hat{a}_{i'}}{\sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'}^{(i)}}.$$

¹⁰Since inequalities (B.14) do not depend on $\{\zeta_i\}_{i \in \mathcal{S}^c}$, we can drop the conditioning.

For any $\tilde{i} \in \mathcal{S}_2$, the first-order condition with respect to $a_{\tilde{i}}$ is given by

$$\sum_{j' \in \mathcal{S}_2} \hat{K}_{h,j'}^{(j)} A_{\tilde{i}j'} - \sum_{j' \in \mathcal{S}_2} \hat{K}_{h,j'}^{(j)} \hat{a}_{\tilde{i}} - \underbrace{\sum_{j' \in \mathcal{S}_2} \hat{K}_{h,j'}^{(j)} \hat{b}_{j'} + \hat{K}_{h,j}^{(j)} A_{\tilde{i}j} - \hat{K}_{h,j}^{(j)} \hat{a}_{\tilde{i}} - \hat{K}_{h,j}^{(j)} \hat{b}_j}_{=0} = 0$$

Thus,

$$\hat{a}_{\tilde{i}} = \frac{\sum_{j' \in \mathcal{S}_2} \hat{K}_{h,j'}^{(j)} A_{\tilde{i}j'} - \hat{K}_{h,j}^{(j)} \hat{b}_j}{\sum_{j' \in \mathcal{S}_2} \hat{K}_{h,j'}^{(j)} + \hat{K}_{h,j}^{(j)}}$$

Then,

$$\begin{aligned} \sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'}^{(i)} \hat{b}_j &= \sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'}^{(i)} A_{i'j} - \sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'}^{(i)} \frac{\sum_{j' \in \mathcal{S}_2} \hat{K}_{h,j'}^{(j)} A_{i'j'} - \hat{K}_{h,j}^{(j)} \hat{b}_j}{\sum_{j' \in \mathcal{S}_2} \hat{K}_{h,j'}^{(j)} + \hat{K}_{h,j}^{(j)}} \\ \Rightarrow \hat{b}_j &= \frac{\sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'}^{(i)} A_{i'j}}{\sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'}^{(i)}} - \frac{\sum_{i' \in \mathcal{S}_2} \sum_{j' \in \mathcal{S}_2} \hat{K}_{h,i'}^{(i)} \hat{K}_{h,j'}^{(j)} A_{i'j'}}{\sum_{i' \in \mathcal{S}_2} \sum_{j' \in \mathcal{S}_2} \hat{K}_{h,i'}^{(i)} \hat{K}_{h,j'}^{(j)}} \end{aligned}$$

Let $\hat{A}_{ij} = \hat{a}_i + \hat{b}_j$. We complete the proof. ■

Lemma B.3 *Under conditions in Theorem 2, for any constant $\delta > 0$, there exist constants $0 < \underline{C}_1 < \bar{C}_1 < \infty$ and $0 < \underline{C}_2 < \bar{C}_2 < \infty$ such that for all $i, j \in \mathcal{S}^c$,*

$$\mathbb{P} \left(\underline{C}_1 n h^{d_\zeta} \leq \min_{i \in \mathcal{S}^c} \sum_{i' \in \mathcal{S}_2} K_{h,i'}^{(i)} \leq \max_{i \in \mathcal{S}^c} \sum_{i' \in \mathcal{S}_2} K_{h,i'}^{(i)} \leq \bar{C}_1 n h^{d_\zeta} \right) \geq 1 - \delta n^{-1/2} \quad (\text{B.19})$$

and

$$\mathbb{P} \left(\underline{C}_2 n h^{d_\zeta} \leq \min_{i \in \mathcal{S}^c} \sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'}^{(i)} \leq \max_{i \in \mathcal{S}^c} \sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'}^{(i)} \leq \bar{C}_2 n h^{d_\zeta} \right) \geq 1 - \delta n^{-1/2} \quad (\text{B.20})$$

In addition, there exists constant $\bar{C}_3 > 0$ such that:

$$\begin{aligned} \mathbb{P} \left(\max_{i \in \mathcal{S}^c} \left| \sum_{i' \in \mathcal{S}_2} K_{h,i'}^{(i)} (\zeta_i - \zeta_{i'})' H_{i'}^{(ij)} (\zeta_i - \zeta_{i'}) - \sum_{i' \in \mathcal{S}_2} \mathbb{E}(K_{h,i'}^{(i)} (\zeta_i - \zeta_{i'})' H_{i'}^{(ij)} (\zeta_i - \zeta_{i'}) \mid \{\zeta_i\}_{i \in \mathcal{S}^c}) \right| \right. \\ \left. \geq \bar{C}_3 \sqrt{n h^{d_\zeta + 4} \log N} \mid \{\zeta_i\}_{i \in \mathcal{S}^c} \right) \leq \delta n^{-1/2} \end{aligned} \quad (\text{B.21})$$

Proof of Lemma B.3. Fixing $\{\zeta_i\}_{i \in \mathcal{S}^c}$, by Assumption 1(iv) and Assumption 4, it is straightforward to verify that for any $i \in \mathcal{S}^c$,

$$\text{Var}(K_{h,i'}^{(i)} - \mathbb{E}(K_{h,i'}^{(i)} \mid \{\zeta_i\}_{i \in \mathcal{S}^c})) = \text{Var}(K_{h,i'}^{(i)} - \mathbb{E}(K_{h,i'}^{(i)} \mid \{\zeta_i\}_{i \in \mathcal{S}^c}) \mid \{\zeta_i\}_{i \in \mathcal{S}^c}) = O(h^{d_\zeta})$$

In addition, by the boundedness of $K_{h,i'}^{(i)}$ (Assumption 4), we apply Bernstein inequality of independent random variables (Wainwright (2019)) to demonstrate that there exists a constant $M > 0$ such that for any $t > 0$,

$$\mathbb{P} \left(\left| \sum_{i' \in \mathcal{S}_2} (K_{h,i'}^{(i)} - \mathbb{E}(K_{h,i'}^{(i)} \mid \{\zeta_i\}_{i \in \mathcal{S}^c})) \right| \geq t \mid \{\zeta_i\}_{i \in \mathcal{S}^c} \right) \leq 2 \exp \left(\frac{-t^2}{M^2 n h^{d_\zeta} + \frac{1}{3} M t} \right)$$

and therefore,

$$\mathbb{P} \left(\max_{i \in \mathcal{S}^c} \left| \sum_{i' \in \mathcal{S}_2} \left(K_{h,i'}^{(i)} - \mathbb{E}(K_{h,i'}^{(i)} \mid \{\zeta_i\}_{i \in \mathcal{S}^c}) \right) \right| \geq t \mid \{\zeta_i\}_{i \in \mathcal{S}^c} \right) \leq 2N \exp \left(\frac{-t^2}{M^2 n h^{d_\zeta} + \frac{1}{3} M t} \right)$$

It follows that we can find a sufficiently large constant C_1 such that when $t = C_1 \sqrt{n h^{d_\zeta} \log N}$,

$$\mathbb{P} \left(\max_{i \in \mathcal{S}^c} \left| \sum_{i' \in \mathcal{S}_2} \left(K_{h,i'}^{(i)} - \mathbb{E}(K_{h,i'}^{(i)} \mid \{\zeta_i\}_{i \in \mathcal{S}^c}) \right) \right| \geq C_1 \sqrt{n h^{d_\zeta} \log N} \right) \leq \delta n^{-1/2} \quad (\text{B.22})$$

In addition, for any $\zeta_i \in \{\zeta_i\}_{i \in \mathcal{S}^c}$, we have

$$\begin{aligned} \mathbb{E}(K_{h,i'}^{(i)} \mid \{\zeta_i\}_{i \in \mathcal{S}^c}) &= \mathbb{E}(K_{h,i'}^{(i)} \mid \zeta_i) = \int K \left(\frac{d_{ii'}}{h} \right) d\mathbb{P}_\zeta(\zeta_{i'}) \\ &= \int_{\{\|\zeta_{i'} - \zeta_i\| \leq \lambda^{-1} h\}} K \left(\frac{d_{ii'}}{h} \right) d\mathbb{P}_\zeta(\zeta_{i'}) \\ &\stackrel{(i)}{\geq} C_2 \mathbb{P}(\|\zeta_i - \zeta_{i'}\| \leq \lambda^{-1} h \mid \zeta_i) \\ &\stackrel{(ii)}{\geq} C_2 \underline{c} (\lambda^{-1} h)^{d_\zeta} \end{aligned} \quad (\text{B.23})$$

Inequality (i) follows from the Lipschitz continuity of kernel $K(\cdot)$ and $K(0) > 0$ (see Assumption 4), and inequality (ii) holds because of Assumption 1(iv). By the boundedness of kernel K we can also show that there exists constant $C_3 > 0$ such that

$$\mathbb{E}(K_{h,i'}^{(i)} \mid \{\zeta_i\}_{i \in \mathcal{S}^c}) \leq C_3 \bar{c} (\lambda^{-1} h)^{d_\zeta}, \quad \forall \zeta_i \quad (\text{B.24})$$

Combining (B.22), (B.23), and (B.24), we have

$$\begin{aligned} \mathbb{P} \left(C_2 \underline{c} n (\lambda^{-1} h)^{d_\zeta} - C_1 \sqrt{n h^{d_\zeta} \log N} \leq \min_{i \in \mathcal{S}^c} \sum_{i' \in \mathcal{S}_2} K_{h,i'}^{(i)} \leq \max_{i \in \mathcal{S}^c} \sum_{i' \in \mathcal{S}_2} K_{h,i'}^{(i)} \leq \right. \\ \left. C_3 \bar{c} n (\lambda^{-1} h)^{d_\zeta} + C_1 \sqrt{n h^{d_\zeta} \log N} \mid \{\zeta_i\}_{i \in \mathcal{S}^c} \right) \geq 1 - \delta n^{-1/2} \end{aligned}$$

Therefore, there exist constants $0 < \underline{C}_1 < \bar{C}_1 < \infty$ such that¹¹

$$\mathbb{P} \left(\underline{C}_1 n h^{d_\zeta} \leq \min_{i \in \mathcal{S}^c} \sum_{i' \in \mathcal{S}_2} K_{h,i'}^{(i)} \leq \max_{i \in \mathcal{S}^c} \sum_{i' \in \mathcal{S}_2} K_{h,i'}^{(i)} \leq \bar{C}_1 n h^{d_\zeta} \mid \{\zeta_i\}_{i \in \mathcal{S}_2} \right) \geq 1 - \delta n^{-1/2}$$

Since the inequality above holds for any realization of $\{\zeta_i\}_{i \in \mathcal{S}_2}$, we drop the conditioning and obtain (B.19).

Now let us turn to the derivation of (B.20). When conditional on event $\mathcal{A}$, or equivalently, with probability at least $1 - \delta n^{-1/2}$, we have

$$\begin{aligned} \max_{i \in \mathcal{S}^c} \left| \sum_{i' \in \mathcal{S}_2} (\hat{K}_{h,i'}^{(i)} - K_{h,i'}^{(i)}) \right| &\stackrel{(i)}{\leq} \max_{i \in \mathcal{S}^c} \sum_{i' \in \mathcal{S}_2} \mathbf{1}(\hat{d}_{ii'} \leq h) \bar{K} |\hat{d}_{ii'} - d_{ii'}|/h \\ &\stackrel{(ii)}{\leq} \max_{i \in \mathcal{S}^c} \sum_{i' \in \mathcal{S}_2} \mathbf{1}(\hat{d}_{ii'} \leq h) \bar{K} \delta_{N,n}/h \end{aligned}$$

where inequality (i) holds because of Assumption 4, and inequality (ii) comes from the definition of $\mathcal{A}$ in

¹¹ $\sqrt{n h^{d_\zeta} \log N}$ is asymptotically negligible compared to $n h^{d_\zeta}$ by conditions in Theorem 2.

as in (B.1). Since we require $\delta_{N,n}/h \rightarrow 0$ in Theorem 2, we employ the same argument used in proving (B.19) to show that there exists constants $0 < C_4 < C_5 < \infty$ such that ¹²

$$\begin{aligned} & \mathbb{P} \left(C_4 n h^{d_\zeta} \leq \min_{i \in \mathcal{S}^c} \sum_{i' \in \mathcal{S}_2} \mathbf{1}(\hat{d}_{ii'} \leq h) \leq \max_{i \in \mathcal{S}^c} \sum_{i' \in \mathcal{S}_2} \mathbf{1}(\hat{d}_{ii'} \leq h) \leq C_5 n h^{d_\zeta} \right) \geq 1 - \delta n^{-1/2} \\ \Rightarrow & \mathbb{P} \left(C_4 n h^{d_\zeta - 1} \delta_{N,n} \leq \min_{i \in \mathcal{S}^c} \left| \sum_{i' \in \mathcal{S}_2} (\hat{K}_{h,i'}^{(i)} - K_{h,i'}^{(i)}) \right| \leq \max_{i \in \mathcal{S}^c} \left| \sum_{i' \in \mathcal{S}_2} (\hat{K}_{h,i'}^{(i)} - K_{h,i'}^{(i)}) \right| \leq C_5 n h^{d_\zeta - 1} \delta_{N,n} \right) \geq 1 - \delta n^{-1/2} \end{aligned} \quad (\text{B.25})$$

Here, we employ Lemma 1 to obtain the second inequality.

Then combining (B.19) and (B.25), there exist $0 < \underline{C}_2 < \bar{C}_2 < \infty$ such that

$$\mathbb{P} \left(\underline{C}_2 n h^{d_\zeta} \leq \min_{i \in \mathcal{S}^c} \sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'}^{(i)} \leq \max_{i \in \mathcal{S}^c} \sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'}^{(i)} \leq \bar{C}_2 n h^{d_\zeta} \right) \geq 1 - \delta n^{-1/2}$$

Therefore, we can obtain (B.20).

To prove (B.21), fixing $\{\zeta_i\}_{i \in \mathcal{S}^c}$, by Assumption 1(iv), Assumption 1(v), and Assumption 4, it is straightforward to verify that

$$\text{Var}(K_{h,i'}^{(i)}(\zeta_i - \zeta_{i'})' H_{i'}^{(ij)}(\zeta_i - \zeta_{i'}) - \mathbb{E}(K_{h,i'}^{(i)}(\zeta_i - \zeta_{i'})' H_{i'}^{(ij)}(\zeta_i - \zeta_{i'}) \mid \{\zeta_i\}_{i \in \mathcal{S}^c}) \mid \{\zeta_i\}_{i \in \mathcal{S}^c}) = O(h^{d_\zeta + 4})$$

By the uniform boundedness of $K(\cdot)$, ζ_i , and $H_{i'}^{(ij)}$ (see Assumption 1(iv), Assumption 1(v), and Assumption 4), we employ Bernstein inequality of independent random variables (Wainwright (2019)) demonstrates that there exists a constant $M > 0$ such that for any $t > 0$,

$$\begin{aligned} & \mathbb{P} \left(\left| \sum_{i' \in \mathcal{S}_2} K_{h,i'}^{(i)}(\zeta_i - \zeta_{i'})' H_{i'}^{(ij)}(\zeta_i - \zeta_{i'}) - \sum_{i' \in \mathcal{S}_2} \mathbb{E}(K_{h,i'}^{(i)}(\zeta_i - \zeta_{i'})' H_{i'}^{(ij)}(\zeta_i - \zeta_{i'}) \mid \{\zeta_i\}_{i \in \mathcal{S}^c}) \right| \geq t \mid \{\zeta_i\}_{i \in \mathcal{S}^c} \right) \\ & \leq 2 \exp \left(\frac{-t^2}{M^2 n h^{d_\zeta + 4} + \frac{1}{3} M t} \right) \end{aligned}$$

and therefore,

$$\begin{aligned} & \mathbb{P} \left(\max_{i \in \mathcal{S}^c} \left| \sum_{i' \in \mathcal{S}_2} K_{h,i'}^{(i)}(\zeta_i - \zeta_{i'})' H_{i'}^{(ij)}(\zeta_i - \zeta_{i'}) - \sum_{i' \in \mathcal{S}_2} \mathbb{E}(K_{h,i'}^{(i)}(\zeta_i - \zeta_{i'})' H_{i'}^{(ij)}(\zeta_i - \zeta_{i'}) \mid \{\zeta_i\}_{i \in \mathcal{S}^c}) \right| \geq t \mid \{\zeta_i\}_{i \in \mathcal{S}^c} \right) \\ & \leq 2N \exp \left(\frac{-t^2}{M^2 n h^{d_\zeta + 4} + \frac{1}{3} M t} \right) \end{aligned}$$

It follows that we can find a sufficiently large constant $\bar{C}_3 > 0$ such that when $t = \bar{C}_3 \sqrt{n h^{d_\zeta + 4} \log n}$,

$$\begin{aligned} & \mathbb{P} \left(\max_{i \in \mathcal{S}^c} \left| \sum_{i' \in \mathcal{S}_2} K_{h,i'}^{(i)}(\zeta_i - \zeta_{i'})' H_{i'}^{(ij)}(\zeta_i - \zeta_{i'}) - \sum_{i' \in \mathcal{S}_2} \mathbb{E}(K_{h,i'}^{(i)}(\zeta_i - \zeta_{i'})' H_{i'}^{(ij)}(\zeta_i - \zeta_{i'}) \mid \{\zeta_i\}_{i \in \mathcal{S}^c}) \right| \right. \\ & \quad \left. \geq \bar{C}_3 \sqrt{n h^{d_\zeta + 4} \log N} \mid \{\zeta_i\}_{i \in \mathcal{S}^c} \right) \leq \delta n^{-1/2}, \end{aligned}$$

which is (B.21). ■

Lemma B.4 *Under conditions in Theorem 2, for any constant $\delta > 0$, there exist constants $0 < \underline{D}_1 <$*

¹²Here we drop the conditioning because it is straightforward to verify that the analysis does not depend on specific realization of $\{\zeta_i\}_{i \in \mathcal{S}^c}$.

$\bar{D}_1 < \infty$ and $0 < \underline{D}_2 < \bar{D}_2 < \infty$ such that for all $i, j \in \mathcal{S}^c$,

$$\mathbb{P}\left(\underline{D}_1 n^2 h^{2d_\zeta} \leq \min_{i,j \in \mathcal{S}^c} \sum_{i',j' \in \mathcal{S}_2} K_{h,i'j'}^{(ij)} \leq \max_{i,j \in \mathcal{S}^c} \sum_{i',j' \in \mathcal{S}_2} K_{h,i'j'}^{(ij)} \leq \bar{D}_1 n^2 h^{2d_\zeta}\right) \leq 1 - \delta n^{-1/2} \quad (\text{B.26})$$

$$\mathbb{P}\left(\underline{D}_2 n^2 h^{2d_\zeta} \leq \min_{i,j \in \mathcal{S}^c} \sum_{i',j' \in \mathcal{S}_2} \hat{K}_{h,i'j'}^{(ij)} \leq \max_{i,j \in \mathcal{S}^c} \sum_{i',j' \in \mathcal{S}_2} \hat{K}_{h,i'j'}^{(ij)} \leq \bar{D}_2 n^2 h^{2d_\zeta}\right) \leq 1 - \delta n^{-1/2} \quad (\text{B.27})$$

In addition, there exists constant $\bar{D}_3 > 0$ such that:

$$\begin{aligned} & \mathbb{P}\left(\max_{i,j \in \mathcal{S}^c} \left| \sum_{i',j' \in \mathcal{S}_2} (K_{h,i'j'}^{(ij)} (\zeta'_i - \zeta'_{i'}, \zeta'_j - \zeta'_{j'})) H_{i'j'}^{(ij)} (\zeta'_i - \zeta'_{i'}, \zeta'_j - \zeta'_{j'})' \right. \right. \\ & \quad \left. \left. - \mathbb{E}(K_{h,i'j'}^{(ij)} (\zeta'_i - \zeta'_{i'}, \zeta'_j - \zeta'_{j'})) H_{i'j'}^{(ij)} (\zeta'_i - \zeta'_{i'}, \zeta'_j - \zeta'_{j'})' \mid \{\zeta_i\}_{i \in \mathcal{S}^c}) \right| \right. \\ & \quad \left. \leq \bar{D}_3 \sqrt{n^3 h^{3d_\zeta + 4} \log N} \mid \{\zeta_i\}_{i \in \mathcal{S}^c} \right) \geq 1 - \delta n^{-1/2} \end{aligned} \quad (\text{B.28})$$

Proof of Lemma B.4. Fix $\zeta_i, \zeta_j \in \{\zeta_i\}_{i \in \mathcal{S}^c}$. Because $K_{h,i'j'}^{(ij)}$ is uniformly bounded and $\mathbb{E}(\mathbb{E}^2(K_{h,i'j'}^{(ij)} \mid \zeta_i, \zeta_j, \zeta_{i'}, \zeta_{j'}) \mid \zeta_i, \zeta_j) = O(h^{3d_\zeta})$, we apply Bernstein's inequality of U-statistics (see [Arcones \(1995, Theorem 2\)](#)) to show that there exists a sufficiently large constant $M > 0$ such that for any $\zeta_i, \zeta_j \in \{\zeta_i\}_{i \in \mathcal{S}^c}$, and for any $t > 0$,

$$\mathbb{P}\left(\left| \sum_{i',j' \in \mathcal{S}_2} K_{h,i'j'}^{(ij)} - \mathbb{E}\left(\sum_{i',j' \in \mathcal{S}_2} K_{h,i'j'}^{(ij)} \mid \{\zeta_i\}_{i \in \mathcal{S}^c}\right) \right| \geq t \mid \{\zeta_i\}_{i \in \mathcal{S}^c}\right) \leq 4 \exp\left(\frac{-t^2}{M^2 n^3 h^{3d_\zeta} + Mnt}\right)$$

and

$$\max_{i,j \in \mathcal{S}^c} \mathbb{P}\left(\left| \sum_{i',j' \in \mathcal{S}_2} K_{h,i'j'}^{(ij)} - \mathbb{E}\left(\sum_{i',j' \in \mathcal{S}_2} K_{h,i'j'}^{(ij)} \mid \{\zeta_i\}_{i \in \mathcal{S}^c}\right) \right| \geq t \mid \{\zeta_i\}_{i \in \mathcal{S}^c}\right) \leq 4N^2 \exp\left(\frac{-t^2}{M^2 n^3 h^{3d_\zeta} + Mnt}\right)$$

We can find a constant D_1 such that when $t = D_1 \sqrt{n^3 h^{3d_\zeta} \log N}$,

$$\mathbb{P}\left(\left| \sum_{i',j' \in \mathcal{S}_2} K_{h,i'j'}^{(ij)} - \mathbb{E}\left(\sum_{i',j' \in \mathcal{S}_2} K_{h,i'j'}^{(ij)} \mid \{\zeta_i\}_{i \in \mathcal{S}^c}\right) \right| \geq D_1 \sqrt{n^3 h^{3d_\zeta} \log N} \mid \{\zeta_i\}_{i \in \mathcal{S}^c}\right) \leq \delta n^{-1/2} \quad (\text{B.29})$$

In addition, for any $\zeta_i, \zeta_j \in \{\zeta_i\}_{i \in \mathcal{S}^c}$, we have

$$\begin{aligned} \mathbb{E}(K_{h,i'j'}^{(ij)} \mid \{\zeta_i\}_{i \in \mathcal{S}^c}) &= \mathbb{E}(K_{h,i'j'}^{(ij)} \mid \zeta_i, \zeta_j) = \int K\left(\frac{d_{ii'}}{h}\right) K\left(\frac{d_{jj'}}{h}\right) d\mathbb{P}_\zeta(\zeta_{i'}) d\mathbb{P}_\zeta(\zeta_{j'}) \\ &= \int_{\{\|\zeta_{i'} - \zeta_i\| \leq \lambda^{-1}h\}} K\left(\frac{d_{ii'}}{h}\right) d\mathbb{P}_\zeta(\zeta_{i'}) \int_{\{\|\zeta_{j'} - \zeta_j\| \leq \lambda^{-1}h\}} K\left(\frac{d_{jj'}}{h}\right) d\mathbb{P}_\zeta(\zeta_{j'}) \\ &\stackrel{(i)}{\geq} D_2 \mathbb{P}(\|\zeta_i - \zeta_{i'}\| \leq \lambda^{-1}h \mid \zeta_i) \mathbb{P}(\|\zeta_j - \zeta_{j'}\| \leq \lambda^{-1}h \mid \zeta_j) \\ &\stackrel{(ii)}{\geq} D_2 \underline{c}^2 (\lambda^{-1}h)^{2d_\zeta} \end{aligned} \quad (\text{B.30})$$

Inequality (i) follows from the Lipschitz continuity of kernel K and $K(0) > 0$ (see Assumption 4), and inequality (ii) holds because Assumption 1(iv). By the boundedness of kernel K we can also show that there exists constant $D_3 > 0$ such that, for any $\zeta_i, \zeta_j \in \{\zeta_i\}_{i \in \mathcal{S}^c}$,

$$\mathbb{E}(K_{h,i'j'}^{(ij)} \mid \{\zeta_i\}_{i \in \mathcal{S}^c}) \leq D_3 \underline{c}^2 (\lambda^{-1}h)^{2d_\zeta} \quad (\text{B.31})$$

Combining (B.29), (B.30), and (B.31),

$$\begin{aligned} \mathbb{P}\left(D_2 \bar{c}^2 n^2 (\lambda^{-1} h)^{2d_\zeta} - D_1 \sqrt{n^3 h^{3d_\zeta} \log N} \leq \min_{i,j \in \mathcal{S}^c} \sum_{i',j' \in \mathcal{S}_2} K_{h,i'j'}^{(ij)} \leq \max_{i,j \in \mathcal{S}^c} \sum_{i',j' \in \mathcal{S}_2} K_{h,i'j'}^{(ij)} \right. \\ \left. \leq D_3 \bar{c}^2 n^2 (\lambda^{-1} h)^{2d_\zeta} + C_1 \sqrt{n^3 h^{3d_\zeta} \log N} \mid \{\zeta_i\}_{i \in \mathcal{S}^c}\right) \leq 1 - \delta n^{-1/2} \end{aligned}$$

Thus, we can find constants $0 < \underline{D}_1 < \bar{D}_1 < \infty$ such that ¹³

$$\mathbb{P}\left(\underline{D}_1 n^2 h^{2d_\zeta} \leq \min_{i,j \in \mathcal{S}^c} \sum_{i',j' \in \mathcal{S}_2} K_{h,i'j'}^{(ij)} \leq \max_{i,j \in \mathcal{S}^c} \sum_{i',j' \in \mathcal{S}_2} K_{h,i'j'}^{(ij)} \leq \bar{D}_1 n^2 h^{2d_\zeta} \mid \{\zeta_i\}_{i \in \mathcal{S}^c}\right) \leq 1 - \delta n^{-1/2}$$

Since the inequality above does not depend on the realization of $\{\zeta_i\}_{i \in \mathcal{S}^c}$, we can drop the conditioning to obtain (B.26).

Now let us turn to the derivation of (B.27). Let $\rho_K := \|K(\cdot)\|_\infty < \infty$. When conditional on event $\mathcal{A}$, or equivalently, with probability at least $1 - \delta n^{-1/2}$ (by Lemma 1),

$$\begin{aligned} \max_{i,j \in \mathcal{S}^c} \left| \sum_{i',j' \in \mathcal{S}_2} (\hat{K}_{h,i'j'}^{(ij)} - K_{h,i'j'}^{(ij)}) \right| &\stackrel{(i)}{\leq} \max_{i,j \in \mathcal{S}^c} \sum_{i',j' \in \mathcal{S}_2} \mathbf{1}(\hat{d}_{ii'} \leq h) \mathbf{1}(\hat{d}_{jj'} \leq h) \bar{K} \rho_K \frac{|\hat{d}_{ii'} - d_{ii'}| + |\hat{d}_{jj'} - d_{jj'}|}{h} \\ &\quad + \max_{i,j \in \mathcal{S}^c} \sum_{i',j' \in \mathcal{S}_2} \mathbf{1}(\hat{d}_{ii'} \leq h) \mathbf{1}(\hat{d}_{jj'} \leq h) \bar{K}^2 \frac{|\hat{d}_{ii'} - d_{ii'}| |\hat{d}_{jj'} - d_{jj'}|}{h^2} \\ &\stackrel{(ii)}{\leq} \max_{i,j \in \mathcal{S}^c} \sum_{i',j' \in \mathcal{S}_2} \mathbf{1}(\hat{d}_{ii'} \leq h) \mathbf{1}(\hat{d}_{jj'} \leq h) \left(\bar{K} \rho_K \frac{\delta_{N,n}}{h} + \bar{K}^2 \frac{\delta_{N,n}^2}{h^2} \right) \end{aligned}$$

where inequality (i) holds because of Assumption 4, and inequality (ii) comes from the definition of $\mathcal{A}$ in as in (B.1). Since we require $\delta_{N,n}/h \rightarrow 0$, we can employ the same argument used in proving (B.26) to show that there exists constants $0 < D_4 < D_5 < \infty$ such that ¹⁴

$$\begin{aligned} \mathbb{P}\left(D_4 n^2 h^{2d_\zeta} \leq \min_{i,j \in \mathcal{S}^c} \sum_{i',j' \in \mathcal{S}_2} \mathbf{1}(\hat{d}_{ii'} \leq h) \mathbf{1}(\hat{d}_{jj'} \leq h) \leq \max_{i,j \in \mathcal{S}^c} \sum_{i',j' \in \mathcal{S}_2} \mathbf{1}(\hat{d}_{ii'} \leq h) \mathbf{1}(\hat{d}_{jj'} \leq h) \leq D_5 n^2 h^{2d_\zeta}\right) \\ \leq 1 - \delta n^{-1/2} \\ \Rightarrow \mathbb{P}\left(D_4 n^2 h^{2d_\zeta-1} \delta_{N,n} \leq \min_{i,j \in \mathcal{S}^c} \left| \sum_{i',j' \in \mathcal{S}_2} (\hat{K}_{h,i'j'}^{(ij)} - K_{h,i'j'}^{(ij)}) \right| \leq \max_{i,j \in \mathcal{S}^c} \left| \sum_{i',j' \in \mathcal{S}_2} (\hat{K}_{h,i'j'}^{(ij)} - K_{h,i'j'}^{(ij)}) \right| \leq D_5 n^2 h^{2d_\zeta-1} \delta_{N,n}\right) \\ \leq 1 - \delta n^{-1/2} \end{aligned} \tag{B.32}$$

Then, by combining (B.26) and (B.32), there exist $0 < \underline{D}_2 < \bar{D}_2 < \infty$ such that

$$\mathbb{P}\left(\underline{D}_2 n^2 h^{2d_\zeta} \leq \min_{i,j \in \mathcal{S}^c} \sum_{i',j' \in \mathcal{S}_2} \hat{K}_{h,i'j'}^{(ij)} \leq \max_{i,j \in \mathcal{S}^c} \sum_{i',j' \in \mathcal{S}_2} \hat{K}_{h,i'j'}^{(ij)} \leq \bar{D}_2 n^2 h^{2d_\zeta}\right) \leq 1 - \delta n^{-1/2}$$

So we complete the proof of (B.27).

To prove (B.28), let $g_{i'j'}^{(ij)}$ denote $K_{h,i'j'}^{(ij)}(\zeta'_i - \zeta'_{i'}, \zeta'_j - \zeta'_{j'}) H_{i'j'}^{(ij)}(\zeta'_i - \zeta'_{i'}, \zeta'_j - \zeta'_{j'})'$ for notational simplicity. Fixing any $\zeta_i, \zeta_j \in \{\zeta_i\}_{i \in \mathcal{S}^c}$, by the boundedness of $K_{h,i'j'}^{(ij)}$ and $H_{i'j'}^{(ij)}$ (Assumption 4 and Assumption 1(v)), we have that $g_{i'j'}^{(ij)}$ is uniformly bounded and $\mathbb{E}(\mathbb{E}^2(g_{h,i'j'}^{(ij)} \mid \zeta_i, \zeta_j, \zeta_{i'}) \mid \zeta_i, \zeta_j) = O(h^{3d_\zeta+4})$. Then, we apply Bernstein's inequality of U-statistics (see Arcones (1995, Theorem 2)) to show that there exists a

¹³ $\sqrt{n^3 h^{3d_\zeta} \log N}$ is asymptotically negligible compared to $n^2 h^{2d_\zeta}$ by conditions in Theorem 2.

¹⁴ Here we drop the conditioning because it is straightforward verify that the analysis does not depend on specific realization of $\{\zeta_i\}_{i \in \mathcal{S}^c}$.

sufficiently large constant M such that for any $t > 0$,

$$\mathbb{P} \left(\left| \sum_{i', j' \in \mathcal{S}_2} \left(g_{i'j'}^{(ij)} - \mathbb{E} \left(g_{i'j'}^{(ij)} \mid \{\zeta_i\}_{i \in \mathcal{S}^c} \right) \right) \right| \geq t \mid \{\zeta_i\}_{i \in \mathcal{S}^c} \right) \leq 4 \exp \left(\frac{-t^2}{M^2 n^3 h^{3d_\zeta+4} + M n h^2 t} \right)$$

and

$$\mathbb{P} \left(\max_{i, j \in \mathcal{S}^c} \left| \sum_{i', j' \in \mathcal{S}_2} \left(g_{i'j'}^{(ij)} - \mathbb{E} \left(g_{i'j'}^{(ij)} \mid \{\zeta_i\}_{i \in \mathcal{S}^c} \right) \right) \right| \geq t \mid \{\zeta_i\}_{i \in \mathcal{S}^c} \right) \leq 4N^2 \exp \left(\frac{-t^2}{M^2 n^3 h^{3d_\zeta+4} + M n h^2 t} \right)$$

It follows that we can find a sufficiently large constant $\bar{D}_3 > 0$ such that when $t = \bar{D}_3 \sqrt{n^3 h^{3d_\zeta+4} \log N}$,

$$\mathbb{P} \left(\max_{i, j \in \mathcal{S}^c} \left| \sum_{i', j' \in \mathcal{S}_2} \left(g_{i'j'}^{(ij)} - \mathbb{E} \left(g_{i'j'}^{(ij)} \mid \{\zeta_i\}_{i \in \mathcal{S}^c} \right) \right) \right| \geq \bar{D}_3 \sqrt{n^3 h^{3d_\zeta+4} \log N} \mid \{\zeta_i\}_{i \in \mathcal{S}^c} \right) \leq \delta n^{-1/2}$$

We complete proving (B.28). ■

C Proofs for Empirical Applications

Proof of Theorem 3.

$$\begin{aligned} \sup_{\alpha} \left\| \psi(\hat{A}, Y_m, W_m, \alpha) - \mu(\alpha) \right\| &\leq \sup_{\alpha} \left\| \frac{1}{M} \sum_{m=1}^M \psi(\hat{A}, W_m, Y_m, \alpha) - \frac{1}{M} \sum_{m=1}^M \psi(P_m, W_m, Y_m, \alpha) \right\| \\ &\quad + \sup_{\alpha} \left\| \frac{1}{M} \sum_{m=1}^M \psi(P_m, W_m, Y_m, \alpha) - \frac{1}{M} \sum_{m=1}^M \psi(A_m, W_m, Y_m, \alpha) \right\| \\ &\quad + \sup_{\alpha} \left\| \frac{1}{M} \sum_{m=1}^M \psi(A_m, W_m, Y_m, \alpha) - \mathbb{E}(\psi(A_m, W_m, Y_m, \alpha)) \right\| \end{aligned}$$

Note that, by Assumption 5(vii) and Assumption 5(viii),

$$\begin{aligned} &\sup_{\alpha} \left\| \frac{1}{M} \sum_{m=1}^M \psi(\hat{A}, W_m, Y_m, \alpha) - \frac{1}{M} \sum_{m=1}^M \psi(P_m, W_m, Y_m, \alpha) \right\| \\ &\leq \frac{1}{M} \sum_{m=1}^M L(W_m, Y_m) N^{-1} \|\hat{F}_{m1} - F_{m2}\|_F \\ &= o_P(1) \end{aligned}$$

In addition, by Assumption 5(iv), we have

$$\sup_{\alpha} \left\| \frac{1}{M} \sum_{m=1}^M \psi(P_m, W_m, Y_m, \alpha) - \frac{1}{M} \sum_{m=1}^M \psi(A_m, W_m, Y_m, \alpha) \right\| = o_P(1)$$

By Assumption 5(vi), we have

$$\sup_{\alpha} \left\| \frac{1}{M} \sum_{m=1}^M \psi(A_m, W_m, Y_m, \alpha) - \mathbb{E}(\psi(A_m, W_m, Y_m, \alpha_0)) \right\| = o_P(1)$$

Thus, we must have $\sup_{\alpha} \left\| \psi(\hat{A}, W_m, Y_m, \alpha) - \mathbb{E}(\psi(A_m, W_m, Y_m, \alpha_0)) \right\| \xrightarrow{p} 0$. Combining the uniform convergence result, Assumption 5(iii), (v), (vi), and Theorem 2.1 in Newey and McFadden (1994), it follows that $\hat{\alpha} \xrightarrow{p} \alpha_0$. ■

Proof of Theorem 4. For any $m \in \{1, \dots, M\}$ and $i = 1, \dots, N_m$, consider the following expansion

of $(\hat{G}_m W_m - G_m W_m)_i$ up to second order¹⁵,

$$\begin{aligned}
(\hat{G}_m W_m - G_m W_m)_i &= \sum_{j \in \mathcal{S}} (\hat{G}_{m,ij} - G_{m,ij}) W_{m,j} = \frac{\sum_{j \in \mathcal{S}} A_{ij} W_{m,j} + \sum_{j \in \mathcal{S}} (\hat{A}_{m,ij} - A_{m,ij}) W_{m,j}}{\sum_{j \in \mathcal{S}} A_{m,ij} + \left(\sum_{j \in \mathcal{S}} \hat{A}_{m,ij} - \sum_{j \in \mathcal{S}} A_{m,ij} \right)} - \frac{\sum_{j \in \mathcal{S}} A_{m,ij} W_{m,j}}{\sum_{j \in \mathcal{S}} A_{m,ij}} \\
&\approx \frac{\sum_{j \in \mathcal{S}} (\hat{A}_{m,ij} - A_{m,ij}) W_{m,j}}{\sum_{j \in \mathcal{S}} A_{m,ij}} + \frac{\sum_{j \in \mathcal{S}} A_{m,ij} W_{m,j}}{\left(\sum_{j \in \mathcal{S}} A_{m,ij} \right)^2} \sum_{j \in \mathcal{S}} (\hat{A}_{m,ij} - A_{m,ij}) \\
&\quad + \frac{\sum_{j \in \mathcal{S}} A_{m,ij} W_{m,j}}{\left(\sum_{j \in \mathcal{S}} A_{m,ij} \right)^2} \left(\sum_{j \in \mathcal{S}} (\hat{A}_{m,ij} - A_{m,ij}) \right) \left(\sum_{j \in \mathcal{S}} (\hat{A}_{m,ij} - A_{m,ij}) W_{m,j} \right) \\
&\quad + \frac{\sum_{j \in \mathcal{S}} A_{m,ij} W_{m,j}}{\left(\sum_{j \in \mathcal{S}} A_{m,ij} \right)^3} \left(\sum_{j \in \mathcal{S}} (\hat{A}_{m,ij} - A_{m,ij}) \right)^2
\end{aligned}$$

Since $f_m(\cdot, \cdot) > 0$ and $\text{supp}(\zeta)$ is compact, we have $\sum_{j \in \mathcal{S}} A_{m,ij} \asymp N_m$ with probability at least $1 - \delta n_m^{-1/2}$. In addition, since W is uniformly bounded, $\sum_{j \in \mathcal{S}} A_{m,ij} W_{m,j} = O(N_m)$ with probability at least $1 - \delta n_m^{-1/2}$. Then, by (C.2), (C.3), and (C.4), we obtain that, with probability at least $1 - \delta n_m^{-1/2}$,

$$\begin{aligned}
\mathbb{E} \left((\hat{G}_m W_m - G_m W_m)_i \mid \{\zeta_{m,i}, W_{m,i}\}_{i=1}^{N_m} \right) &= O \left(h^2 + \frac{\log N_m}{n_m^2 h^{2d_\zeta}} \right) \\
\text{Var} \left((\hat{G}_m W_m - G_m W_m)_{m,i} \mid \{\zeta_{m,i}, W_{m,i}\}_{i=1}^{N_m} \right) &= O \left(\delta_{N,n}^2 h^2 + \frac{\log N_m}{n_m^2 h^{2d_\zeta}} \right)
\end{aligned}$$

One can also show that, using concentration inequalities in Lemma B.3 and Lemma B.4, $(\hat{G}_m W_m - G_m W_m)_i = O \left(h^2 + \frac{\log N_m}{n_m^2 h^{2d_\zeta}} \right)$ with probability at least $1 - \delta n_m^{-1/2}$. Also, it is straightforward to extend the result to $(\hat{G}_m Y_m - G_m Y_m)_i$. In addition, $(\hat{G}_m W_m)' \hat{G}_m W_m - (G_m W_m)' G_m W_m$ can be approximated by¹⁶

$$(\hat{G}_m W_m)' \hat{G}_m W_m - (G_m W_m)' G_m W_m \approx 2(G_m W_m)' (\hat{G}_m W_m - G_m W_m)$$

It follows that, with probability at least $1 - \delta n_m^{-1/2}$,

$$\begin{aligned}
\mathbb{E} \left((\hat{G}_m W_m)' \hat{G}_m W_m - (G_m W_m)' G_m W_m \mid \{\zeta_{m,i}, W_{m,i}\}_{i=1}^{N_m} \right) &= N_m O \left(h^2 + \frac{\log N_m}{n_m^2 h^{2d_\zeta}} \right) \\
\text{Var} \left((\hat{G}_m W_m)' \hat{G}_m W_m - (G_m W_m)' G_m W_m \mid \{\zeta_{m,i}, W_{m,i}\}_{i=1}^{N_m} \right) &= N_m^2 O \left(\delta_{N,n}^2 h^2 + \frac{\log N_m}{n_m^2 h^{2d_\zeta}} \right)
\end{aligned}$$

and $(\hat{G}_m W_m)' \hat{G}_m W_m - (G_m W_m)' G_m W_m = O \left(h^2 + \frac{\log N_m}{n_m^2 h^{2d_\zeta}} \right)$ with probability at least $1 - \delta n^{-1/2}$. Define

$$\begin{aligned}
\Delta_{m,1} &:= \hat{V}_m' \hat{Z}_m \Sigma_m \hat{Z}_m' \hat{V}_m - V_m' Z_m \Sigma_m Z_m' V_m \\
\Delta_{m,2} &:= \hat{V}_m' \hat{Z}_m \Sigma_m \hat{Z}_m' Y_m - V_m' Z_m \Sigma_m Z_m' Y_m
\end{aligned}$$

It is straightforward to extend the above results to show that, for $k \in \{1, 2\}$, with probability at least

¹⁵It is straightforward to verify that, by (C.2), (C.3), and (C.4), higher order expansion does not (asymptotically) contribute the bias and variance of $(\hat{G}_m W_m - G_m W_m)_i$.

¹⁶By the bias, the variance, and the probability bound of $(\hat{G}_m W_m - G_m W_m)_i$, higher order expansion does not (asymptotically) contribute the bias and variance.

$$1 - \delta n_m^{-1/2},$$

$$\begin{aligned}\mathbb{E}(\Delta_{m,k} \mid \{\zeta_{m,i}, W_{m,i}\}_{i=1}^{N_m}) &= N_m O\left(h^2 + \frac{\log N_m}{n_m^2 h^{2d_\zeta}}\right), \\ \text{Var}(\Delta_{m,k} \mid \{\zeta_{m,i}, W_{m,i}\}_{i=1}^{N_m}) &= N_m^2 O\left(\delta_{N,n}^2 h^2 + \frac{\log N_m}{n_m^2 h^{2d_\zeta}}\right), \\ \Delta_{m,k} &= O\left(h^2 + \frac{\log N_m}{n_m^2 h^{2d_\zeta}}\right),\end{aligned}\tag{C.1}$$

For notational simplicity, denote

$$\begin{aligned}\Delta_1 &:= \sum_{m=1}^M \hat{V}_m' \hat{Z}_m \Sigma_m \hat{Z}_m' \hat{V}_m - \sum_{m=1}^M V_m' Z_m \Sigma_m Z_m' V_m \\ \Delta_2 &:= \sum_{m=1}^M \hat{V}_m' \hat{Z}_m \Sigma_m \hat{Z}_m' Y_m - \sum_{m=1}^M V_m' Z_m \Sigma_m Z_m' Y_m,\end{aligned}$$

and $\hat{\alpha}$ can be approximated by¹⁷

$$\begin{aligned}\hat{\alpha} &= \left(\sum_{m=1}^M \hat{V}_m' \hat{Z}_m \Sigma_m \hat{Z}_m' \hat{V}_m \right)^{-1} \left(\sum_{m=1}^M \hat{V}_m' \hat{Z}_m \Sigma_m \hat{Z}_m' Y_m \right) \\ &= \left(\sum_{m=1}^M V_m' Z_m \Sigma_m Z_m' V_m + \Delta_1 \right)^{-1} \left(\sum_{m=1}^M V_m' Z_m \Sigma_m Z_m' Y_m + \Delta_2 \right) \\ &\approx \left(\sum_{m=1}^M V_m' Z_m \Sigma_m Z_m' V_m \right)^{-1} \left(I + \Delta_1 \left(\sum_{m=1}^M V_m' Z_m \Sigma_m Z_m' V_m \right)^{-1} \right) \left(\sum_{m=1}^M V_m' Z_m \Sigma_m Z_m' Y_m + \Delta_2 \right) \\ &\approx \underbrace{\left(\sum_{m=1}^M V_m' Z_m \Sigma_m Z_m' V_m \right)^{-1} \sum_{m=1}^M V_m' Z_m \Sigma_m Z_m' Y_m}_{U_1} + \underbrace{\left(\sum_{m=1}^M V_m' Z_m \Sigma_m Z_m' V_m \right)^{-1} \Delta_2}_{U_2} \\ &\quad + \underbrace{\left(\sum_{m=1}^M V_m' Z_m \Sigma_m Z_m' V_m \right)^{-1} \Delta_1 \left(\sum_{m=1}^M V_m' Z_m \Sigma_m Z_m' V_m \right)^{-1} \left(\sum_{m=1}^M V_m' Z_m \Sigma_m Z_m' Y_m \right)}_{U_3}\end{aligned}$$

Here, by CLT, $\frac{1}{\sqrt{M}}(U_1 - \alpha) \xrightarrow{d} N(0, \Omega)$, where Ω is the asymptotic variance when using complete network data. Let $N := \frac{1}{M} \sum_{m=1}^M N_m$ and $n := \frac{1}{M} \sum_{m=1}^M n_m$. By (C.1), Assumption 6(i), Assumption 6(iii), and Assumption 6(v), with probability at least $1 - \delta n^{-1/2}$, we obtain

$$\begin{aligned}\mathbb{E}(U_2 + U_3) &= O\left(h^2 + \frac{\log N_m}{n_m^2 h^{2d_\zeta}}\right) \\ \text{Var}(U_2 + U_3) &= \frac{1}{M} O\left(h^2 \delta_{N,n}^2 + \frac{\log N_m}{n_m^2 h^{2d_\zeta}}\right)\end{aligned}$$

Therefore, we complete the proof. ■

Lemma C.1 *Under conditions in Theorem 2 and Assumption 6, for any $i \in \mathcal{S}^c$, with probability at least $1 - \delta n^{-1/2}$,*

¹⁷It is straightforward to verify that, by the bias, the variance, and the probability bound of $\Delta_{m,k}$, higher order expansion does not (asymptotically) contribute the bias and variance of $\hat{\alpha}$.

$$\mathbb{E} \left(\sum_{j \in \mathcal{S}} (A_{ij} - \hat{A}_{ij}) \mid \{\zeta_i, W_i\}_{i=1}^N \right) = O(Nh^2), \quad \mathbb{E} \left(\sum_{j \in \mathcal{S}} (A_{ij} - \hat{A}_{ij}) W_j \mid \{\zeta_i, W_i\}_{i=1}^N \right) = O(Nh^2) \quad (\text{C.2})$$

and

$$\begin{aligned} \text{Var} \left(\sum_{j \in \mathcal{S}} (\hat{A}_{ij} - A_{ij}) \mid \{\zeta_i, W_i\}_{i=1}^N \right) &= O \left(N^2 \left(\delta_{N,n}^2 h^2 + \frac{\log N}{n^2 h^{2d_\zeta}} \right) \right), \\ \text{Var} \left(\sum_{j \in \mathcal{S}} (\hat{A}_{ij} - A_{ij}) W_j \mid \{\zeta_i, W_i\}_{i=1}^N \right) &= O \left(N^2 \left(\delta_{N,n}^2 h^2 + \frac{\log N}{n^2 h^{2d_\zeta}} \right) \right) \end{aligned} \quad (\text{C.3})$$

In addition, for any $\delta > 0$, there exist constant $E_1, E_2 \geq 0$, such that

$$\mathbb{P} \left(\left| \sum_{j \in \mathcal{S}} (\hat{A}_{ij} - A_{ij}) \right| \geq E_1 N \left(h^2 + \frac{\log N}{n h^{d_\zeta}} \right) \right) \leq \delta n^{-1/2} \quad (\text{C.4})$$

Proof of Lemma C.1. We only present the proof of (C.2) and (C.3), as (C.4) follows directly by using concentration inequalities in Lemma B.3 and Lemma B.4. For any $i, j \in \mathcal{S}^c$ and $i', j' \in \mathcal{S}^c$, define $Q_2^{(ij)}$ and $Q_3^{(ij)}$ as in (B.3), and $\hat{R}_3^{(ij)}$ as in (B.4), define $\hat{R}_1^{(ij)}$, $\hat{R}_2^{(ij)}$, and $\hat{R}_3^{(ij)}$ as in (B.4), define $R_1^{(ij)}$, $R_2^{(ij)}$, and $R_3^{(ij)}$ as in (B.10), and define $H_{i'}^{(ij)}$, $H_{j'}^{(ij)}$, and $H_{i'j'}^{(ij)}$ as in (B.5). Note that

$$\sum_{j \in \mathcal{S}} (\hat{A}_{ij} - A_{ij}) = \sum_{j \in \mathcal{S}^c} \sum_{k=1,2,3} R_k^{(ij)} + \sum_{j \in \mathcal{S}^c} \sum_{k=1,2,3} (\hat{R}_k^{(ij)} - R_k^{(ij)}) + \sum_{j \in \mathcal{S}^c} Q_2^{(ij)} + \sum_{j \in \mathcal{S}^c} Q_3^{(ij)} - \sum_{j \in \mathcal{S}^c} \epsilon_{ij}$$

Bias By (B.6), (B.7), Assumption 1(ii), and Assumption 6(ii), we have, with probability at least $1 - \delta n^{-1/2}$,

$$\mathbb{E} \left(\sum_{j \in \mathcal{S}^c} \sum_{k=1,2,3} \hat{R}_k^{(ij)} \mid \{\zeta_i, W_i\}_{i=1}^N \right) \leq 6N \rho_{f''} \eta^2 h^2$$

and

$$\mathbb{E} \left(\sum_{j \in \mathcal{S}^c} Q_2^{(ij)} - \sum_{j \in \mathcal{S}^c} \epsilon_{ij} \mid \{\zeta_i, W_i\}_{i=1}^N \right) = 0$$

In addition, with probability at least $1 - \delta n^{-1/2}$,

$$\begin{aligned} \sum_{j \in \mathcal{S}^c} Q_3^{(ij)} &= \sum_{j \in \mathcal{S}^c} \frac{\sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'i'}^{(ij)} f_{i'i'}}{\sum_{i', j' \in \mathcal{S}_2} \hat{K}_{h,i'j'}^{(ij)}} \stackrel{(i)}{\leq} \frac{1}{D_2 n^2 h^{2d_\zeta}} \sum_{j \in \mathcal{S}^c} \left| \sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'i'}^{(ij)} f_{i'i'} \right| \\ &\leq \frac{1}{D_2 n^2 h^{2d_\zeta}} \left| \sum_{i' \in \mathcal{S}_2} \sum_{j \in \mathcal{S}^c} \hat{K}_{h,i'i'}^{(ij)} f_{i'i'} \right| \\ &\leq \frac{\rho_K^2 \rho_f}{D_2 n^2 h^{2d_\zeta}} \left| \sum_{i' \in \mathcal{S}_2} \mathbf{1}(d_{ii'} \leq h) \sum_{j' \in \mathcal{S}^c} \mathbf{1}(d_{ij} \leq h) \right| \\ &\stackrel{(ii)}{\leq} \frac{\rho_K^2 \rho_f}{D_2 n^2 h^{2d_\zeta}} \bar{C}_1^2 n N h^{2d_\zeta} \\ &= \frac{\bar{C}_1^2 \rho_K^2 \rho_f N}{D_2 n} \end{aligned} \quad (\text{C.5})$$

Here, (i) holds with probability at least $1 - \delta n^{-1/2}$ because of (B.27) in Lemma B.26, (ii) follows from (B.19) by replacing $K(\cdot)$ with a uniform kernel. Therefore, we conclude that, with probability

at least $1 - \delta n^{-1/2}$,

$$\mathbb{E} \left(\sum_{j \in \mathcal{S}} (\hat{A}_{ij} - f_{ij}) \mid \{\zeta_i, W_i\}_{i=1}^N \right) \leq O \left(Nh^2 + \frac{N}{n} \right) = O(Nh^2)$$

The last equality holds because of conditions in Theorem 2 and $d_\zeta \geq 2$. Since W_i is uniformly bounded (Assumption 6(iv)), it immediately follows that, with probability at least $1 - \delta n^{-1/2}$,

$$\mathbb{E} \left(\sum_{j \in \mathcal{S}} (\hat{A}_{ij} - f_{ij}) W_j \mid \{\zeta_i, W_i\}_{i=1}^N \right) = O(Nh^2)$$

Variance Note that

$$\begin{aligned} & \text{Var} \left(\sum_{j \in \mathcal{S}} (\hat{A}_{ij} - A_{ij}) \mid \{\zeta_i, W_i\}_{i=1}^N \right) \\ &= 4 \text{Var} \left(\sum_{j \in \mathcal{S}^c} \sum_{k=1,2,3} (\hat{R}_k^{(ij)} - R_k^{(ij)}) \mid \{\zeta_i, W_i\}_{i=1}^N \right) + 4 \text{Var} \left(\sum_{j \in \mathcal{S}^c} Q_2^{(ij)} \mid \{\zeta_i, W_i\}_{i=1}^N \right) \\ & \quad + 4 \text{Var} \left(\sum_{j \in \mathcal{S}^c} Q_3^{(ij)} \mid \{\zeta_i, W_i\}_{i=1}^N \right) + 4 \text{Var} \left(\sum_{j \in \mathcal{S}^c} \epsilon_{ij} \mid \{\zeta_i, W_i\}_{i=1}^N \right) \end{aligned}$$

Step 1: Bound on $\text{Var} \left(\sum_{k=1,2,3} (\hat{R}_k^{(ij)} - R_k^{(ij)}) \mid \{\zeta_i, W_i\}_{i=1}^N \right)$ By (B.12), we have, with probability at least $1 - \delta n^{-1/2}$,

$$\begin{aligned} \text{Var} \left(\sum_{j \in \mathcal{S}^c} \hat{R}_1^{(ij)} - R_1^{(ij)} \mid \{\zeta_i, W_i\}_{i=1}^N \right) &\leq N^2 \max_{i,j \in \mathcal{S}^c} \text{Var} \left(\hat{R}_1^{(ij)} - R_1^{(ij)} \mid \{\zeta_i, W_i\}_{i=1}^N \right) = O(N^2 \delta_{N,n}^2 h^2) \\ \text{Var} \left(\sum_{j \in \mathcal{S}^c} \hat{R}_2^{(ij)} - R_2^{(ij)} \mid \{\zeta_i, W_i\}_{i=1}^N \right) &\leq N^2 \max_{i,j \in \mathcal{S}^c} \text{Var} \left(\hat{R}_2^{(ij)} - R_2^{(ij)} \mid \{\zeta_i, W_i\}_{i=1}^N \right) = O(N^2 \delta_{N,n}^2 h^2) \end{aligned}$$

Using the same argument, we obtain that, with probability at least $1 - \delta n^{-1/2}$,

$$\text{Var} \left(\sum_{j \in \mathcal{S}^c} \hat{R}_3^{(ij)} - R_3^{(ij)} \mid \{\zeta_i, W_i\}_{i=1}^N \right) \leq N^2 \max_{i,j \in \mathcal{S}^c} \text{Var} \left(\hat{R}_3^{(ij)} - R_3^{(ij)} \mid \{\zeta_i, W_i\}_{i=1}^N \right) = O(N^2 \delta_{N,n}^2 h^2)$$

Thus, with probability at least $1 - \delta n^{-1/2}$,

$$\text{Var} \left(\sum_{k=1,2,3} (\hat{R}_k^{(ij)} - R_k^{(ij)}) \mid \{\zeta_i, W_i\}_{i=1}^N \right) = O(N^2 \delta_{N,n}^2 h^2)$$

Step 2: Bound on $\text{Var} \left(\sum_{j \in \mathcal{S}^c} Q_2^{(ij)} \mid \{\zeta_i, W_i\}_{i=1}^N \right)$ Note that

$$\text{Var} \left(\sum_{j \in \mathcal{S}^c} \frac{\sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'}^{(i)} \epsilon_{i'j}}{\sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'}^{(i)}} \mid \{\zeta_i, W_i\}_{i=1}^N \right) = \text{Var} \left(\frac{\sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'}^{(i)} \sum_{j \in \mathcal{S}^c} \epsilon_{i'j}}{\sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'}^{(i)}} \mid \{\zeta_i, W_i\}_{i=1}^N \right)$$

Since $\sum_{j \in \mathcal{S}^c} \epsilon_{i'j}$ is independent across $i' \in \mathcal{S}_2$ and sub-Gaussian with $\sigma \sim \sqrt{n}$, we employ the method used in proving (B.14) to show that, with probability at least $1 - \delta n^{-1/2}$,

$$\text{Var} \left(\sum_{j \in \mathcal{S}^c} \frac{\sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'}^{(i)} \epsilon_{i'j}}{\sum_{i' \in \mathcal{S}_2} \hat{K}_{h,i'}^{(i)}} \mid \{\zeta_i, W_i\}_{i=1}^N \right) = O \left(\frac{N \log N}{n h^{d_\zeta}} \right)$$

Also, with probability at least $1 - \delta n^{-1/2}$,

$$\begin{aligned} \text{Var} \left(\sum_{j \in \mathcal{S}^c} \frac{\sum_{j' \in \mathcal{S}_2} \hat{K}_{h,j'}^{(j)} \epsilon_{ij'}}{\sum_{j' \in \mathcal{S}_2} \hat{K}_{h,j'}^{(j)}} \mid \{\zeta_i, W_i\}_{i=1}^N \right) &= \text{Var} \left(\frac{\sum_{j' \in \mathcal{S}_2} \left(\sum_{j \in \mathcal{S}^c} \hat{K}_{h,j'}^{(j)} \right) \epsilon_{ij'}}{\sum_{j' \in \mathcal{S}_2} \hat{K}_{h,j'}^{(j)}} \mid \{\zeta_i, W_i\}_{i=1}^N \right) \\ &\leq \frac{\max_{j' \in \mathcal{S}_2} \left(\sum_{j \in \mathcal{S}^c} \hat{K}_{h,j'}^{(j)} \right)^2}{\min_{j' \in \mathcal{S}_2} \left(\sum_{j \in \mathcal{S}^c} \hat{K}_{h,j'}^{(j)} \right)^2} \mathbb{E} \left(\left(\sum_{j' \in \mathcal{S}_2} \epsilon_{ij'} \right)^2 \right) \\ &\stackrel{(i)}{\leq} O \left(\frac{N^2 h^{2d_\zeta}}{n^2 h^{2d_\zeta}} n \log N \right) \\ &= O \left(\frac{N^2}{n} \log N \right) \end{aligned}$$

In addition, with probability at least $1 - \delta n^{-1/2}$,

$$\begin{aligned} \sum_{j \in \mathcal{S}^c} \text{Var} \left(\sum_{j' \in \mathcal{S}^c} \frac{\sum_{i', j'' \in \mathcal{S}_2} \hat{K}_{h,i'j''}^{(ij)} \epsilon_{i'j''}}{\sum_{i', j'' \in \mathcal{S}_2} \hat{K}_{h,i'j''}^{(ij)}} \mid \{\zeta_i, W_i\}_{i=1}^N \right) &\leq N^2 \max_{i, j \in \mathcal{S}^c} \text{Var} \left(\frac{\sum_{i', j'' \in \mathcal{S}_2} \hat{K}_{h,i'j''}^{(ij)} \epsilon_{i'j''}}{\sum_{i', j'' \in \mathcal{S}_2} \hat{K}_{h,i'j''}^{(ij)}} \mid \{\zeta_i, W_i\}_{i=1}^N \right) \\ &\stackrel{(i)}{=} O \left(N^2 \frac{\log N}{n^2 h^{2d_\zeta}} \right) \end{aligned}$$

where (i) follows from (B.15).

Therefore, we conclude that, when $d_\zeta \geq 2$, with probability at least $1 - \delta n^{-1/2}$,

$$\begin{aligned} \text{Var} \left(\sum_{j \in \mathcal{S}^c} Q_2^{(ij)} \mid \{\zeta_i, W_i\}_{i=1}^N \right) &= O \left(\frac{N^2 \log N}{N n h^{d_\zeta}} + \frac{N^2}{n} \log N + N^2 \frac{\log N}{n^2 h^{2d_\zeta}} \right) \\ &= O \left(N^2 \frac{\log N}{n^2 h^{2d_\zeta}} \right) \end{aligned}$$

Step 3: Bound on $\text{Var} \left(\sum_{j \in \mathcal{S}^c} Q_3^{(ij)} \mid \{\zeta_i, W_i\}_{i=1}^N \right)$ By (C.5), with probability at least $1 - \delta n^{-1/2}$,

$$\text{Var} \left(\sum_{j \in \mathcal{S}^c} Q_3^{(ij)} \mid \{\zeta_i, W_i\}_{i=1}^N \right) = O \left(\frac{N^2}{n^2} \right)$$

Step 4: Bound on $\text{Var} \left(\sum_{j \in \mathcal{S}^c} \epsilon_{ij} \mid \{\zeta_i, W_i\}_{i=1}^N \right)$ Since $\{\epsilon_{ij} \mid i < j, i, j \in \mathcal{S}_2 \cup \mathcal{S}^c\}$, it is straightforward to verify that, with probability at least $1 - \delta n^{-1/2}$,

$$\text{Var} \left(\sum_{j \in \mathcal{S}^c} \epsilon_{ij} \mid \{\zeta_i, W_i\}_{i=1}^N \right) = O(N \log N)$$

Combining these results above, we conclude that, with probability at least $1 - \delta n^{-1/2}$,

$$\begin{aligned}\text{Var} \left(\sum_{j \in \mathcal{S}} (\hat{A}_{ij} - A_{ij}) \mid \{\zeta_i, W_i\}_{i=1}^N \right) &= O \left(N^2 \left(\delta_{N,n}^2 h^2 + \frac{\log N}{n^2 h^{2d_\zeta}} + \frac{1}{n^2} + \frac{1}{N} \right) \right) \\ &= O \left(N^2 \left(\delta_{N,n}^2 h^2 + \frac{\log N}{n^2 h^{2d_\zeta}} \right) \right)\end{aligned}$$

Since W_i is uniformly bounded (Assumption 6(iv)), it immediately follows that, with probability at least $1 - \delta n^{-1/2}$,

$$\text{Var} \left(\sum_{j \in \mathcal{S}} (\hat{A}_{ij} - A_{ij}) W_j \mid \{\zeta_i, W_i\}_{i=1}^N \right) = O \left(N^2 \left(\delta_{N,n}^2 h^2 + \frac{\log N}{n^2 h^{2d_\zeta}} \right) \right)$$

This completes the proof.

■

D Supplementary Proofs

Lemma D.1 Let $\zeta_i \in \{\bar{\zeta}_1, \dots, \bar{\zeta}_G\}$ be i.i.d. random variables such that $0 \leq \bar{\zeta}_g \leq 1$ for any $g = 1, 2, \dots, G$, then $f(\cdot, \cdot)$ satisfies Assumption 2 except in the degenerate case where there exist $g \neq g'$ such that $f(\bar{\zeta}_g, \zeta) = f(\bar{\zeta}_{g'}, \zeta)$ for all $\zeta \in \{\bar{\zeta}_1, \dots, \bar{\zeta}_G\}$.

Proof of Lemma D.1. Without loss of generality, suppose that $f(\bar{\zeta}_1, \zeta_{g'}) > 0$ for all g' , then for any $g, g' \in \{1, \dots, G\}$,

$$\begin{aligned} & \max_{\zeta \in \{\bar{\zeta}_1, \dots, \bar{\zeta}_G\}} \left| \int f(\zeta, \zeta') (f(\bar{\zeta}_g, \zeta') - f(\bar{\zeta}_{g'}, \zeta')) d\mathbb{P}(\zeta') \right| \\ & \geq \max \left\{ \left| \int f(\bar{\zeta}_g, \zeta') (f(\bar{\zeta}_g, \zeta') - f(\bar{\zeta}_{g'}, \zeta')) d\mathbb{P}(\zeta') \right|, \left| \int f(\bar{\zeta}_{g'}, \zeta') (f(\bar{\zeta}_g, \zeta') - f(\bar{\zeta}_{g'}, \zeta')) d\mathbb{P}(\zeta') \right| \right\} \end{aligned}$$

Then expression above equals 0 if and only if

$$\int f^2(\bar{\zeta}_g, \zeta) d\mathbb{P}(\zeta) = \int f^2(\bar{\zeta}_{g'}, \zeta) d\mathbb{P}(\zeta) = \int f(\bar{\zeta}_g, \zeta) f(\bar{\zeta}_{g'}, \zeta) d\mathbb{P}(\zeta)$$

which implies that $f(\bar{\zeta}_g, \zeta) = f(\bar{\zeta}_{g'}, \zeta)$ for all $\zeta \in \{\bar{\zeta}_1, \dots, \bar{\zeta}_G\}$. ■

Lemma D.2 Let $\zeta_i \in [0, 1]^{d_\zeta}$ be i.i.d. random vectors such that the components of $\zeta_i = (\zeta_{i1}, \dots, \zeta_{id_\zeta})'$ are mutually independent and each follows the uniform distribution on $[0, 1]$, then

$$f(\zeta_i, \zeta_j) = \frac{\exp(\omega(\zeta_i, \zeta_j)' \beta)}{1 + \exp(\omega(\zeta_i, \zeta_j)' \beta)}$$

satisfies Assumption 2 when

$$\omega(\zeta_i, \zeta_j)' \beta = \zeta_{i1} + \zeta_{j1} + \sum_{k=2}^d \beta_k |\zeta_{ik} - \zeta_{jk}| + \sum_{k=d+1}^{d_\zeta} \beta_k (\zeta_{ik} - \zeta_{jk})^2, \quad \beta_2, \dots, \beta_k \leq 0$$

Proof of Lemma D.2. Since $\zeta_i \in [0, 1]^{d_\zeta}$ be i.i.d. random vectors such that the components of $\zeta_i = (\zeta_{i1}, \dots, \zeta_{id_\zeta})'$ are mutually independent and each follows the uniform distribution on $[0, 1]$, it is sufficiently to check the Assumption 2 under the following three specific case.

Case 1: $d_\zeta = 1$ and $\omega(\zeta_i, \zeta_j) = \zeta_i + \zeta_j$ In this case, for each $\zeta_1 < \zeta_2 \in [0, 1]$ we have

$$f(\zeta_1, \zeta_2) = \frac{\exp(\zeta_1 + \zeta_2)}{1 + \exp(\zeta_1 + \zeta_2)}$$

and

$$\begin{aligned} d_{12} &= \sup_{\zeta} \left| \int f(\zeta, \zeta') (f(\zeta_1, \zeta') - f(\zeta_2, \zeta')) d\zeta' \right| \\ &\geq \left| \int f(0, \zeta') (f(\zeta_1, \zeta') - f(\zeta_2, \zeta')) d\zeta' \right| \\ &\geq \left(\inf_{\zeta \in [0, 1]} f(0, \zeta) \right) \left| \int_0^1 \int_{\zeta_1}^{\zeta_2} f'(\zeta, \zeta') d\zeta d\zeta' \right| \\ &\stackrel{(i)}{\geq} \left(\inf_{\zeta \in [0, 1]} f(0, \zeta) \right) \left(\inf_{\zeta, \zeta' \in [0, 1]} f'(\zeta, \zeta') \right) |\zeta_2 - \zeta_1| \end{aligned}$$

where inequality (i) holds because $f'(\cdot, \cdot) > 0$ on $[0, 1]^2$. It is straightforward to check that $(\inf_{\zeta \in [0, 1]} f(0, \zeta)) > 0$ and $(\inf_{\zeta, \zeta' \in [0, 1]} f'(\zeta, \zeta')) > 0$, thus, let $\lambda = (\inf_{\zeta \in [0, 1]} f(0, \zeta)) (\inf_{\zeta, \zeta' \in [0, 1]} f'(\zeta, \zeta'))$, we prove that

$$d_{12} \geq \lambda |\zeta_2 - \zeta_1|$$

Case 2: $d_\zeta = 1$ and $\omega(\zeta_i, \zeta_j) = -|\zeta_i - \zeta_j|$ In this case, for each $\zeta_1 < \zeta_2 \in [0, 1]$ we have

$$f(\zeta_1, \zeta_2) = \frac{\exp(-|\zeta_1 - \zeta_2|)}{1 + \exp(-|\zeta_1 - \zeta_2|)}$$

and define

$$h_0(\zeta) = \int f(0, \zeta') f(\zeta, \zeta') d\zeta', \quad h_1(\zeta) = \int f(1, \zeta') f(\zeta, \zeta') d\zeta'$$

and it follows that

$$d_{12} = \sup_{\zeta} \left| \int f(\zeta, \zeta') (f(\zeta_1, \zeta') - f(\zeta_2, \zeta')) d\zeta' \right| \geq \max \{ |h_0(\zeta_1) - h_0(\zeta_2)|, |h_1(\zeta_1) - h_1(\zeta_2)| \}$$

Thus, it suffices to prove that there exists a constant $\lambda > 0$ such that for any $\zeta_1, \zeta_2 \in [0, 1]$:

$$\max \{ |h_0(\zeta_1) - h_0(\zeta_2)|, |h_1(\zeta_1) - h_1(\zeta_2)| \} \geq \lambda |\zeta_1 - \zeta_2|$$

It is straightforward to check that

$$\begin{aligned} h'_0(\zeta) &= - \int f(0, \zeta') \text{sign}(\zeta - \zeta') f(\zeta, \zeta') (1 - f(\zeta, \zeta')) d\zeta' \\ h_0(\zeta) &= h_1(1 - \zeta), \quad h'_1(\zeta) = -h'_0(1 - \zeta) \end{aligned}$$

Also, $h_0(\cdot)$ achieves its maximum at $\zeta_0^* < 1/2$ and $h_1(\cdot)$ achieves its maximum at $\zeta_1^* = 1 - \zeta_0^* > 1/2$. $h_0(\cdot)$ is a strictly increasing function ζ when $\zeta < \zeta_0^*$, and is a strictly decreasing function ζ when $\zeta > \zeta_0^*$.

Step 1 $\zeta_1 \in [\zeta_0^*, \zeta_1^*] \vee \zeta_2 \in [\zeta_0^*, \zeta_1^*]$ Fix an arbitrary positive constant $\delta < 1/2 - \zeta_0^*$, define

$$\lambda_0 = \inf_{\zeta \in [\zeta_0^* + \delta, 1]} |h'_0(\zeta)|$$

One can check that $\lambda_0 > 0$ because $h'_0(\cdot)$ is strictly negative on $[\zeta_0^* + \delta, 1]$. Then:

- (i) When $\zeta_1 \in [0, \zeta_0^*]$ and $\zeta_2 \in [\zeta_0^*, \zeta_1^* - \delta_0]$, we have $|h_1(\zeta_1) - h_1(\zeta_2)| \geq \lambda_0 |\zeta_1 - \zeta_2|$ by mean value theorem.
- (ii) When $\zeta_1 \in [0, \zeta_0^*]$ and $\zeta_2 \in [\zeta_1^* - \delta_0, \zeta_1^*]$, $|h_1(\zeta_1) - h_1(\zeta_2)| \geq \lambda_0 (\zeta_1^* - \zeta_2^* - \delta)$ and thus, $|h_1(\zeta_1) - h_1(\zeta_2)| \geq \lambda_0 (\zeta_1^* - \zeta_2^* - \delta) |\zeta_1 - \zeta_2|$ because $|\zeta_1 - \zeta_2| \leq 1$.
- (iii) When $\zeta_1, \zeta_2 \in [\zeta_0^*, \zeta_1^*]$, since $\delta < 1/2 - \zeta_0^*$, it follows that

$$\max \{ |h_0(\zeta_1) - h_0(\zeta_2)|, |h_1(\zeta_1) - h_1(\zeta_2)| \} \geq \lambda_0 |\zeta_1 - \zeta_2|$$

- (iv) When $\zeta_1 \in [\zeta_0^*, \zeta_0^* + \delta]$ and $\zeta_2 \in [\zeta_1^*, 1]$, $|h_0(\zeta_1) - h_0(\zeta_2)| \geq \lambda_0 (\zeta_1^* - \zeta_2^* - \delta)$ and thus, $|h_0(\zeta_1) - h_0(\zeta_2)| \geq \lambda_0 (\zeta_1^* - \zeta_2^* - \delta) |\zeta_1 - \zeta_2|$ because $|\zeta_1 - \zeta_2| \leq 1$.
- (v) When $\zeta_1 \in [0, \zeta_0^*]$ and $\zeta_2 \in [\zeta_0^*, \zeta_1^* - \delta_0]$, we have $|h_0(\zeta_1) - h_0(\zeta_2)| \geq \lambda_0 |\zeta_1 - \zeta_2|$ by mean value theorem.

Therefore, we can show that

$$\max\{|h_0(\zeta_1) - h_0(\zeta_2)|, |h_1(\zeta_1) - h_1(\zeta_2)|\} \geq \lambda_0(\zeta_1^* - \zeta_0^* - \delta)|\zeta_1 - \zeta_2|$$

Step 2 $\zeta_1 \in [0, \zeta_0^*] \wedge \zeta_2 \in [\zeta_1^*, 1]$ Define

$$\lambda_1 = \inf_{\zeta \in [0, \zeta_0^*]} |h_0(\zeta) - h_1(\zeta)|$$

We have $\lambda_1 > 0$ because $h_0(\zeta) > h_1(\zeta)$ when $\zeta \in [0, \zeta_0^*]$. Thus, for any $\zeta_1 \in [0, \zeta_0^*] \wedge \zeta_2 \in [\zeta_1^*, 1]$, we have

(i) When $|h_0(\zeta_1) - h_0(\zeta_2)| \leq \lambda_1$, we have $|h_1(\zeta_1) - h_1(\zeta_2)| \geq 2\lambda_1 - \lambda_1 = \lambda_1$.

(ii) When $|h_1(\zeta_1) - h_1(\zeta_2)| \leq \lambda_1$, we have $|h_0(\zeta_1) - h_0(\zeta_2)| \geq 2\lambda_1 - \lambda_1 = \lambda_1$.

Therefore, we have $\max\{|h_0(\zeta_1) - h_0(\zeta_2)|, |h_1(\zeta_1) - h_1(\zeta_2)|\} \geq \lambda_1 \geq \lambda_1|\zeta_1 - \zeta_2|$ when $\zeta_1 \in [0, \zeta_0^*] \wedge \zeta_2 \in [\zeta_1^*, 1]$.

Combining the conclusion in two steps yields that

$$d_{12} \geq \max\{|h_0(\zeta_1) - h_0(\zeta_2)|, |h_1(\zeta_1) - h_1(\zeta_2)|\} \geq \min\{\lambda_0(\zeta_1^* - \zeta_0^* - \delta), \lambda_1\}|\zeta_2 - \zeta_1|$$

Case 3: $d_\zeta = 1$ and $\omega(\zeta_i, \zeta_j) = -(\zeta_i - \zeta_j)^2$ In this case, for each $\zeta_1 < \zeta_2 \in [0, 1]$ we have

$$f(\zeta_1, \zeta_2) = \frac{\exp(-(\zeta_1 - \zeta_2)^2)}{1 + \exp(-(\zeta_1 - \zeta_2)^2)}$$

and define

$$h_0(\zeta) = \int f(0, \zeta')f(\zeta, \zeta')d\zeta', \quad h_1(\zeta) = \int f(1, \zeta')f(\zeta, \zeta')d\zeta'$$

and it follows that

$$d_{12} = \sup_{\zeta} \left| \int f(\zeta, \zeta')(f(\zeta_1, \zeta') - f(\zeta_2, \zeta'))d\zeta' \right| \geq \max\{|h_0(\zeta_1) - h_0(\zeta_2)|, |h_1(\zeta_1) - h_1(\zeta_2)|\}$$

Thus, it suffices to prove that there exists a constant $\lambda > 0$ such that for any $\zeta_1, \zeta_2 \in [0, 1]$:

$$\max\{|h_0(\zeta_1) - h_0(\zeta_2)|, |h_1(\zeta_1) - h_1(\zeta_2)|\} \geq \lambda|\zeta_1 - \zeta_2|$$

It is straightforward to check that

$$\begin{aligned} h'_0(\zeta) &= -2 \int f(0, \zeta')(\zeta - \zeta')f(\zeta, \zeta')(1 - f(\zeta, \zeta'))d\zeta' \\ h_0(\zeta) &= h_1(1 - \zeta), \quad h'_1(\zeta) = -h'_0(1 - \zeta) \end{aligned}$$

Also, $h_0(\cdot)$ achieves its maximum at $\zeta_0^* < 1/2$ and $h_1(\cdot)$ achieves its maximum at $\zeta_1^* = 1 - \zeta_0^* > 1/2$. $h_0(\cdot)$ is a strictly increasing function ζ when $\zeta < \zeta_0^*$, and is a strictly decreasing function ζ when $\zeta > \zeta_0^*$.

Step 1 $\zeta_1 \in [\zeta_0^*, \zeta_1^*] \vee \zeta_2 \in [\zeta_0^*, \zeta_1^*]$ Fix an arbitrary positive constant $\delta < 1/2 - \zeta_0^*$, define

$$\lambda_0 = \inf_{\zeta \in [\zeta_0^* + \delta, 1]} |h'_0(\zeta)|$$

One can check that $\lambda_0 > 0$ because $h'_0(\cdot)$ is strictly negative on $[\zeta_0^* + \delta, 1]$. Then:

- (i) When $\zeta_1 \in [0, \zeta_0^*]$ and $\zeta_2 \in [\zeta_0^*, \zeta_1^* - \delta_0]$, we have $|h_1(\zeta_1) - h_1(\zeta_2)| \geq \lambda_0 |\zeta_1 - \zeta_2|$ by mean value theorem.
- (ii) When $\zeta_1 \in [0, \zeta_0^*]$ and $\zeta_2 \in [\zeta_1^* - \delta_0, \zeta_1^*]$, $|h_1(\zeta_1) - h_1(\zeta_2)| \geq \lambda_0(\zeta_1^* - \zeta_2^* - \delta)$ and thus, $|h_1(\zeta_1) - h_1(\zeta_2)| \geq \lambda_0(\zeta_1^* - \zeta_2^* - \delta) |\zeta_1 - \zeta_2|$ because $|\zeta_1 - \zeta_2| \leq 1$.
- (iii) When $\zeta_1, \zeta_2 \in [\zeta_0^*, \zeta_1^*]$, since $\delta < 1/2 - \zeta_0^*$, it follows that

$$\max\{|h_0(\zeta_1) - h_0(\zeta_2)|, |h_1(\zeta_1) - h_1(\zeta_2)|\} \geq \lambda_0 |\zeta_1 - \zeta_2|$$

- (iv) When $\zeta_1 \in [\zeta_0^*, \zeta_0^* + \delta]$ and $\zeta_2 \in [\zeta_1^*, 1]$, $|h_0(\zeta_1) - h_0(\zeta_2)| \geq \lambda_0(\zeta_1^* - \zeta_2^* - \delta)$ and thus, $|h_0(\zeta_1) - h_0(\zeta_2)| \geq \lambda_0(\zeta_1^* - \zeta_2^* - \delta) |\zeta_1 - \zeta_2|$ because $|\zeta_1 - \zeta_2| \leq 1$.
- (v) When $\zeta_1 \in [0, \zeta_0^*]$ and $\zeta_2 \in [\zeta_0^*, \zeta_1^* - \delta_0]$, we have $|h_0(\zeta_1) - h_0(\zeta_2)| \geq \lambda_0 |\zeta_1 - \zeta_2|$ by mean value theorem.

Therefore, we can show that

$$\max\{|h_0(\zeta_1) - h_0(\zeta_2)|, |h_1(\zeta_1) - h_1(\zeta_2)|\} \geq \lambda_0(\zeta_1^* - \zeta_0^* - \delta) |\zeta_1 - \zeta_2|$$

Step 2 $\zeta_1 \in [0, \zeta_0^*] \wedge \zeta_2 \in [\zeta_1^*, 1]$ Define

$$\lambda_1 = \inf_{\zeta \in [0, \zeta_0^*]} |h_0(\zeta) - h_1(\zeta)|$$

We have $\lambda_1 > 0$ because $h_0(\zeta) > h_1(\zeta)$ when $\zeta \in [0, \zeta_0^*]$. Thus, for any $\zeta_1 \in [0, \zeta_0^*] \wedge \zeta_2 \in [\zeta_1^*, 1]$, we have

- (i) When $|h_0(\zeta_1) - h_0(\zeta_2)| \leq \lambda_1$, we have $|h_1(\zeta_1) - h_1(\zeta_2)| \geq 2\lambda_1 - \lambda_1 = \lambda_1$.
- (ii) When $|h_1(\zeta_1) - h_1(\zeta_2)| \leq \lambda_1$, we have $|h_0(\zeta_1) - h_0(\zeta_2)| \geq 2\lambda_1 - \lambda_1 = \lambda_1$.

Therefore, we have $\max\{|h_0(\zeta_1) - h_0(\zeta_2)|, |h_1(\zeta_1) - h_1(\zeta_2)|\} \geq \lambda_1 \geq \lambda_1 |\zeta_1 - \zeta_2|$ when $\zeta_1 \in [0, \zeta_0^*] \wedge \zeta_2 \in [\zeta_1^*, 1]$.

Combining the conclusion in two steps yields that

$$d_{12} \geq \max\{|h_0(\zeta_1) - h_0(\zeta_2)|, |h_1(\zeta_1) - h_1(\zeta_2)|\} \geq \min\{\lambda_0(\zeta_1^* - \zeta_0^* - \delta), \lambda_1\} |\zeta_2 - \zeta_1|$$

■

Lemma D.3 *There are two sets of independent sub-Gaussian random variables $\{X_i\}_{i=1}^N$ and $\{Y_i\}_{i=1}^N$, such that there exist $\sigma_X, \sigma_Y > 0$:*

$$\sup_{1 \leq i \leq N} \mathbb{E}(e^{\lambda X_i}) \leq e^{\frac{\lambda^2 \sigma_X^2}{2}}, \quad \sup_{1 \leq i \leq N} \mathbb{E}(e^{\lambda Y_i}) \leq e^{\frac{\lambda^2 \sigma_Y^2}{2}}, \quad \forall \lambda \in \mathbb{R}$$

$\{X_i\}_{i=1}^N$ and $\{Y_i\}_{i=1}^N$ are independent of each other. Then for all $\epsilon > 0$,

$$\mathbb{P}\left(\left|\frac{1}{N} \sum_{i=1}^N X_i Y_i\right| \geq \epsilon\right) \leq 2 \exp\left(-\frac{N \epsilon^2}{2(\sigma_X^2 \sigma_Y^2 + \sigma_X \sigma_Y \epsilon)}\right)$$

Proof. It is no harm to assume $\mathbb{E}(X_i) = \mathbb{E}(Y_i) = 0$ for all i . Then

$$\mathbb{E} \exp(\lambda X_i Y_i) = \mathbb{E}(\mathbb{E}(\exp(\lambda X_i Y_i) \mid X_i)) \leq \mathbb{E} \exp\left(\frac{\lambda^2 X_i^2 \sigma_Y^2}{2}\right) \leq \frac{1}{\sqrt{1 - \lambda^2 \sigma_Y^2 \sigma_X^2}}, \quad \forall 0 \leq \lambda^2 \leq \frac{1}{\sigma_Y^2 \sigma_X^2}$$

When $\lambda^2 < \frac{1}{\sigma_Y^2 \sigma_X^2}$, we have

$$\frac{1}{\sqrt{1 - \lambda^2 \sigma_X^2 \sigma_Y^2}} \leq \exp \left(\frac{\lambda^2 \sigma_X^2 \sigma_Y^2}{2(1 - \lambda^2 \sigma_X^2 \sigma_Y^2)} \right) \leq \exp \left(\frac{\lambda^2 \sigma_X^2 \sigma_Y^2}{2(1 - \lambda \sigma_X \sigma_Y)} \right)$$

Therefore, $X_i Y_i$ is satisfies Bernstein's condition because

$$\mathbb{E}(\lambda X_i Y_i) \leq \exp \left(\frac{\lambda^2 \sigma_X^2 \sigma_Y^2}{2(1 - \lambda \sigma_X \sigma_Y)} \right), \quad \forall |\lambda| < \frac{1}{\sigma_X \sigma_Y}$$

It immediately follows that

$$\mathbb{E} \left(\frac{1}{N} \sum_{i=1}^N \lambda X_i Y_i \right) \leq \exp \left(\frac{\lambda^2 \sigma_X^2 \sigma_Y^2}{2N(1 - \lambda \sigma_X \sigma_Y)} \right), \quad \forall |\lambda| < \frac{1}{\sigma_X \sigma_Y}$$

and

$$\mathbb{P} \left(\left| \frac{1}{N} \sum_{i=1}^N X_i Y_i \right| \geq \epsilon \right) \leq 2 \exp \left(- \frac{N \epsilon^2}{2(\sigma_X^2 \sigma_Y^2 + \sigma_X \sigma_Y \epsilon)} \right)$$

■