

CASE REPORT

Malignant peripheral nerve sheath tumor in the paraspinal region mimicking a benign peripheral nerve sheath tumor: a case report

Weibo Pan¹ · Bin Feng¹ · Zhan Wang¹ · Nong Lin¹ · Zhaoming Ye¹

Received: 1 August 2016 / Revised: 4 September 2016 / Accepted: 18 September 2016
© Springer-Verlag Berlin Heidelberg 2016

Abstract

Purpose Malignant peripheral nerve sheath tumors are extremely rare in the general population and display a predilection for metastasis to the lungs. Here, we present a rare case of a malignant peripheral nerve sheath tumor located in the paraspinal region and highlight the importance of preoperative biopsy in diagnosis of spinal epidural peripheral nerve sheath tumors.

Methods We describe the clinical course of the patient as well as the radiological and pathological findings of the tumor.

Results A 14-year-old girl presented with a six-month history of sacral pain. Occasionally she experienced left leg pain and abnormal gait. General physical examination revealed sensorial loss in the L5–S1 regions. T1-weighted sagittal MRI showed a hypointense oval mass and the contrast-enhanced T1-weighted axial MRI image showed heterogeneous enhancement of the tumor. On CT imaging, this tumor characteristically appears as a dumbbell-like mass with punctate calcification and widening L5–S1 intervertebral foramen. Complete resection was performed using an anterior approach. Intraoperative pathological examination revealed evidence of malignancy and subsequent immunohistochemical analysis of the tumor confirmed the diagnosis of MPNST. The postoperative course

was uneventful and the patient has had significant improvement in her symptoms 1 month postoperatively.

Conclusions Preoperative biopsy should be routinely performed for pathological differential diagnosis of spinal epidural PNSTs as well as surgical decision-making. Furthermore a combination of clinical manifestation, radiological findings and biopsy should also be pursued for diagnosing these tumors.

Keywords Malignant peripheral nerve sheath tumor (MPNST) · Benign peripheral nerve sheath tumor (BPNST) · Spinal epidural peripheral nerve sheath tumors (spinal epidural PNST) · Biopsy · Diagnosis

Introduction

Malignant peripheral nerve sheath tumors (MPNSTs) are malignant tumors derived from a major or minor peripheral nerve branch or sheath of peripheral nerve fibers. MPNST usually occurs between 20 and 50 years of age, comprising about 5–10 % of soft tissue sarcomas [1, 2]. Although MPNST is uncommon in young patients, it also represents about 5–10 % of all soft tissue sarcomas of childhood [3, 4]. MPNSTs are most commonly located in the trunk and extremities [5]. Spinal MPNST is rare and primary spinal MPNST in children are rarer. Given its aggressive behavior, prompt and accurate diagnosis is critical for improving patient outcomes. Therefore, preoperative biopsy is necessary and helpful for surgical decision-making and preoperative preparation. In general, biopsies of spinal epidural peripheral nerve sheath tumors are performed in patients with presumed malignant tumors according to the clinical and imaging findings. However, preoperative biopsies of spinal epidural MPNSTs are often omitted due to their nonspecific and benign imaging findings.

✉ Zhaoming Ye
yezhaomingz2@163.com

¹ Department of Orthopaedics, Centre for Orthopaedic Research, Orthopedics Research Institute of Zhejiang University, The Second Affiliated Hospital of Zhejiang University School of Medicine, Zhejiang University, 88 Jiefang Road, Hangzhou, Zhejiang 310000, People's Republic of China

Here, we are describing a case of spinal epidural MPNST in the paraspinal region mimicking a BPNST to highlight the importance of preoperative biopsy in diagnosis of spinal epidural peripheral nerve sheath tumors (spinal epidural PNST).

Case report

A 14-year-old girl presented with a six-month history of sacral pain. Occasionally she experienced left leg pain and abnormal gait. During the last four months, her symptoms were aggravated and came to our hospital for treatment. General physical examination revealed sensorial loss in the L5–S1 regions. Bilateral deep tendon reflexes at both knees and ankles were normal. There were no signs of fasciculation, muscle atrophy, and upper extremity motor neuron dysfunction on her physical examination. There were no dermatologic signs of neurofibromatosis. She had no history of trauma. There were no defecation and urinary system complaints.

A plain radiograph demonstrated a mass lesion on the left sacrum and presacral space (Fig. 1). Lumbosacral CT showed a big and well-circumscribed mass extending from the L5–S1 intervertebral foramen into the left lateral aspect of the vertebra. A dumbbell-like mass with punctate calcification and widening L5–S1 intervertebral foramen was also revealed. Psoas muscle was pushed forward by the mass (Fig. 2a). After contrast enhancement, the mass presented a dumbbell-like shape and heterogeneous enhancement (Fig. 2b). The destructive and invasion of the spinal canal and erosion of the vertebral body was not revealed by CT. Moreover, CT of the chest showed no metastatic

lesions in both lungs. MRI of the lumbar spine without contrast revealed a level. Preoperative MRI showed a well-circumscribed paraspinal lesion with its size $41 \times 49 \times 49$ mm, no soft tissue involvement and complete tumor capsule. T1 weighted sagittal MRI showed a hypointense oval mass (Fig. 3a). After contrast enhancement, the mass presented a heterogeneous enhancement on T1-weighted axial MRI (Fig. 3b). On T2 weighted MRI without contrast, the mass showed a mixed signal of high and medium intensity (Fig. 3c).

On the basis of these imaging findings and clinical manifestation, we mostly suspected paraspinal benign peripheral nerve sheath tumors (BPNSTs) such as schwannoma and neurofibroma, although malignancy was not completely excluded. Then surgical resection was performed using an anterior approach. However, the intraoperative tumor capsule was incomplete and intraoperative pathological examination of a frozen section revealed evidence of malignancy. Thus, complete excision of the tumor was preferentially performed. Subsequent immunohistochemical reevaluation of the specimen showed that the tumor cells were positive for S-100, vimentin, CD31, CD34, CD56, CD117, Syn, Bcl-2, NSE, EMA, and negative for SMA, HMB45, melan-A, CD99, CgA, CK (AE1/AE3), CAM5.2, GFAP and desmin. The MIB-1 labeling index (Ki-67), which represents a marker of cell proliferation, was 30–40 %. Based on these findings, we finally diagnosed the tumor as a MPNST.

Postoperative MRI showed gross-total resection with some peri-resection site changes. The patient had a rapid recovery with no complications. Her gait is normal and her pain has completely resolved. She has no neurological deficits with no evidence of bowel/bladder dysfunction.




Fig. 1 Simple radiograph shows a mass lesion located in the left sacrum and presacral space (a, b)




Fig. 2 Computed tomography showed a big mass extending from the L5–S1 intervertebral foramen into the left lateral aspect of the vertebra. A *dumbbell-like* and soft tissue mass with punctate calcification and widening L5–S1 intervertebral foramen was also

revealed. Psoas muscle was pushed forward by the mass (a). After contrast enhancement, the mass presented a *dumbbell-like* shape and heterogeneous enhancement (b)

Follow-up MRI at 1 month showed the postoperative changes, with no evidence of residual tumor.

Discussion

MPNST has been previously known as neurofibrosarcoma, neurogenic sarcoma, malignant schwannoma, and malignant neurilemmoma [6]. The major differential diagnosis is BPNSTs. The mainstay treatment for MPNST is complete resection with the aim of achieving clear surgical margins [7]. While BPNSTs well respond to marginal excision with preservation of neurologic function [8]. Thus, the early and accurate diagnosis for spinal epidural PNSTs is particularly important for the choice of optimal treatment.

The preferred modality for imaging of MPNSTs is MRI for the primary lesion. However, benign imaging findings of our case with MPNST were actually shown on MRI and CT. T1 weighted sagittal MRI showed a hypointense oval mass and the contrast-enhanced T1-weighted axial MRI image showed heterogeneous enhancement of the tumor. CT chest was done to exclude pulmonary metastases. On CT imaging, this tumor characteristically appears as a dumbbell-like mass with punctate calcification and widening L5–S1 intervertebral foramen. As we all know, benign spinal dumbbell tumors (SDTs) such as schwannomas and neurofibromas occur frequently while malignant SDTs, including MPNSTs, are extremely rare [9]. So, paraspinal BPNSTs such as schwannoma and neurofibroma were mostly suspected. We agreed that preoperative

diagnosis based on imaging findings would be helpful in selecting therapeutic methods for SDTs. Surprisingly, intraoperative pathological examination of a frozen section revealed evidence of malignancy and postoperative immunohistochemical reevaluation of the specimen showed MPNST. Additionally, the intraoperative tumor capsule is incomplete, which was contrary to preoperative MRI findings. Recently, evidence shows that MRI is unreliable to determine the diagnosis of MPNST and guide surgical decision-making [10]. Therefore, pre-operative biopsy is essential for precise diagnosis of spinal epidural PNSTs.

Matsumoto Y et al. has recently established a new scoring system, the dumbbell scoring system (DSS), for preoperative evaluation of the malignant potential of SDTs. This scoring system based on imaging findings includes four predictors namely size, boundary, irregularly shape and osteolytic bone destruction. If the DSS score is equal to or higher than 3, biopsies was recommended to confirm the histological diagnosis [11]. According to the rules of DSS, the DSS score of our case was 0, which meant preoperative biopsy was not definitely recommended. However, our patient without biopsy was finally diagnosed as a MPNST. Of course, the limitations of DSS included its retrospective design and small sample size, which was mentioned in the study [11]. Ozawa H et al. found that malignant dumbbell tumors were more common in children younger than 10 years of age than in older patients [9]. So other factors may also be considered in creating the new DSS in the future studies. Currently, however, we hold the




Fig. 3 T1-weighted sagittal MRI showed a hypointense oval mass (a). The contrast-enhanced T1-weighted axial MRI image showed heterogeneous enhancement of the tumor (b). The mass showed a

mixed signal of high and medium intensity on T2-weighted MRI without contrast (c)

view that a combination of clinical manifestation, radiological findings and biopsy should be used for preoperative diagnosis of SDTs.

Biopsy of tumors located in the spine may be associated with pain, risk, expense, and needle tract seeding. But Pianta M et al. [12] reported that CT-guided core biopsy of PNSTs is a safe procedure with excellent diagnostic accuracy (100 %). The only occurring complication (12 %) was pain exacerbation but resolved within 1 week or after a longer duration. Needle tract seeding refers to implantation of tumor cells by contamination when biopsy needles are used to obtain tumor tissues. To our knowledge, while this iatrogenic

phenomenon is reported in treating spinal metastases [13, 14], it has not been reported in spinal epidural PNSTs. Moreover, the incidence of the needle tract seeding may be avoided with precautionary measures and adopting the local treatment methods such as wide excision and radiotherapy which can remove or kill the implanted tumor cells along the needle tract [15]. In brief, biopsy of spinal epidural PNSTs is a safe and accurate procedure for diagnosing these tumors. Therefore, while weighing the pros and cons of preoperative biopsy, we believe it should be conducive to confirming the pathological nature of spinal epidural PNSTs, and developing a comprehensive treatment procedure.

Conclusions

On the whole, we recommend that preoperative biopsy should be routinely performed for pathological differential diagnosis of spinal epidural PNSTs as well as surgical decision-making. Furthermore a combination of clinical manifestation, radiological findings and biopsy should also be pursued for diagnosing these tumors.

Compliance with ethical standards

Conflict of interest None of the authors has any potential conflict of interest.

Funding No funds were received in support of this work.

Informed consent The patient and/or her family were informed that data from the case would be submitted for publication, and gave their consent.

References

1. Doorn PF, Molenaar WM, Buter J, Hoekstra HJ (1995) Malignant peripheral nerve sheath tumors in patients with and without neurofibromatosis. *Eur J Surg Oncol* 21(1):78–82
2. Kar M, Deo SV, Shukla NK, Malik A, DattaGupta S, Mohanti BK, Thulkar S (2006) Malignant peripheral nerve sheath tumors (MPNST)—clinicopathological study and treatment outcome of twenty-four cases. *World J Surg Oncol* 4:55
3. deCou JM, Rao BN, Parham DM, Lobe TE, Bowman L, Pappo AS, Fontanesi J (1995) Malignant peripheral nerve sheath tumors: the St. Jude Children's Research Hospital experience. *Ann Surg Oncol* 2(6):524–529
4. Stucky CC, Johnson KN, Gray RJ, Pockaj BA, Ocal IT, Rose PS, Wasif N (2012) Malignant peripheral nerve sheath tumors (MPNST): the Mayo clinic experience. *Ann Surg Oncol* 19(3):878–885
5. Anghileri M et al (2006) Malignant peripheral nerve sheath tumors: prognostic factors and survival in a series of patients treated at a single institution. *Cancer* 107(5):1065–1074
6. Gupta G, Mammis A, Maniker A (2008) Malignant peripheral nerve sheath tumors. *Neurosurg Clin N Am* 19(4):533–543, v
7. Valentin T, Le Cesne A, Ray-Coquard I, Italiano A, Decanter G, Bompas E, Isambert N, Thariat J, Linassier C, Bertucci F, Bay JO, Bellesoeur A, Penel N, Le Guellec S, Filleron T, Chevreau C (2016) Management and prognosis of malignant peripheral nerve sheath tumors: the experience of the French Sarcoma Group (GSF-GETO). *Eur J Cancer* 56:77–84
8. Huang JH, Zhang J, Zager EL (2005) Diagnosis and treatment options for nerve sheath tumors. *Expert Rev Neurother* 5(4):515–523
9. Ozawa H, Kokubun S, Aizawa T, Hoshikawa T, Kawahara C (2007) Spinal dumbbell tumors: an analysis of a series of 118 cases. *J Neurosurg Spine* 7(6):587–593
10. Karsy M, Guan J, Ravindra VM, Stilwill S, Mahan MA (2016) Diagnostic quality of magnetic resonance imaging interpretation for peripheral nerve sheath tumors: can malignancy be determined? *J Neurol Surg A Cent Eur Neurosurg*. doi:[10.1055/s-0036-1584213](https://doi.org/10.1055/s-0036-1584213) (**Epub ahead of print**)
11. Matsumoto Y1, Harimaya K, Kawaguchi K, Hayashida M, Okada S, Doi T, Iwamoto Y (2016) Dumbbell scoring system: a new method for the differential diagnosis of malignant and benign spinal dumbbell tumors. *Spine (Phila Pa 1976)*. doi:[10.1097/BRS.0000000000001582](https://doi.org/10.1097/BRS.0000000000001582) (**Epub ahead of print**)
12. Pianta M, Chock E, Schlicht S, McCombe D (2015) Accuracy and complications of CT-guided core needle biopsy of peripheral nerve sheath tumours. *Skeletal Radiol* 44(9):1341–1349
13. Lee CH, Lee JW, Hyun SJ, Kim KJ, Jahng TA, Kim HJ (2014) Needle-tract seeding after percutaneous vertebroplasty: a case report. *Spine (Phila Pa 1976)* 39(12):E752–756
14. Chen YJ, Chang GC, Chen WH, Hsu HC, Lee TS (2007) Local metastases along the tract of needle: a rare complication of vertebroplasty in treating spinal metastases. *Spine (Phila Pa 1976)* 32(21):E615–618
15. Tyagi R, Dey P (2014) Needle tract seeding: an avoidable complication. *Diagn Cytopathol* 42(7):636–640