



## IMAGES OF SPINE CARE

## Extranodal non-Hodgkin's lymphoma with multiple vertebral and hepatic lesions without involvement of lymph nodes

A 55-year-old man was admitted to the hospital because of severe lumbago. Laboratory findings including soluble interleukin-2 receptor revealed almost normal levels except for alkaline phosphatase (367 U/L) and lactate dehydrogenase (511 U/L). Computed tomography, magnetic resonance imaging, and bone scintigraphy demonstrated bone lesions at Th7, 11, 12, and L4 (Fig. 1A–C). Because of the progressive paralyses of bilateral lower extremities, the pedicle of Th7 was urgently removed. Multiple masses in the liver were also observed (Fig. 1D), without splenomegaly or enlarged lymph nodes.

Metastatic cancer to the bone and liver was highly suspected because: he had no B symptoms associated with aggressive lymphoma; serum chemistry revealed elevated levels of ALP and LDH, which are frequently observed in

bone tumors; and no other lesions in lymph node or organs were demonstrated. From the histologic examinations of the liver and the pedicle of Th7; however, we diagnosed the patient as diffuse large B-cell lymphoma (Fig. 2). Bone marrow examination from the right posterior iliac crest revealed no invasion by lymphoma cells.

The patient received six courses of chemotherapy with rituximab, which was followed by the complete remission.

Primary lymphoma of the bone is rare, representing 1% of malignant lymphoma. The long bones are biased for involvement, such as the femur and humerus [1]. In our case, the lesions were located at Th7, 11, 12, and L4, with no other bone lesions. Also, the extension of lymphoma cells to the spines from the whole bone marrow infiltration was ruled out. Although the incidence of a single vertebral lymphoma is reported to be 1.7% of all primary lymphomas of the bone, multiple primary vertebral lymphoma without involvement to other bones is extremely rare as we can find only two cases in the literature [2,3]. Primary hepatic lymphoma is also rare, representing 0.016% of non-Hodgkin's lymphoma (NHL) [4,5]. A previous report on a rare case of NHL of the bone and liver without




Fig. 1. Images of bone and liver. (A–C) Bone lesions. Magnetic resonance imaging of the spine (A, B) demonstrates bone masses at Th7 (B), 11, and L4 (arrows). The lesions were detected by  $^{99m}\text{Tc}$ -bone scintigraphy (C), with no other bone lesions. The highest accumulation of  $^{99m}\text{Tc}$  is observed at Th7. (D) Hepatic lesions. Multiple large masses are observed in the enhanced CT. The spleen is not enlarged and has no masses, and no enlarged lymph nodes are observed.




Fig. 2. Histologic findings. In the histologic studies of the pedicle of Th7 (A) and liver (needle-biopsied; (B)), the lesion was characterized by a diffuse growth pattern of lymphocytes without follicles at a low-power field (A, B). At a high-power field, diffuse proliferation of medium-to-large-sized lymphoma cells with large nucleus was observed (C). An immunohistochemical study demonstrated that the lymphoma cells were negative for CD3, CD4, CD5, CD8, CD10, CD30, and ALK-1 and positive for CD20 (D), BCL6 (E), and MUM-1 (F). Chromosomal analysis and flow cytometric analysis were not available because of the limited size of the samples.

lymphadenopathy [6], showed that the affected bones were widely spread. However, to the best of our knowledge, this is the first report of NHL in only the vertebral bones and hepatic lesions. Clinicians should be aware of NHL that mimics metastatic cancer to the bone and liver in images and clinical findings, and future accumulation of similar cases is expected.

## References

- [1] Heyning FH, Hogendoorn PC, Kramer MH, Hermans J, Kluin-Nelemans JC, Noordijk EM, et al. Primary non-Hodgkin's lymphoma of bone: a clinicopathological investigation of 60 cases. *Leukemia* 1999;13:2094–8.
- [2] Ebus SCM, Bensen HJJA, Norel van GJ, Donk R. Primary non-Hodgkin's lymphoma in multiple vertebrae presenting as a lumbar radicular syndrome. *Spine* 2002;27:E271–3.
- [3] Tabata R, Tabata C, Numa M, Nagai T. Extranodal follicular lymphoma exclusively featured by diffuse pattern originating from multiple vertebral bones. *Ann Hematol* 2008;87:581–3.
- [4] Kikuma K, Watanabe J, Oshiro Y, Shimogama T, Honda Y, Okamura S, et al. Etiological factors in primary hepatic B-cell lymphoma. *Virchows Arch* 2012;460:379–87.
- [5] Noronha V, Shafi NQ, Obando JA, Kummar S. Primary non-Hodgkin's lymphoma of the liver. *Crit Rev Oncol Hematol* 2005;53:199–207.
- [6] Cheng G, Servaes S, Chamroonrat W, Li J, Zhuang H. Non-Hodgkin's lymphoma of the bone and the liver without lymphadenopathy revealed on FDG-PET/CT. *Clin Imaging* 2010;34:476–9.

Rie Tabata, MD<sup>a</sup>

Chiharu Tabata, MD<sup>b</sup>

Takami Ishikawa, MD<sup>c</sup>

Ryoji Yasumizu, MD<sup>d</sup>

<sup>a</sup>Department of Hematology and Oncology  
Hyogo Prefectural Tsukaguchi Hospital  
6-8-17 Minamitsukaguchi  
Amagasaki, Hyogo, 661-0012 Japan

<sup>b</sup>Hyogo College of Medicine Cancer Center  
1-1 Mukogawa-cho  
Nishinomiya, Hyogo  
663-8501 Japan

<sup>c</sup>Department of Orthopedics  
Hyogo Prefectural Tsukaguchi Hospital  
6-8-17 Minamitsukaguchi  
Amagasaki, Hyogo, Japan

<sup>d</sup>Department of Pathology  
Hyogo Prefectural Tsukaguchi Hospital  
6-8-17 Minamitsukaguchi  
Amagasaki, Hyogo, Japan

FDA device/drug status: Approved (rituximab, doxorubicin hydrochloride, vincristine sulfate, cyclophosphamide, prednisolone).

Author disclosures: **RT**: Nothing to disclose. **CT**: Nothing to disclose.

**TI**: Nothing to disclose. **RY**: Nothing to disclose.

The authors declare that they have no conflicts of interest.

The authors declare that they have no funding support.