

Case Report

Osteoid osteoma presenting as thoracic scoliosis

Masashi Uehara, MD, Jun Takahashi, MD*, Shugo Kuraishi, MD, Masayuki Shimizu, MD,
Shota Ikegami, MD, Toshimasa Futatsugi, MD, Hiroyuki Kato, MD

Department of Orthopaedic Surgery, Shinshu University School of Medicine, 3-1-1 Asahi, Matsumoto-City, Nagano 390-8621, Japan

Received 11 May 2015; revised 28 July 2015; accepted 27 August 2015

Abstract

BACKGROUND CONTEXT: Osteoid osteoma of the thoracic spine is relatively uncommon and is often difficult to diagnose, especially when patients do not complain of pain.

PURPOSE: This study aims to describe an unusual case of scoliosis caused by osteoid osteoma of the thoracic spine that was challenging to diagnose.

STUDY DESIGN/SETTING: A case report of a 12-year-old girl who presented with scoliosis caused by osteoid osteoma of the thoracic spine without apparent pain was carried out.

METHODS: Diagnosis of the lesion was made using computed tomography (CT) and magnetic resonance imaging as well as the Scoliosis Research Society-22 (SRS-22) patient-based questionnaire.

RESULTS: A preoperative CT myelogram revealed a mass lesion in the lamina of the 10th thoracic vertebra that was considered to be osteoid osteoma. This diagnosis was histologically confirmed following tumor excision. The patient's spinal deformity and SRS-22 scores were both improved at 5 months postoperatively.

CONCLUSIONS: Osteoid osteoma of the thoracic spine may present as non-painful scoliosis. Tumor resection is effective. Clinicians should bear this uncommon lesion in mind during recommended CT examination before scoliosis surgery. © 2015 Elsevier Inc. All rights reserved.

Keywords:

Osteoid osteoma; Scoliosis; Spinal deformity; Tumor excision; Vertebral rotation; CT myelogram

Introduction

Osteoid osteoma was first described by Jaffe in 1935 [1]. It is a benign reactive bone lesion that most frequently occurs in male adolescents between 10 and 20 years of age [2,3]. The predominant cause of painful scoliosis in adolescents is osteoid osteoma of the spine [4]. Approximately 10%–25% of osteoid osteoma cases occur in the spine [5–9]: 60% of tumors are found in the lumbar spine, 27% in the cervical spine, 12% in the thoracic spine, and 2% on the sacrum. There is a very strong correlation between the neoplasm and scoliosis because two-thirds of cases manifest as painful scoliosis [10]. However, osteoid osteoma is commonly missed in a large,

albeit undetermined, number of patients [11] because the lesion is often recognized only after months or years of ineffective bracing [12].

Osteoid osteoma of the thoracic spine is relatively uncommon. It is often difficult to diagnose, especially when the patient does not complain of pain. We herein describe the clinical outcome of a girl who presented with scoliosis caused by osteoid osteoma of the thoracic spine that was detected by preoperative computed tomography (CT).

Case report

A 12-year-old girl was suspected to have scoliosis in a routine school health screening in May 2013. Because her condition worsened despite the use of a brace, she visited our institution for scoliosis correction surgery in April 2014. Her height was 154 cm and her weight was 40 kg at presentation. She had pervasive developmental disorders along with an expressive language impediment, although her intelligence quotient score was average. She possessed no family history of scoliosis. She complained of no additional

FDA device/drug status: Not applicable.

Author disclosures: **MU:** Nothing to disclose. **JT:** Nothing to disclose. **SK:** Nothing to disclose. **MS:** Nothing to disclose. **SI:** Nothing to disclose. **TF:** Nothing to disclose. **HK:** Nothing to disclose.

* Corresponding author. Department of Orthopaedic Surgery, Shinshu University School of Medicine, 3-1-1 Asahi, Matsumoto-City, Nagano 390-8621, Japan. Tel.: (81) 263-37-2659; fax: (81) 263-35-8844.

E-mail address: jtaka@shinshu-u.ac.jp (J. Takahashi).




Fig. 1. (Left) Apical trunk rotation (ATR) was 19° as determined by a scoliometer. (Right) ATR had improved to 9° at 5 months postoperation.

symptoms, including back pain. Physical examination revealed right shoulder elevation. Apical trunk rotation, as determined by a scoliometer, was 19° (Fig. 1, Left). We observed no neurologic deficits. Whole spine radiographs revealed a Lenke-type 1AN curve with a Cobb angle of 45° from T5 to L2 (main thoracic) and 28° from L2 to L5 (lumbar) (Fig. 2, Left). Risser grade was 2.

Magnetic resonance imaging of the whole spine did not reveal any evidence of Arnold-Chiari malformation, syringomyelia, or dural ectasia (Fig. 3). As we had planned to perform posterior correction surgery for the patient's scoliosis, a CT examination was performed after myelography, which demonstrated a mass lesion in the lamina of the 10th thoracic vertebra with a visible nidus (Fig. 4). A re-examination of the MRI results verified this finding. The patient's Scoliosis Research Society-22 (SRS-22) patient questionnaire scores for function, pain, self-image, mental health, and subtotal were 4.5, 5.0, 2.4, 4.0, and 3.9, respectively. This enabled a revised diagnosis of osteoid osteoma, and the lesion was

surgically excised. Histologic tissue findings confirmed our diagnosis (Fig. 5).

Postoperative brace therapy improved the patient's spinal deformity. Her shoulder imbalance was diminished and her apical trunk rotation had improved to 9° at 5 months after surgery (Fig. 1, Right). At 5 months postoperatively, Cobb angle from T5 to L2 was ameliorated at 30° (Fig. 2, Right) and SRS-22 scores for function, pain, self image, mental health, subtotal, satisfaction, and total had improved to 4.8, 5.0, 3.4, 4.6, 4.5, 4.0, and 4.4, respectively.

The patient and her family have provided consent for the data in this case to be published.

Discussion

Osteoid osteoma is a benign osteoblastic neoplasm that is characterized by a well-demarcated core of <1 cm and a distinctive surrounding zone of reactive bone formation [13]. Although the lesions may occur anywhere in the cortex or




Fig. 2. (Left) Whole spine radiographs showed a Lenke type 1AN curve with a Cobb angle of 45°. (Right) Cobb angle had improved to 30° at 5 months postoperation.




Fig. 3. MRI of the whole spine showed no signs of Arnold-Chiari malformation, syringomyelia, or dural ectasia.




Fig. 4. CT revealed a mass lesion in the lamina of the 10th thoracic vertebra. The lesion had a visible nidus (arrow).

medulla of the skeleton, they usually affect the lower extremities. Approximately 10%–25% of osteoid osteoma cases manifest in the spine [5–9], with a tendency to involve the posterior elements that range from 70% to 100% [14–16]. At an incidence of only 12% of spinal cases, osteoid osteoma of the thoracic spine is relatively rare [10].

Pain is the primary symptom of initial and recurrent disease [17], although reports of osteoid osteoma without the presence of pain exist [18]. In this case report, the SRS-22 questionnaire before surgery revealed a pain score of 5.0. It is difficult to diagnose osteoid osteoma without the presence of pain.

We first became aware of the osteoid osteoma lesion in a CT image. Indeed, preoperative CT is indispensable for not overlooking the presence of osteoid osteoma. Furthermore, 3D-CT revealed that the spinal curvature in this patient had

less rotation than that in typical adolescent idiopathic scoliosis (AIS) (Fig. 6), which confirmed the utility of CT before possibly unnecessary scoliosis surgery. Surgical removal of the lesion improved the patient's deformities and survey scores. Retrospective examination of preoperative radiographs showed that pedicle width on the convex side was wide in the apex region. Because the pedicle width on the concave side is narrow due to spinal rotation in AIS patients, pain-related scoliosis may be considered when no vertebral rotation is evident.

Scoliosis with early age onset, long C curve, and left thoracic curve is often associated with a Chiari malformation or syringomyelia [19], whereas a short, sharp, angular curve is usually present in cases of neurofibromatosis type 1 [20]. Meanwhile, scoliosis in Marfan syndrome tends to exhibit longer kyphosis extending through the thoracolumbar junction [21]. Careful examination of curve types is essential for




Fig. 5. Histologic examination of the resected tumor with hematoxylin and eosin (HE) staining showed calcific and non-calcific osteoid tissue, which confirmed the diagnosis of osteoid osteoma.



Fig. 6. 3D-CT revealed that the spinal curvature in the patient had less rotation than that in a typical adolescent idiopathic scoliosis.

the correct diagnosis of AIS. A limitation of this case report is a short follow-up period.

Conclusions

We encountered an unusual case of osteoid osteoma of the thoracic spine presenting as scoliosis that was difficult to diagnose. We successfully treated the patient by tumor excision, but continued follow-up is needed. Clinicians should bear in mind the possibility of osteoid osteoma before scoliosis surgery, even in patients without apparent pain, and carefully examine CT findings for evidence of tumor involvement.

References

- [1] Jaffe HL. Osteoid osteoma. A benign osteoblastic tumor composed of osteoid and atypical bone. *Arch Surg* 1935;31:709–28.
- [2] Aegerter EE, Kirkpatrick JA Jr, editors. *Orthopedic diseases*. 3rd ed. Philadelphia: WB Saunders; 1968.
- [3] Mehta MH. Pain provoked scoliosis. Observations on the evolution of the deformity. *Clin Orthop Relat Res* 1978;135:58–65.
- [4] Zileli M, Cagli S, Basdemir G, Ersahin Y. Osteoid osteomas and osteoblastomas of the spine. *Neurosurg Focus* 2003;15:E5.
- [5] Azouz EM, Kozlowski K, Marton D, Sprague P, Zerhouni A, Asselah F. Osteoid osteoma and osteoblastoma of the spine in children. Report of 22 cases with brief literature review. *Pediatr Radiol* 1986;16:25–31.
- [6] Barei DP, Moreau G, Scarborough MT, Neel MD. Percutaneous radiofrequency ablation of osteoid osteoma. *Clin Orthop Relat Res* 2000;373:115–24.
- [7] Boriani S, Weinstein JN. Differential diagnosis and surgical treatment of primary benign and malignant neoplasms. In: Frymoyer JN, editor. *The adult spine. Principles and practice*. 2nd ed. Philadelphia, PA: Lippincott Raven; 1997. p. 950–87.
- [8] Jackson RP, Reckling PW, Mants FA. Osteoid osteoma and osteoblastoma: similar histology lesions with different natural histories. *Clin Orthop Relat Res* 1977;128:303–13.
- [9] Lindner NJ, Ozaki T, Roedl R, Gosheger G, Winkelmann W, Wörtler K. Percutaneous radiofrequency ablation in osteoid osteoma. *J Bone Joint Surg Br* 2001;83:391–6.
- [10] Janin Y, Epstein JA, Carras R, Khan A. Osteoid osteomas and osteoblastomas of the spine. *Neurosurgery* 1981;8:31–8.
- [11] Keim HA, Reina EG. Osteoid-osteoma as a cause of scoliosis. *J Bone Joint Surg Am* 1975;57:159–63.
- [12] Kirwan EO, Hutton PA, Pozo JL, Ransford AO. Osteoid osteoma and benign osteoblastoma of the spine. Clinical presentation and treatment. *J Bone Joint Surg Br* 1984;66:21–6.
- [13] Unni KK, editors. *Dahlin's bone tumors: general aspects and data on 11,087 cases*. 5th ed. Philadelphia, PA: Lippincott-Raven; 1996.
- [14] Crouzet G, Mnif J, Vasdev A, Pascal-Ortiz D, Chirossel JP, Pasquier B. Osteoid osteoma of the spine: radiological aspects and value of arteriography. Four cases. *J Neuroradiol* 1989;16:145–59.
- [15] Fountain EM, Burge CH. Osteoid osteoma of the cervical spine. A review and case report. *J Neurosurg* 1961;18:380–3.
- [16] Marcove R, Heelan R, Huvos A, Healey J, Lindeque BG. Osteoid osteoma: diagnosis, localization and treatment. *Clin Orthop Relat Res* 1991;267:197–201.
- [17] Atesok KI, Alman BA, Schemitsch EH, Peyser A, Mankin H. Osteoid osteoma and osteoblastoma. *J Am Acad Orthop Surg* 2011;19:678–89.
- [18] McDermott MB, Kyriakos M, McEnery K. Painless osteoid osteoma of the rib in an adult. A case report and a review of the literature. *Cancer* 1996;77:1442–9.
- [19] Wu L, Qiu Y, Wang B, Zhu ZZ, Ma WW. The left thoracic curve pattern: a strong predictor for neural axis abnormalities in patients with “idiopathic” scoliosis. *Spine* 2010;15:182–5.
- [20] Singh K, Samartzis D, An HS. Neurofibromatosis type 1 with severe dystrophic kyphoscoliosis and its operative management via a simultaneous anterior-posterior approach: a case report and review of the literature. *Spine J* 2005;5:461–6.
- [21] Sponseller PD, Hobbs W, Riley LH 3rd, Pyeritz RE. The thoracolumbar spine in Marfan syndrome. *J Bone Joint Surg Am* 1995;77:867–76.