

Accepted Manuscript

Title: Surgical management of gorham disease involving the upper cervical spine with occipito-cervical-thoracic fusion: a case report

Author: Adam Schell, John M. Rhee, Abigail Allen, Lindsay Andras, Feifei Zhou

PII: S1529-9430(16)00373-9

DOI: <http://dx.doi.org/doi: 10.1016/j.spinee.2016.02.020>

Reference: SPINEE 56908

To appear in: *The Spine Journal*

Received date: 29-1-2016

Accepted date: 4-2-2016

Please cite this article as: Adam Schell, John M. Rhee, Abigail Allen, Lindsay Andras, Feifei Zhou, Surgical management of gorham disease involving the upper cervical spine with occipito-cervical-thoracic fusion: a case report, *The Spine Journal* (2016), <http://dx.doi.org/doi: 10.1016/j.spinee.2016.02.020>.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Surgical Management of Gorham Disease involving the Upper Cervical Spine with occipito-cervical-thoracic fusion: A Case Report

Adam Schell MD¹, John M. Rhee MD^{2*}, Abigail Allen MD³, Lindsay Andras MD⁴, and Feifei Zhou MD^{2,5}

*To whom correspondence should be addressed
Department of Orthopaedic Surgery
Emory Spine Center
Emory University
Atlanta, GA

1. University Hospitals of Cleveland, Case Medical Center, Case Western Reserve University School of Medicine, Cleveland, Ohio 44106, USA
2. Department of Orthopedic Surgery, Emory Spine Center, Emory University, Atlanta, GA 30329, USA
3. Pediatric Orthopaedic Clinic Leni & Peter May Department of Orthopaedics, Icahn School of Medicine at Mount Sinai, New York, NY 10029, USA
4. Children's Orthopedic Center, Children's Hospital Los Angeles, Los Angeles, CA 90027, USA
5. Orthopaedics Department, Peking University Third Hospital, Beijing 100191, P.R. China

Keywords: cervical spine, osteolysis, gorham disease, occipito-cervicalthoracic fusion

Abstract

Background context: Gorham disease is a rare osteolytic condition with only 17 reported cases primarily affecting the cervical spine.

Purpose: We report the case of a 31-year-old woman with severe kyphotic instability and myelopathy of the upper cervical spine secondary to massive osteolysis of the posterior elements. The surgical management, clinical outcome, and review of the relevant literature are discussed.

Study Design: Case report and literature review

Patient Sample: One patient with Gorham disease of the cervical spine.

Methods: We report the case of a 31-year-old woman with Gorham disease affecting the upper cervical spine. Combined posterior and anterior stabilization and fusion was performed from the occiput to the thoracic spine.

Results: Six-year follow-up with annual CT imaging showed solid fusion from occiput to T2. No spread or local recurrence has developed to date.

Conclusions: Correction of severe instability with myelopathy due to Gorham disease of the upper cervical spine was achieved through posterior and anterior occipito-cervical-thoracic fusion.

Introduction

Sub-acute osteolysis in an otherwise healthy individual is an unusual clinical finding. In the absence of malignancy, nephropathy, infection, or metabolic disease, a small group of disorders can cause this phenomenon(1). One such disorder is Gorham disease of bone, a rare condition of unregulated bone resorption related to focal areas of non-neoplastic vascular tissue proliferation(2). Gorham disease was first described in 1838(3), and since then more than 300 cases have been reported. It is known by at least ten names, including “vanishing bone disease,” “massive osteolysis,” and “phantom bone disease” among others. While substantial effort has been devoted to understanding this disorder, the etiology and underlying pathophysiology remain unclear(4).

The natural history varies widely among individuals and depends on the skeletal location of disease and time of diagnosis. The most common site of involvement is the shoulder, followed by the pelvis, ribs, and the skull. To date, approximately 50 cases affecting the spine have been reported, but only 17 of those primarily involved the cervical spine(5-7). A spectrum of clinical outcomes is reported when cervical vertebrae are affected, ranging from fatalities due to complications of osteolysis(8) to spontaneous osteolytic arrest after only conservative management(9).

Treatment of the underlying disorder includes both medical and surgical strategies, and spinal involvement typically necessitates surgical fusion to maintain stability and prevent neurologic injury. Historically, bone grafting in Gorham patients has been met with limited success due to graft osteolysis and poor graft ingrowth(10, 11). Herein we report the case of a 31-year-old woman with severe cervical instability secondary to Gorham disease affecting the upper cervical spine, and we provide clinical outcomes up to 6 years following circumferential fusion from the occiput to the thoracic spine.

Case Report

A 31 year old woman with a 7-year history of axial neck pain presented to the senior author with a one month history of rapidly progressive exacerbation of neck pain associated with bilateral hand numbness. The numbness was most severe while sitting upright and relieved by lying supine, and this sensory deficit made holding objects in her hands difficult. She also described gradual worsening of chronic neck pain in recent years, for which she had previously sought treatment by numerous physicians and chiropractors to no avail. Her past medical history was negative for arthritides, malignancy, endocrine disease, or familial disorders.

On physical exam, the posterior elements in the upper cervical spine were notably absent to palpation. Neurological exam was non-focal with no motor impairment or long tract signs. Her laboratory studies were within normal limits and showed no evidence of infection or systemic inflammatory disease. Flexion-extension lateral radiographs revealed the dorsal elements and facet complex were both missing at the levels of C2 and

C3 but had been present on historical radiographs from only 3 years prior (**Figure 1**). The arch of C1 was thinned, and the proximal portion of the C4 lamina and spinous process were missing as well. Computed tomographic imaging revealed similar findings (**Figure 2**). Magnetic resonance imaging showed no evidence of infection, malignancy, or cord compression. However, the posterior elements were notably missing. Given the clinical and radiographic findings of progressive osteolysis without identifiable cause, a diagnosis of Gorham's vanishing bone disease was made. Due to the significant instability, emergent stabilization and fusion was recommended.

The first stage operation involved a posterior fusion and instrumentation from the occiput to T2. The proximal and distal extents of the fusion construct were chosen to widely bridge the pathologically involved segments due to the potential for local recurrence and progressive osteolysis at adjacent levels. During dissection, the arch of C1 was approximately one-third its expected size, and only a thin membrane overlaid C2-C3. However, there was no intra-operative evidence of malignant, infectious, or inflammatory processes. Because there was no observable mass, there was no specific material to biopsy – specimens sent for pathology were non-specific and consistent with bone, muscle, and fibroconnective tissue. In order to bridge the large bony defect present posteriorly, an autologous rib graft was harvested and fashioned to strut from the occiput to the spinous process of C5. Supplemental morsellized iliac crest bone graft as well as InFuse (rhBMP-2; Medtronic) were used posteriorly. An aggressive approach to bone grafting was taken given the potential for graft osteolysis and progressive local disease. The patient then underwent staged multilevel anterior discectomy and fusion from C2 to T2 (**Figure 3**) two days later, using a combination of iliac crest bone graft, allograft, and

plates. Anterior fusion was thought necessary in order to enhance the probability of a successful arthrodesis throughout the entire length of the construct, as well as to provide anterior fusion in case progressive osteolysis occurred.

The patient underwent radiation oncology consultation postoperatively, but radiation therapy was not elected. At most recent follow up (6 years), CT demonstrated solid fusion. There has been no evidence of progressive disease or osteolysis either through, or adjacent to, the fusion mass (**Figures 4**).

Discussion

Gorham disease is an exceedingly rare osteolytic disease that has received substantial attention from the medical and surgical communities since it was first comprehensively described in 1955(12). Despite more than 300 published cases and many rigorous investigations into the molecular basis of disease(4), the precise etiology is still unknown. There is no evidence to suggest heritability, and both genders are affected in similar frequency. It can occur at any age, but the majority of cases occur before age 40(2).

The pathophysiology is unknown but appears to involve both spontaneous and post-traumatic angiomatosis. Osteolysis is thought to occur as a result of angiomatoid proliferation in the affected bone, generating a lymphovascular network that reduces osseous matrix pH and local oxygen tension. The resulting environment leads to activation of hydrolytic enzymes and phosphatase secretion from osteoblasts, and a destructive remodelling imbalance favoring excessive bone resorption is cultivated(13).

Both osteoclastosis and production of interleukin-6 reportedly contribute to this osteolytic cascade(14-16).

Gorham disease is a diagnosis of exclusion, and more common causes of osteolysis (such as infection, occult malignancy, and hyperparathyroidism) must be ruled out early in the evaluation. Heffez and colleagues developed a set of criteria that may aid in establishing a diagnosis if other osteolytic processes are suspected(17). Clinical signs are often subtle and may only become apparent late in the course of the disease. Like the patient in this report, some individuals experience chronic pain long before mechanical instability arises. Meanwhile, others will notice an acute pain or deformity, especially after sustaining trauma to the affected area. For this reason, the natural history varies greatly among affected individuals, and patient outcomes are optimized when diagnosis is established early in the course of disease.

The shoulder and pelvis are the most frequently affected sites, but involvement of humerus, sternum, clavicle, skull, ribs, femur, mandible, spine, hands, and feet are each described in multiple case reports(2). Morbidity and mortality are highest when the bony thorax and/or spine are involved. A well-known complication of thoracic involvement is lymphangiomatous extension into pleural space, which may lead to development of chylothorax(18). Morbidity is also higher when the spine is affected, due to instability, neurological decline, and the resultant comorbidities that may arise in the immobilized patient. Spinal fusion in this disease setting has been associated with a high incidence of graft resorption(11), fusion failure requiring reoperation(19), and unpredictable disease progression(10).

When mechanical instability arises, surgical correction is necessary, but mechanically stable lesions may be treated non-surgically using medical and radiation therapies alone. Reported medical treatment strategies include single or combination regimens of agents such as bisphosphonates(20), alpha-interferon 2b(21), and various cytotoxic agents(22). Radiation therapy has been employed as an adjunct to surgical treatment, and reports show conventionally fractionated doses as low as 40 Gy to slow progression of active Gorham lesions when administered following radical resection of affected tissues(23, 24). In this case, clinical evidence was most consistent with an arrested Gorham process, and after consultation with a radiation oncologist it was decided to forgo XRT in the absence of disease recurrence or progression.

Sixteen cases of cervical spine involvement are reported in the literature, and eight of those cases occurred at the C2 and/or C3 level as seen in this patient(10, 11, 25-30). This vertebral level appears to pose an increased risk of mortality, as three of the five patients who have died from complications of Gorham disease of the cervical spine had disease involving these levels. In these cases, mortality may be attributed to upper cervical cord injury, cerebrovascular accident, and complications following osteolytic progression and multiple unsuccessful fusion attempts. However, favorable outcomes are reported in the majority of cases where adequate spinal fusion is achieved(5, 31). In one case, diagnosis was obtained early in the disease process, and a combination of halo traction and radiation therapy was sufficient for reversal of osteolysis(9).

The goals of treatment in this patient were to achieve spinal stability through instrumentation and arthrodesis. An aggressive approach was taken in terms of both spinal fixation and bone grafting due to the potential for local recurrence and spread of

Gorham disease, as well as the possibility that the osteolytic process might impair the process of fusion. Therefore, the fusion and fixation were performed circumferentially along with the addition of autograft rib and iliac crest. RhBMP-2 was used posteriorly as well to try to assure solid arthrodesis.

In conclusion, a case of advanced Gorham osteolysis affecting the upper cervical spine is described. While the exact etiology and pathophysiology of this disease remain mysterious, swift intervention may prevent neurologic decline in cases of osteolytic cervical instability. In the present case, correction of severe cervical instability and prevention of neurologic injury were achieved through emergent circumferential fusion from occiput to T2. Although extremely rare, Gorham disease should be considered in patients presenting with osteolytic cervical instability in the absence of a mass or infectious process. Maximal fixation and bone grafting strategies are recommended in order to ensure healing in the setting of a disease that may present with local recurrence.

References

- 1 Resnick D. Diagnosis of bone and joint disorders. 4th ed. Philadelphia: Saunders; 2002.
- 2 Patel DV. Gorham's disease or massive osteolysis. Clinical medicine & research. 2005 May;**3**(2):65-74.
- 3 Jackson J. A boneless arm. Boston Med Surg J. 1838;**1838**(18):368-9.
- 4 Dellinger MT, Garg N, Olsen BR. Viewpoints on vessels and vanishing bones in Gorham-Stout disease. Bone. 2014 Jun;**63**:47-52.
- 5 Aizawa T, Sato T, Kokubun S. Gorham disease of the spine: a case report and treatment strategies for this enigmatic bone disease. The Tohoku journal of experimental medicine. 2005 Feb;**205**(2):187-96.
- 6 Kakuta Y, Iizuka H, Kobayashi R, et al. Gorham disease of the lumbar spine with an abdominal aortic aneurysm: a case report. The spine journal : official journal of the North American Spine Society. 2014 Jan;**14**(1):e5-9.

- 7 Lala S, Mulliken JB, Alomari AI, Fishman SJ, Kozakewich HP, Chaudry G. Gorham-Stout disease and generalized lymphatic anomaly--clinical, radiologic, and histologic differentiation. *Skeletal radiology*. 2013 Jul;**42**(7):917-24.
- 8 Ellis DJ, Adams TO. Massive Osteolysis - Report of Case. *Journal of Oral Surgery*. 1971 1971;**29**(9):659-&.
- 9 Mawk JR, Obukhov SK, Nichols WD, Wynne TD, Odell JM, Urman SM. Successful conservative management of Gorham disease of the skull base and cervical spine. *Childs Nervous System*. 1997 Nov-Dec;**13**(11-12):622-5.
- 10 Woodward HR, Chan DPK, Lee J. Massive Osteolysis of the Cervical-Spine - a Case-Report of Bone-Graft Failure. *Spine*. 1981 1981;**6**(6):545-9.
- 11 Castleman B, Macausland WR, Aufranc OE, et al. Weekly Clinicopathological Exercises - Case 16-1964. *New England Journal of Medicine*. 1964 1964;**270**(14):731-&.
- 12 Gorham LW, Stout AP. Massive Osteolysis (Acute Spontaneous Absorption of Bone, Phantom Bone, Disappearing Bone) - Its Relation to Hemangiomatosis. *Journal of Bone and Joint Surgery-American Volume*. 1955 1955;**37**(5):985-1004.
- 13 Dickson GR, Mollan RA, Carr KE. Cytochemical localization of alkaline and acid phosphatase in human vanishing bone disease. *Histochemistry*. 1987;**87**(6):569-72.
- 14 Gorham LW, Stout AP. Massive osteolysis (acute spontaneous absorption of bone, phantom bone, disappearing bone); its relation to hemangiomatosis. *The Journal of bone and joint surgery American volume*. 1955 Oct;**37-A**(5):985-1004.
- 15 Devlin RD, Bone HG, 3rd, Roodman GD. Interleukin-6: a potential mediator of the massive osteolysis in patients with Gorham-Stout disease. *The Journal of clinical endocrinology and metabolism*. 1996 May;**81**(5):1893-7.
- 16 Moller G, Priemel M, Amling M, Werner M, Kuhlmeier AS, Delling G. The Gorham-Stout syndrome (Gorham's massive osteolysis). A report of six cases with histopathological findings. *The Journal of bone and joint surgery British volume*. 1999 May;**81**(3):501-6.
- 17 Heffez L, Doku HC, Carter BL, Feeney JE. Perspectives on Massive Osteolysis - Report of a Case and Review of the Literature. *Oral Surgery Oral Medicine Oral Pathology Oral Radiology and Endodontics*. 1983 1983;**55**(4):331-43.
- 18 Tie ML, Poland GA, Rosenow EC, 3rd. Chylothorax in Gorham's syndrome. A common complication of a rare disease. *Chest*. 1994 Jan;**105**(1):208-13.
- 19 Chong Ng L, Sell P. Gorham disease of the cervical spine-a case report and review of the literature. *Spine (Phila Pa 1976)*. 2003 Sep 15;**28**(18):E355-8.
- 20 Zheng MW, Yang M, Qiu JX, et al. Gorham-Stout syndrome presenting in a 5-year-old girl with a successful bisphosphonate therapeutic effect. *Experimental and therapeutic medicine*. 2012 Sep;**4**(3):449-51.
- 21 Takahashi A, Ogawa C, Kanazawa T, et al. Remission induced by interferon alfa in a patient with massive osteolysis and extension of lymph-hemangiomatosis: a severe case of Gorham-Stout syndrome. *J Pediatr Surg*. 2005 Mar;**40**(3):E47-50.
- 22 Holroyd I, Dillon M, Roberts GJ. Gorham's disease: a case (including dental presentation) of vanishing bone disease. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 2000 Jan;**89**(1):125-9.
- 23 Gupta RK, Kumar M, Verma A, et al. Gorham disease of mandible treated with post-operative radiotherapy. *South Asian journal of cancer*. 2012 Oct;**1**(2):95-7.

- 24 Handl-Zeller L, Hohenberg G. Radiotherapy of Morbus Gorham-Stout: the biological value of low irradiation dose. *Br J Radiol.* 1990 Mar;**63**(747):206-8.
- 25 Puertas EB, Wajchenberg M, Jesus-Garcia R, Reis Rodrigues LM. From Imaging and Surgical Outcome.... *The Spine Journal.* 2006;**6**(4):468-70.
- 26 Kai B, Ryan A, Munk PL, Dunlop P. Gorham disease of bone: three cases and review of radiological features. *Clinical radiology.* 2006 Dec;**61**(12):1058-64.
- 27 Frame B, Herrera LF, Mitchell DC, Fine G. Massive osteolysis and tumoral calcinosis. *The American journal of medicine.* 1971 Mar;**50**(3):408-12.
- 28 Foulst H, Goupille P, Aesch B, Valat JP, Burdin P, Jan M. Massive osteolysis of the cervical spine. A case report. *Spine (Phila Pa 1976).* 1995 Jul 15;**20**(14):1636-9.
- 29 Hambach R, Pujman J, Maly V. Massive osteolysis due to hemangiomatosis; report of a case of Gorham's disease with autopsy. *Radiology.* 1958 Jul;**71**(1):43-7.
- 30 Bohlman HH, Sachs BL, Carter JR, Riley L, Robinson RA. Primary Neoplasms of the Cervical-Spine - Diagnosis and Treatment of 23 Patients. *Journal of Bone and Joint Surgery-American Volume.* 1986 Apr;**68A**(4):483-94.
- 31 Puertas EB, Wajchenberg M, Jesus-Garcia R, Rodrigues LM. Where cord was compressed by disappearing bone. *The spine journal : official journal of the North American Spine Society.* 2006 Jul-Aug;**6**(4):468-70.

Figure 1. Lateral cervical spine radiographs. A) Neutral upright radiograph taken three years prior to presentation. At that time, the patient was complaining of gradually worsening axial neck pain but had no neurologic symptoms. Close inspection of the lateral radiograph demonstrates a somewhat mottled appearance to the C2 pars, facet, and lamina, as well as an indistinct cortical margin along the upper border of the C2 posterior elements. (B) Lateral upright radiograph taken at the time of presentation. The C1 laminar arch is notably thin. The facets, pars, lamina, and spinous process of C2 are completely absent. Large areas of the posterior elements of C3 and C4 are missing. Spondyloslistheses are noted at C2-3 and C3-4, along with a generalized kyphosis.

Figure 2. Axial Computed tomography images. There is severe destruction within the upper cervical spine and complete absence of the lamina, pedicles, and posterior elements at the level of C2 (Top left). Similar, but less extensive, bone loss is seen at C3 and C4.

Figure 3. Post-operative radiographs from AP and lateral views one year after circumferential fusion.

Figure 4. Post-operative computed tomography imaging in the sagittal plane 1 year post-op. A solid fusion is present throughout. Rib struts have reconstructed the large bony defect in the upper cervical spine and incorporated from the occiput to C5.

Accepted Manuscript