



Aggressive thoracic vertebral lesion with intraosseous and epidural components

A 51-year-old woman presented with history of low back-ache and progressive spastic paraparesis of 1 year's duration. Magnetic resonance imaging of the thoracic spine (Fig. 1) demonstrated an enhancing lesion in the T8 body that was hyperintense on T1 and T2 images and had prominent secondary trabeculations. These signal changes were seen extending into the lamina, pedicles, and the articular pro-

cesses. There was an associated soft tissue component seen in the paravertebral and epidural region compressing the cord at this level. Computed tomography (Fig. 2A, B) images revealed a decreased T8 vertebral body height and multiple lucencies, coarse trabeculations, and a “polka-dot” appearance of the body. There was a breach in the posterior cortex of the body. Spinal angiogram (Fig. 2C) showed a tumor blush arising from the bilateral T8 segmental artery feeders. Following an endovascular embolization of the lesion, the patient underwent a T8 corpectomy, excision of the epidural component, and insertion of a cage and pedicle screws (Fig. 2D). Histopathology confirmed the lesion to be a vertebral hemangioma.




Fig. 1. MRI of the thoracic spine: (A) Sagittal T1 sequence demonstrating a heterogeneous lesion within the T8 body and a soft tissue component in the spinal canal. (B) The lesion is noted to be hyperintense on T2-weighted sequences. (C) Enhancement with contrast is noted on gadolinium-enhanced T1 sequences. (D) Axial T2 sequence demonstrating the posterior extent of the lesion.




Fig. 2. CT thoracic spine showing (A) lytic change in the T8 body; (B) “polka-dot” appearance of the body and multiple lucencies; (C) spinal angiogram showing the tumour blush; and (D) postoperative CT showing the cage and pedicle screws in situ.

Sumit Thakar, MCh^a

Narayanam Anantha Sai Kiran, MCh^a

Dilip Mohan, MCh, DNB^a

B.T. Pushpa, DNB^b

Vinay Hegde, MD^b

Alangar S. Hegde, MCh, PhD^a

^aDepartment of Neurosurgery

Sri Sathya Sai Institute of Higher Medical Sciences

EPIP Area, Whitefield

Bangalore 560066, Karnataka State, India

^bDepartment of Radiology

Sri Sathya Sai Institute of Higher Medical Sciences

EPIP Area, Whitefield

Bangalore 560066, Karnataka State, India

FDA device/drug status: Not applicable.

Author disclosures: **ST**: Nothing to disclose. **NASK**: Nothing to disclose. **DM**: Nothing to disclose. **BTP**: Nothing to disclose. **VH**: Nothing to disclose. **ASH**: Nothing to disclose.

The authors do not report any conflict of interest concerning the materials and methods used in this study or the findings specified in this paper.

Source of support: The authors have not received any support in the form of grant from any source for preparation of this paper or in the course of the study.

This paper has not been published before and has not been submitted for publication to any other journal in part or full. It has not been presented in any meeting/conference. This paper has been prepared using guidelines laid down by the Institute.