

Accepted Manuscript

Title: Transoral vertebroplasty for a c2 aneurysmal bone cyst

Author: Liberto Brage, Héctor Roldán, Julio Plata-Bello, Martel Diego, Víctor García-Marín

PII: S1529-9430(16)00432-0

DOI: <http://dx.doi.org/doi: 10.1016/j.spinee.2016.02.042>

Reference: SPINEE 56930

To appear in: *The Spine Journal*

Received date: 6-9-2015

Revised date: 28-1-2016

Accepted date: 23-2-2016



Please cite this article as: Liberto Brage, Héctor Roldán, Julio Plata-Bello, Martel Diego, Víctor García-Marín, Transoral vertebroplasty for a c2 aneurysmal bone cyst, *The Spine Journal* (2016), <http://dx.doi.org/doi: 10.1016/j.spinee.2016.02.042>.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Title

Transoral vertebroplasty for a C2 Aneurysmal Bone Cyst

Authors

Liberto Brage¹, Héctor Roldán¹, Julio Plata-Bello¹, Diego Martel¹, Víctor García-Marín¹.

Affiliation

¹ Hospital Universitario de Canarias (Department of Neurosurgery), S/C de Tenerife, Spain.

³ Hospital Universitario de Canarias (Department of Neurology), S/C de Tenerife, Spain.

Corresponding author

Julio Plata Bello

Address: Hospital Universitario de Canarias (Neuroscience department), Calle Ofra s/n La Cuesta. CP 38320. La Laguna, S/C de Tenerife. Spain. Phone number: +34 922 255 544 / +34 646 625 973. E – mail address: jplata5@hotmail.com

ABSTRACT

BACKGROUND: Aneurysmal bone cysts at the cervical spine represent a real challenge both diagnostically and therapeutically, especially in young patients.

PURPOSE: To present an unusual case of a C2 aneurysmal bone cyst expanding the body in its entirety in a young adult successfully treated with a transoral vertebroplasty.

STUDY DESIGN: Case Report.

METHODS: We report the case of a 17 year-old woman with a history of cervical pain and occipital headache after a car accident. Routine X-rays disclosed a C2 lesion. Her neurologic examination was normal. Computed tomography showed a lytic lesion occupying almost the entire body of the C2 vertebra. The cortical bone was intact but notably thinned. MRI imaging revealed a cystic image with blood inside. Transoral vertebroplasty was selected among other surgical options for the following reasons: 1) to improve the clinical symptoms, and 2) to prevent future vertebral collapse with devastating neurological consequences. Under general anesthesia and continuous neurophysiological monitoring, we conducted a fluoroscopic-guided transoral vertebroplasty through a Jamshidi needle. A cytology sample from the cystic lesion was taken through the needle.

RESULTS: The blood smear showed no tumoral cellularity. There were no complications during surgery or postoperative infections. After 4 years of follow-up, the patient is pain-free and leads a normal life.

CONCLUSIONS: Transoral vertebroplasty seems to be a direct, safe, and effective technique to stabilize cystic lesions that endanger the stability of C2 and to improve symptoms. Aneurysmal bone cysts should be included in the differential diagnosis of lytic lesions at the vertebral body of C2.

Keywords: Axis, Aneurysmal Bone Cyst, Stability, Transoral, Vertebroplasty, Antibiotics.

INTRODUCTION

Aneurysmal bone cysts (ABC), representing 1.4% of all primary bone tumors and 15% of all primary spine tumors [1], are a rare benign condition, sometimes self-limiting, and generally located in the posterior arch of lumbar vertebrae, followed by the thoracic, cervical, and sacral segments [2]. Despite their benign nature, they can be locally aggressive, and can result in pathological fractures and neurological complications [3]. They can appear as primary bone lesions (in 70 % of the cases) or secondary (in about 30 %), when ABC-like areas are encountered inside other bone conditions (giant cell tumors, chondroblastoma, telangiectatic osteosarcoma, osteoblastoma) [4]. Although not pathognomonic, ABC are usually diagnosed based on their typical radiological appearance, consisting of multilocular cysts with fluid-fluid levels on MRI and bony septations on computed tomography [5].

The treatment of ABC is controversial. Multiple treatment modalities have been tried with variable improvement and recurrence rates. The options include curettage with or without bone grafting, complete excision, arterial embolization, intralesional drug injections (steroid and calcitonin) and radiation [6]. Intralesional curettage is the technique most often used in the spine [7,8]. Nevertheless, complete excision is the optimal approach to achieve local control of the tumor as it prevents recurrence, but this treatment exposes the patient to high surgical morbidity [9,10]. Radiotherapy has been indicated in unresectable lesions, as it has numerous and severe complications [11]. Arterial embolization has been used preoperatively to decrease surgical bleeding and also as a stand-alone approach for inoperable cases [12].

We report an unusual case of an ABC at C2 affecting the entire vertebral body successfully treated with transoral cementation in a young adult, in addition to discussing the rationale for the selected treatment.

METHODS

Our case is a 17 year-old female who reported a long history of cervical pain and occipital headaches. She had previously sought out medical attention, but no clear diagnosis had been achieved. She was subsequently involved in a car accident and sustained a whiplash injury. On routine cervical X-rays, a radiolucent lesion in the vertebral body of C2 was found (Fig. 1A). For this reason she was referred to our hospital. On examination, the only physical finding was neck pain without any neurological deficits. A computed tomography scan (CT-scan) showed a lytic expansile lesion occupying almost the entire body of C2 with intact but notably thinned anterior and posterior cortical boundaries. Multiple partitions could be seen inside the cavity (Fig. 1B,C). Magnetic resonance imaging (MRI) revealed a cystic lesion containing blood (Fig. 1C).

Given the symptoms, patient age and structural compromise of the integrity of C2 by the lytic lesion, we decided to perform a transoral vertebroplasty with three main objectives: 1) to improve the clinical symptoms, 2) to prevent future vertebral collapse with devastating neurological consequences, and 3) to obtain a pathological sample of the content of the lesion.

The surgery was performed under general anesthesia and continuous neurophysiological monitoring. The patient underwent systemic and local antibiotic prophylaxis in the oral cavity. The patient was placed supine with the head hyperextended and a Dingman mouth gag was used. The inner part of the mouth was draped. The entry point was located in the throat with the aid of biplanar fluoroscopy (Fig. 2A). A Jamshidi needle was used to reach the lesion allowing the evacuation of the hematic content, which was sent to the Pathology Department for postoperative diagnosis (Fig. 2B). We then performed the fluoroscopy-guided vertebroplasty by introducing approximately 1 cc of PMMA (Confidence Spinal Cement System™, DePuy Synthes, Inc., Warsaw, IN) (Fig. 2C), which easily filled in all the cavities. Once the final result was verified, the Jamshidi needle was withdrawn and the pharynx was inspected. A small bleeding point was controlled by applying low pressure with gauze. The procedure lasted

an hour and a half, and the patient awoke from anesthesia without complications. Oral rinses with chlorhexidine and intravenous cefazolin were prescribed for 72 hours after surgery, and no local or systemic infection signs occurred. A plain cervical radiography was taken confirming the integrity of both cortical boundaries and the patient was discharged from hospital after three days of admission.

The blood smear was normal without any tumoral cells. This confirmed our diagnosis of ABC.

After four years of follow-up, the patient is pain-free and leads a normal life. Plain X-Rays, CT-scan, and MRI show no signs of recurrence or progression of the C2 lesion (Fig. 3A,B,C).

DISCUSSION

To our knowledge this is the second case reported of a transoral cementation of an ABC at C2 [16]. This article describes the case of a 58 year-old male who had a C2 fracture secondary to a lytic lesion, treated with vertebroplasty through a transoral route. Our case is the first one in a young adult.

The main pathological feature of ABC is the presence of many blood cavities separated by thin fibrous septa. This fact explains the radiological image of a great cavity with many hollows in the interior filled with blood [9]. Our case showed, on both MRI and CT-scan, the typical radiological appearance of an ABC. MRI demonstrated a hyperintense signal at T2WI consistent with blood content (Fig. 1C), while the CT-scan showed the extent of the bone destruction with integrity of the cortical boundaries and several cavities filled with fluid (Fig. 1B,C). Blood was aspirated from the vertebral body of C2 through the Jamshidi needle. Pathological analysis was normal without any tumoral cells, which was therefore consistent with ABC [20].

Instability is fairly clear in literature when there is a fracture at C2 or a

C1-C2 or C2-C3 subluxation. Moreover, acute cord compression is reported in the literature in the absence of complete vertebral collapse [21]. We decided to indicate an invasive treatment in this case because the patient was young, active, symptomatic, and the very thin outer walls of the C2 body posed a significant risk of fracture with potentially catastrophic consequences. Stand-alone embolization and infiltration of fibrosing agents have been advocated for treating ABC [6,12]. However, these procedures were discarded from the beginning as none of them stabilizes C2 and the presence of very thin walls might not contain fluids from extravasation into the spinal canal. What to do from this point was a matter of thoughtful debate.

Open surgery of cervical ABC implies a combined approach. The anterior approach is used to accomplish the curettage, electrocautery and bone grafting, whereas the posterior approach is necessary to stabilize the segment [20]. Usually, at C2, the posterior stabilization requires an occipitocervical fusion [20]. The disadvantages include being a double procedure with a fairly complicated postoperative course and a high risk of intraoperative bleeding [22,23]. In fact, many authors advise to carry out a selective arterial embolization to prevent hemorrhage [12,20]. We also considered this patient as a high-risk case for curettage, as the boundaries were very thin and easy to pierce, with eventual neurological consequences.

In our case, as ABC show huge cavities filled with blood, vertebroplasty seemed to be an option. Sometimes bone septa of ABC are not complete, and the cavities are separated only by thin fibrous walls that can be torn by pressurized cement, so one injection can be enough to fill in all the cavities. Nevertheless, even though vertebroplasty at C2 seems to represent an important tool to control the pain and to prevent vertebral collapse and spinal cord compression, only a small number of cases have been published [14,15,17,18,19,24,25]. The advantages over open procedures include a quicker and easier method without an external incision in a young patient [26]. Percutaneous vertebroplasties at the thoracic and lumbar vertebral bodies are usually performed through a transpedicular route. At the subaxial cervical spine, an anterolateral oblique

approach is typically used [15]. To the best of our knowledge, the transpedicular approach to reach the body of C2 is not a safe option due to the proximity of the vertebral arteries and the small size of the pedicles [19]. The most straightforward access is provided by the transoral route, which has the advantage of the short distance to the C2 vertebral body, thereby avoiding important vascular structures [15,16]. The most significant drawback is crossing a naturally contaminated area. Infection rates for open transoral procedures vary widely from 0% to 50%, depending on preoperative site preparation, prophylactic antibiotic use, duration of the procedure, and the extent of tissue disruption [13]. In our patient we did not have any infectious complications. We provided systemic and local antibiotic prophylaxis, performing a rapid procedure through a minimum puncture in the oropharynx and were very careful not to touch the rest of the oral cavity with the Jamshidi needle (we draped the inner part of the mouth to prevent this). Cefazolin was continued for 72 hours after surgery until discharge. Daily pharyngeal inspection was conducted until discharge, and no signs of infection were noted. Some authors have described mixing PMMA with local antibiotics [27]. We did not do this as we were not certain about the effect of using a different substance on PMMA final consistency, and we were concerned about PMMA leakage at such a critical level as C2.

To accomplish the percutaneous vertebroplasty in our patient we selected polymethyl-methacrylate (PMMA) as opposed to biological cement for several reasons. Firstly, the use of biological cement in complex cases is not well known, with only a handful of papers describing the long-term effects of these materials, especially in case of leakage [28]. Secondly, even with integrity of cortical boundaries, we have more confidence in high-viscosity cement, as we do know how it changes over time until it becomes dough so it can be safely applied under radiological guidance [29]. Thirdly, the exothermic polymerization process that occurs with PMMA, reaching temperatures superior to the critical level for protein denaturation, can provide a tumoricidal effect in a lesion with high rates of recurrence [30].

1 As stand-alone cementation is not an established treatment for ABC, the
2 ability to achieve local control of the lesion cannot be predicted beforehand, so
3 patients must be followed up carefully. Nevertheless, cases of resolution of ABC
4 after minor intralesional procedures have been published, which probably
5 reinforces the nontumoral nature of these lesions [2].

6 7 8 **CONCLUSION**

9
10 Transoral vertebroplasty may constitute a safe and effective treatment to
11 easily stabilize cystic lesions that compromise the stability of C2. ABCs affecting
12 the vertebral body need to be considered when making the differential diagnosis
13 of cystic lesions at C2 in young adults.

1 REFERENCES

- 2
3
4 1. Ameli NO, Abbassioun K, Saleh H, et al. Aneurysmal bone cysts of the spine.
5 Report of 17 cases. J Neurosurg 1985;63:685–90.
- 6 2. Liu JK, Brockmeyer DL, Dailey AT, Schmidt MH. Surgical management of
7 aneurysmal bone cysts of the spine. Neurosurg Focus 2003;15:E4.
- 8 3. Burch S, Hu S, Berven S. Aneurysmal bone cysts of the spine. Neurosurg Clin N
9 Am 2008;19:41–7.
- 10 4. Boriani S, De Iure F, Campanacci L, Gasbarrini A, Bandiera S, Biagini R, et al.
11 Aneurysmal bone cyst of the mobile spine. Report of 41 cases. Spine
12 2001;26:27–35.
- 13 5. Kitamura T, Ikuta K, Senba H, Komiya N, Shidahara S. Long-term follow-up of
14 a case of aneurysmal bone cyst in the lumbar spine. Spine J 2013;13:e55-8.
- 15
16 6. Saccomanni B. Aneurysmal bone cyst of spine: a review of literature. Arch
17 Orthop Trauma Surg 2008;128:1145–7.
- 18 7. Boriani S, Lo SF, Puvanesarajah V, Fisher CG, Varga PP, Rhines LD, et al;
19 AOSpine Knowledge Forum Tumor. Aneurysmal bone cysts of the spine:
20 treatment options and considerations. J Neurooncol 2014;120:171-8.
- 21 8. Sebaaly A, Ghostine B, Kreichati G, Mallet JF, Glorion C, Moussa R, et al.
22 Aneurysmal Bone Cyst of the Cervical Spine in Children: A Review and a Focus
23 on Available Treatment Options. J Pediatr Orthop 2015;35:693-702.
- 24 9. Papagelopoulos PJ, Currier BL, Shaughnessy WJ, Sim FH, Ebersold MJ, Bond
25 JR, et al. Aneurysmal bone cyst of the spine. Management and outcome. Spine
26 1998;23:621–8.
- 27 10. Boriani S, Bandiera S, Donthineni R, Amendola L, Cappuccio M, De Iure F, et
28 al. Morbidity of en bloc resections in the spine. Eur Spine J 2010;19:231–41.
- 29 11. Vergel de Dios AM, Bond JR, Shives TC, McLeod RA, Unni KK. Aneurysmal

- bone cyst. A clinicopathologic study of 238 cases. *Cancer* 1992;69:2921–31.
12. Rossi G, Rimondi E, Bartalena T, Gerardi A, Alberghini M, Staals EL, et al. Selective arterial embolization of 36 aneurysmal bone cysts of the skeleton with N-2-butyl cyanoacrylate. *Skeletal Radiol* 2010;39:161–7.
13. Sachs DC, Inamasu J, Mendel EE, Guiot BH. Transoral vertebroplasty for renal cell metastasis involving the axis: case report. *Spine* 2006;34:E925-8.
14. Mont'Alverne F, Vallée JN, Cormier E, Guillevin R, Barragan H, Jean B, et al. Percutaneous vertebroplasty for metastatic involvement of the axis. *AJNR Am J Neuroradiol* 2005;26:1641-5.
15. Martin JB, Gailloud P, Dietrich PY, Luciani ME, Somon T, Sappino PA, et al. Direct transoral approach to C2 for percutaneous vertebroplasty. *Cardiovasc Intervent Radiol* 2002;25:517-9.
16. Gailloud P, Martin JB, Olivi A, Rüfenacht DA, Murphy KJ. Transoral vertebroplasty for a fractured C2 aneurysmal bone cyst. *J Vasc Interv Radiol* 2002;13:340-1.
17. Dorman JK. Vertebroplasty of the C2 vertebral body and dens using an anterior cervical approach: technical case report. *Neurosurgery* 2010;67:E1143-6.
18. Yoon JY, Kim TK, Kim KH. Anterolateral percutaneous vertebroplasty at C2 for lung cancer metastasis and upper cervical facet joint block. *Clin J Pain* 2008;24:641-6.
19. Monterumici DA, Narne S, Nena U, Sinigaglia R. Transoral kyphoplasty for tumors in C2. *Spine J* 2007;7:666-70.
20. Levin DA, Hensinger RN, Graziano GP. Aneurysmal bone cyst of the second cervical vertebrae causing multilevel upper cervical instability. *J Spinal Disord Tech* 2006;19:73-5.
21. Chan MS, Wong YC, Yuen MK, Lam D. Spinal aneurysmal bone cyst causing acute cord compression without vertebral collapse: CT and MRI findings. *Pediatr Radiol* 2002;32:601-4.

- 1 22. Mehdian H, Weatherley C. Combined anterior and posterior resection and spinal
2 stabilization for aneurysmal bone cyst. *Eur Spine J* 1995;4:123-5.
- 3 23. Nicastro JF, Leatherman KD. Two-stage resection and spinal stabilization for
4 aneurysmal bone cyst. A report of two cases. *Clin Orthop Relat Res*
5 1983;(180):173-8.
- 6 24. Lieberman I, Reinhardt MK. Vertebroplasty and kyphoplasty for osteolytic
7 vertebral collapse. *Clin Orthop Relat Res* 2003;(415 Suppl):S176-86.
- 8 25. Fourney DR, Schomer DF, Nader R, Chlan-Fourney J, Suki D, Ahrar K, et al.
9 Percutaneous vertebroplasty and kyphoplasty for painful vertebral body
10 fractures in cancer patients. *J Neurosurg* 2003;98(1 Suppl):21-30.
- 11 26. Bonney G, Williams JP. Trans-oral approach to the upper cervical spine. A
12 report of 16 cases. *J Bone Joint Surg Br* 1985;67:691-8.
- 13 27. Webb JC, Spencer RF. The role of polymethylmethacrylate bone cement in
14 modern orthopaedic surgery. *J Bone Joint Surg Br* 2007;89:851-7.
- 15 28. Martínez-Quñones JV, Medina-Imbroda JM, Aso-Escario J, Pons Bosque J,
16 Consolini FE, Arregui-Calvo R. [Biological cement extravasation into the spinal
17 canal following kyphoplasty. Long-term follow-up. Report of one case and
18 review of the literature]. *Neurocirugia (Astur)* 2011;22:588-99.
- 19 29. Georgy BA. Clinical experience with high-viscosity cements for percutaneous
20 vertebral body augmentation: occurrence, degree, and location of cement
21 leakage compared with kyphoplasty. *AJNR Am J Neuroradiol* 2010;31:504-8.
- 22 30. Malawer MM, Dunham W. Cryosurgery and acrylic cementation as surgical
23 adjuncts in the treatment of aggressive (benign) bone tumors. Analysis of 25
24 patients below the age of 21. *Clin Orthop Relat Res* 1991;(262):42-57.

FIGURE LEGENDS

Fig. 1. Preoperative images: (A) Plain radiography showing a lytic lesion at C2 body. (B), (C) Sagittal and axial CT-scan demonstrating an expansile osteolytic lesion. Note the integrity of the thin cortical boundaries and multiple partitions in its interior. (D) Preoperative T2WI MRI showing blood content inside the lesion.

Fig. 2. Intraoperative images: (A) Image of the operating room showing patient's position under biplanar fluoroscopy. (B) Blood coming out of the lesion through the Jamshidi needle. No pathological cells were found postoperatively. (C) Fluoroscopic image depicting the Jamshidi needle inside the lesion and the cement filling the cavities.

Fig. 3. Postoperative images: (A), (B) Plain radiographs showing the final result of the cementation. (C) CT-scan demonstrating that the lesion is completely full of cement. No extravasation of cement occurred.