



Gorham's disease—vanishing bone of the cervical spine

A 25-year-old male patient presented with neck pain. Cervical spine magnetic resonance imaging demonstrated a mass

isointense to muscle on T1- and T2-weighted images with moderate enhancement in the C6 posterior elements (Figs. 1 and 2). A small right scapula cystic lesion was also seen (Fig. 2). Computed tomography revealed the C6 lesion as a destructive vanishing bone tumor, as well as small lucent lesions in the vertebrae, right scapula, and second right




Fig. 1. Sagittal T2-weighted MR images of the cervical spine through the level of C6 (Left, Middle, Right) showed a mass isointense to muscle on T1- and T2-weighted images with moderate enhancement (arrows).




Fig. 2. The mass (arrowheads) was isointense to muscle on T2-weighted axial MR image (Left) and showed moderate enhancement on postcontrast axial T1-weighted MR image (Middle). A small right scapula cystic lesion (arrow) was also seen (Right).



Fig. 3. Sagittal reformatted CT image (A) showed a destructive vanishing bone tumor in the C6 posterior elements (arrow). Coronal reformatted CT images showed small lucent lesions (arrows) in the right scapula (B), vertebra (C), and second right anterior costa (D).




Fig. 4. Axial CT scans showed a destructive vanishing bone tumor in the C6 vertebra (A), lucent lesions in the T1 vertebra (B), and second right anterior costa (C) (arrows). Fat density in the second right anterior costa lesion suggested benignity. A very low density mass compatible with lymphangioma was found in the anterior mediastinum (D) (arrow).

anterior costa (Figs. 3 and 4). A fat density in the second right anterior costa lesion suggested benignity (Fig. 4). A low density mass compatible with lymphangioma was found in the anterior mediastinum (Fig. 4). The radiological diagnosis was Gorham's disease. A resection of the C6 vertebra lesion was performed and pathology demonstrated an angiomatous lesion.

Gorham's disease (idiopathic massive osteolysis, vanishing bone disease, or disappearing bone disease), an aggressive form of osseous angiomatosis, is very rare in the spine [1].

Reference

- [1] Barman A, Bhide R, Viswanathan A, George J, Thomas R, Tharion G. Gorham's disease of the spine. *Neurorehabilitation* 2013;33:121–6.

Mustafa Kemal Demir, MD^a

Özlem Yapıcıer, MD^b

Zafer Orkun Toktaş, MD^c

Baran Yılmaz, MD^c

Akın Akakın, MD^c

Deniz Konya, MD^c

^aDepartment of Radiology

Bahçeşehir University School of Medicine

Göztepe Medical Park Hospital, 11. kısım, Yasemin Apt
D blok. Daire 35 Ataköy-Bakırköy, İstanbul 34158, Turkey

^bDepartment of Pathology

Acıbadem University School of Medicine

Kerem Aydınlar Kampüsü, Patoloji Ana Bilim Dalı
Kayışdağı Cad. No:32 Ataşehir, İstanbul 34752, Turkey

^cDepartment of Neurosurgery

Bahçeşehir University School of Medicine

Göztepe Medical Park Hospital

Göztepe Medikal Park Hastanesi, Nöroşirürji Kliniği E5

Üzeri 23 Nisan Sok. No: 17 Merdivenköy Kadıköy
İstanbul 34732, Turkey

FDA device/drug status: Not applicable.

Author disclosures: **MKD**: Nothing to disclose. **ÖY**: Nothing to disclose. **ZOT**: Nothing to disclose. **BY**: Nothing to disclose. **AA**: Nothing to disclose. **DK**: Nothing to disclose.

Conflict of interest: The authors declare that they have no conflict of interest.