

Accepted Manuscript

Diastematomyelia with the owl sign (Type I split cord malformation)

Ayşe Gul Alimli, MD, Cigdem Oztunali, MD, Oznur Leman Boyunaga, MD, Betül Emine Derinkuyu, MD, Cagrı Damar, MD, Ayfer Aslan, MD, Murat Ucar, MD



PII: S1529-9430(15)00547-1

DOI: [10.1016/j.spinee.2015.05.025](https://doi.org/10.1016/j.spinee.2015.05.025)

Reference: SPINEE 56347

To appear in: *The Spine Journal*

Received Date: 14 May 2015

Accepted Date: 19 May 2015

Please cite this article as: Alimli AG, Oztunali C, Boyunaga OL, Derinkuyu BE, Damar C, Aslan A, Ucar M, Diastematomyelia with the owl sign (Type I split cord malformation), *The Spine Journal* (2015), doi: 10.1016/j.spinee.2015.05.025.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Diastematomyelia with the owl sign (Type I split cord malformation)

Ayşe Gul Alimli, MD¹, Cigdem Oztunali, MD¹, Oznur Leman Boyunaga, MD¹,
Betül Emine Derinkuyu, MD¹, Çağrı Damar MD¹, Ayfer Aslan MD², Murat
Ucar MD³

¹Gazi University, School of Medicine, Department of Pediatric Radiology,
Ankara, Turkey

²Gazi University, School of Medicine, Department of Neurosurgery, Ankara,
Turkey

³Gazi University, School of Medicine, Department of Radiology, Ankara,
Turkey

*Corresponding author:

Ayşe Gul Alimli, MD

Gazi University, School of Medicine

Department of Pediatric Radiology

Besevler, 06510

Ankara, Turkey

+90 505 441 9314

drsahin80@hotmail.com

Diastematomyelia with the owl sign (Type I split cord malformation)

A 4-year old female patient was admitted our Neurosurgery Department with the complaints of weakness in her left leg, difficulty in walking and falling down frequently while walking. She had a sacral dimple since her birth but it was not investigated. Physical examination revealed sacral dimple, scoliosis and slight dorsal flexion of the left foot. Magnetic resonance imaging (MRI) of the spine was performed. MR images showed thoracolumbar levoscoliosis. Additionally, syringomyelia was present throughout the T5-T7 vertebrae (**Figure 1**) and the spinal cord was tethered with a low-lying conus medullaris at L4 level. MR images showed congenital fusions of the T7-T8 and T12-L1 vertebral bodies (**Figure 2**). Vertebral anomalies including hemivertebrae and butterfly vertebrae at the level of T9-11 were observed (**Figure 1-2**). Spinal canal was ectatic at the lower thoracic and lumbar levels (**Figure 3**). There was diastematomyelia with a bony spur continuing throughout the T 11- L2 levels (the owl sign) (**Figure 3**). At these levels, the spur in the spinal canal divided the spine into two hemicords (Type I split cord malformation with two dural sacs) (**Figure 3**). A T11-12 laminectomy with the removal of the bony spur to repair the diastematomyelia was performed. Dural tubes were then repaired and a common thecal sac was subsequently formed. A coccygeal dermal sinus tract that is not extending into the spinal canal was observed during the operation and the tract was excised. The patient was discharged uneventfully following the surgery.

Figure Legends

Figure 1: Sagittal T2 weighted MR image shows syringomyelia (black arrows) and vertebral anomalies (white arrows).

Figure 2: Sagittal T2 weighted MR image shows congenital fusion of the T12-L1 posterior vertebral bodies (white arrow) and vertebral anomalies (black arrows).

Figure 3: Axial (right) and coronal (left) T2 weighted MR images show diastematomyelia with a bony spur (asterix) (the owl sign) and two hemicords (black arrows) with the ectatic spinal canal (white arrows) .





