

# Accepted Manuscript

Title: Langerhans cell histiocytosis of the cervical spine

Author: Nancy Abu-Bonsrah, C. Rory Goodwin, Genevieve M. Crane, Godwin Abiola, Daniel M. Sciubba

PII: S1529-9430(15)01220-6

DOI: <http://dx.doi.org/doi: 10.1016/j.spinee.2015.08.010>

Reference: SPINEE 56516

To appear in: *The Spine Journal*



Please cite this article as: Nancy Abu-Bonsrah, C. Rory Goodwin, Genevieve M. Crane, Godwin Abiola, Daniel M. Sciubba, Langerhans cell histiocytosis of the cervical spine, *The Spine Journal* (2015), <http://dx.doi.org/doi: 10.1016/j.spinee.2015.08.010>.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

## Langerhans Cell Histiocytosis of the Cervical Spine

Nancy Abu-Bonsrah<sup>1</sup> BS, C. Rory Goodwin<sup>1</sup>, MD, PhD, Genevieve M. Crane<sup>2</sup>, MD,  
PhD, Godwin Abiola<sup>1</sup>, BS, Daniel M. Sciubba<sup>1</sup>, MD

<sup>1</sup>Departments of Neurosurgery and <sup>2</sup>Pathology, The Johns Hopkins University School of  
Medicine,  
Baltimore, MD, USA.

**Disclosure:** CRG is a UNCF Merck Postdoctoral Fellow and has received an award from the Burroughs Wellcome Fund. This article reflects the views of the author CRG and should not be construed to represent the FDA's views or policies. NAB, GMC and GA have no disclosures. DS is a consultant for DePuy, Medtronic, Orthofix, and Globus.

The authors report no external source of funding for this manuscript.

Corresponding author:  
C. Rory Goodwin, MD  
600 North Wolfe Street  
Meyer 5-185  
Baltimore, MD 21287, USA  
Phone 410-955-4424  
Fax: 410 502 5768  
rory@jhmi.edu

# 1 Manuscript Text

2 31-year-old male with history of diffuse Langerhans cell histiocytosis (LCH) involving  
 3 the left mastoid base status post resection and radiation presented with progressive severe  
 4 right-sided neck pain. CT and MRI revealed an osteolytic lesion at C5 (Figure 1) and due  
 5 to concern for mechanical instability, the patient underwent a two-stage anterior C5  
 6 corpectomy and posterior cervical fusion with instrumentation (Figure 2). Final  
 7 pathology of the surgical tissue confirmed LCH (Figure 3). Post-operatively, the patient's  
 8 pain improved and he continues to undergo chemotherapy for his systemic disease. LCH  
 9 is a rare disease commonly affecting the bones and spinal involvement occurs in 6.5-25%  
 10 of all skeletal cases with 11% affecting the cervical spine<sup>1</sup>. Patients commonly present  
 11 with localized neck pain, restricted range of motion or neurological deficits<sup>2-4</sup>. Imaging  
 12 frequently reveals an osteolytic lesion, though osteoblastic lesions can occur<sup>2</sup>. LCH of the  
 13 spine may be treated conservatively with observation, immobilization, NSAIDS and  
 14 casting or with radiation, steroids or chemotherapy—especially in systemic disease or  
 15 multifocal spine disease<sup>1-2, 4-5</sup>. Surgical intervention—i.e. curettage with or without bone  
 16 grafting, internal fixation and fusion or percutaneous vertebroplasty—may be indicated in  
 17 cases of severe mechanical instability, deformity and/or neurological deficits caused by  
 18 compression<sup>1, 2</sup>.

19

20 **Figure 1:** MRI (A) and CT (B) imaging showing an expansile lytic lesion associated with  
 21 erosive changes within the C5 vertebral body

22 **Figure 2:** Post-operative CT imaging, with and without contrast, showing stable  
 23 corpectomy at C5 with posterior fusion of C4-C6

**Figure 3:** Histopathology of LCH involving the cervical spine. A histiocytic lesion with numerous admixed eosinophils is seen diffusely involving the vertebral body and partially eroding bone (A, H&E, 20X lens, bone indicated by ‘\*’). The histiocytic cells were strongly CD1A positive (B, immunoperoxidase, 20X lens). Closer examination of the neoplastic cells revealed characteristic Langerhans cells with elongated and folded nuclei (C, H&E, 40X, arrow indicating cells with more prominent nuclear features).

## References

1. Jiang L, Liu ZJ, Liu XG, et al. Langerhans cell histiocytosis of the cervical spine: a single Chinese institution experience with thirty cases. *Spine*. 2010;35(1):E8-15.
2. Feng F, Tang H, Chen H, Jia P, Bao L, Li JJ. Percutaneous vertebroplasty for Langerhans cell histiocytosis of the lumbar spine in an adult: Case report and review of the literature. *Exp Ther Med*. 2013;5(1):128-132.
3. Chaudhary V, Bano S, Aggarwal R, et al. Neuroimaging of Langerhans cell histiocytosis: a radiological review. *Jpn J Radiol*. 2013;31(12):786-96.
4. Sayhan S, Altinel D, Erguden C, Kizmazoglu C, Guray M, Acar U. Langerhans cell histiocytosis of the cervical spine in an adult: a case report. *Turk Neurosurg*. 2010;20(3):409-12.
5. Cantu MA, Lupo PJ, Bilgi M, Hicks MJ, Allen CE, McClain KL. Optimal therapy for adults with Langerhans cell histiocytosis bone lesions. *PLoS ONE*. 2012;7(8):e43257.