

Accepted Manuscript

Title: Lumbar pedicle lesion causing radicular pain

Author: Laxminadh Sivaraju, Saritha Aryan, Narayanam Anantha Sai Kiran,
Alangar S Hegde

PII: S1529-9430(15)01213-9

DOI: <http://dx.doi.org/doi: 10.1016/j.spinee.2015.08.003>

Reference: SPINEE 56509

To appear in: *The Spine Journal*



Please cite this article as: Laxminadh Sivaraju, Saritha Aryan, Narayanam Anantha Sai Kiran, Alangar S Hegde, Lumbar pedicle lesion causing radicular pain, *The Spine Journal* (2015), <http://dx.doi.org/doi: 10.1016/j.spinee.2015.08.003>.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Title page**Title: Lumbar pedicle lesion causing radicular pain**

Article Type: Special feature: Images of spine care

Authors

Laxminadh Sivaraju, M.Ch

Email: Laxminadh.sivaraju@gmail.com

Saritha Aryan, M.Ch

Email: saaryan@gmail.com

Narayanam Anantha Sai Kiran, M.Ch

Email: Email: saikiransssihms@gmail.com

Alangar S Hegde, M.Ch, PhD

Email: alangars@rediffmail.com

First Author: Laxminadh Sivaraju, M.Ch

Order of Authors: Laxminadh Sivaraju, M.Ch; Saritha Aryan, M.Ch; Narayanam Anantha Sai Kiran,

M.Ch; Alangar S Hegde, M.Ch,Phd

Department and institution

Departments of Neurosurgery

Sri Sathya Sai Institute of Higher Medical
Sciences, Whitefield, Bangalore- 560066, India

Corresponding Author

Dr. Laxminadh Sivaraju,
Department of Neurosurgery,
Sri Sathya Sai Institute of Higher Medical
Sciences, Whitefield, Bangalore- 560066, India.

Email: laxminadh.sivaraju@gmail.com

Phone: 91 8884436640

Fax: 8411503

Source of support: The authors have not received any support, in the form of grant, from any source for preparation of this article. Neither do the authors, nor does the institute have any personal or institutional financial interest in drugs, materials, or devices described in their submissions.

Disclosure: This paper has not been published before and has not been submitted for publication to any other journal in part or full. The authors concur with this submission and there is no conflict of interest arising from this article.

1
2
3
4
5
6
7
8 A twenty six year old male presented with history of right lower limb radicular
9 pain, parasthesias in right thigh and back ache for two years. He had antalgic
10 gait and lumbosacral scoliosis with wasting of paraspinal muscles on the right
11 side. Computed tomography (CT) showed a lytic lesion with surrounding
12 sclerosis in the right L3 pedicle. There was soft tissue mass lesion arising
13 inferomedially and causing foraminal stenosis (Figure 1). On magnetic
14 resonance (MR) imaging it was hypointense in both T1 and T2-weighted
15 images measuring 2.3X2 cm. There was moderate contrast enhancing soft tissue
16 extending into right L3-4 neural foramen compressing the exiting root. There
17 was marrow edema seen in short tau inversion recovery (STIR) images (Figure
18 2). He underwent right L3 hemilaminectomy and excision of the tumor. It had
19 both bony and soft tissue components. L3 inferior facet was excised.
20 Postoperative period was uneventful and radicular pain subsided. Postoperative
21 dynamic X-rays showed no instability. He was advised to follow up and was
22 explained about requirement of instrumentation in future if instability occurs.
23 The lesion was sent for histopathological examination. Histopathology reported

as osteoid formation with intervening fibrovascular stroma. There were foci of osteoclastic giant cells with hemosiderin laden macrophages. Overall features were consistent with osteoblastoma, a benign tumor resulting in remodelling of osteoid tissue and bone[1]. At ten months followup the patient was doing well.

References

[1]. Li Z, Zhao Y, Hou S, Mao N, Yu S, Hou T. Clinical features and surgical management of spinal osteoblastoma: a retrospective study in 18 cases. PLOS One. 2013;8:e74635.

[2]. Kirwan EO, Hutton PA, Pozo JL, Ransford AO. Osteoid osteoma and benign osteoblastoma of the spine. Clinical presentation and treatment. J Bone Joint Surg Br. 1984;66:21-26.

Figure legends:

Figure 1: CT images sagittal (A,B) and axial (C,D) showing the lytic lesion with surrounding sclerosis at the right L3 pedicle extending into neural foramen.

Figure 2: MR T2-weighted sagittal and axial (A,B) images showing hypo intense lesion in the right L3 pedicle. Contrast sagittal, axial and coronal (C-F) images showing moderate enhancement of the lesion.