

Accepted Manuscript

Title: A multicentre cohort study of spinal osteoid osteomas: results of surgical treatment and analysis of local recurrence

Author: Nasir A. Quraishi, Stefano Boriani, Silviu Sabou, Peter P. Varga, Alessandro Luzzati, Ziya L. Gokaslan, Michael G. Fehlings, Charles G. Fisher, Mark B. Dekutoski, Laurence D. Rhines, Jeremy J. Reynolds, Niccole M. Germscheid, Chetan Bettegowda, Richard P. Williams



PII: S1529-9430(16)31016-6
DOI: <http://dx.doi.org/doi: 10.1016/j.spinee.2016.10.010>
Reference: SPINEE 57181

To appear in: *The Spine Journal*

Received date: 8-11-2015
Revised date: 27-7-2016
Accepted date: 12-10-2016

Please cite this article as: Nasir A. Quraishi, Stefano Boriani, Silviu Sabou, Peter P. Varga, Alessandro Luzzati, Ziya L. Gokaslan, Michael G. Fehlings, Charles G. Fisher, Mark B. Dekutoski, Laurence D. Rhines, Jeremy J. Reynolds, Niccole M. Germscheid, Chetan Bettegowda, Richard P. Williams, A multicentre cohort study of spinal osteoid osteomas: results of surgical treatment and analysis of local recurrence, *The Spine Journal* (2016), <http://dx.doi.org/doi: 10.1016/j.spinee.2016.10.010>.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

A Multicentre Cohort Study of Spinal Osteoid Osteomas: Results of Surgical Treatment and Analysis of Local Recurrence

By, Nasir A. Quraishi LLM, FRCS^a, Stefano Boriani, MD^b, Silviu Sabou^a, Peter P. Varga, MD^c, Alessandro Luzzati, MD^d, Ziya L. Gokaslan, MD, FAANS, FACS^e, Michael G. Fehlings, MD, PhD, FRCSC, FACS^f, Charles G. Fisher, MD, MHSc, FRCSC^g, Mark B. Dekutoski, MD^h, Laurence D. Rhines, MDⁱ, Jeremy J. Reynolds, FRCS^j, Niccole M. Gersmeyer MSc^k, Chetan Bettegowda, MD, PhD^l, Richard P. Williams, Fracs^m

^a Centre for Spinal Studies and Surgery, Queens Medical Centre Campus of Nottingham University Hospitals NHS Trust, Derby Road, Nottingham, NG7 2UH, United Kingdom

^b Department of Oncologic and Degenerative Spine Surgery, Rizzoli Institute, via G.C. Pupilli, 1 - 40136 Bologna, Italy

^c The National Center for Spinal Disorders Budapest, Királyhágó u. 1, 1126 Hungary

^d Centro di Chirurgia Ortopedica Oncologica e Ricostruttiva del Rachide Istituto Orthopedico Galeazzi Via Riccardo Galeazzi, 4, 20161 Milano, Italy

^e Department of Neurosurgery, The Warren Alpert Medical School of Brown University, 222 Richmond St, Providence, RI 02903, United States

^f Division of Neurosurgery, Toronto Western Hospital, University Health Network, University of Toronto 399 Bathurst St, Toronto, ON M5T 2S8, Canada

^g University of British Columbia, Combined Neurosurgery and Orthopaedic Spine Program, Blusson Spinal Cord Centre, 6th Floor, 818 West 10th Avenue, Vancouver, British Columbia V5Z 1M9, Canada

^h Department of Orthopaedic Surgery, The CORE Institute 14520 W Granite Valley Dr #210, Sun City West, AZ 85375, United States

ⁱ Department of Neurosurgery, Division of Surgery, The University of Texas MD
Anderson Cancer Center, 1515 Holcombe Blvd # 442, Houston, TX 77030, United
States

^j Spinal Division, Nuffield Orthopedic Centre, University of Oxford Windmill Road,
Headington, Oxford OX3 7HE, United Kingdom

^k Research Department, AOSpine International, Clavadelerstrasse 87270 Davos
Platz, Switzerland

^l Department of Neurosurgery, Johns Hopkins University School of Medicine,
Baltimore, Maryland, United States, 733 N Broadway, Baltimore, MD 21205, United
States

^m Princess Alexandra Hospital, Brisbane, Australia 237 Ipswich Rd, Woolloongabba
QLD 4102, Australia

Corresponding Author:

Mr Nasir A. Quraishi, LL.M, FRCS

Consultant Spinal Surgeon

Honorary Clinical Associate Professor

Centre for Spinal Studies and Surgery

Queens Medical Centre Campus of Nottingham University Hospitals NHS Trust

West Block, D Floor

Queens Medical Centre

Derby Road

Nottingham NG7 2UH

UK

Tel +44 (0) 115 924 9924, Extn 65160 / 61713

1 Fax +44 (0) 115 970 9991

2 Email:nasquraishi@hotmail.com

3
4 Acknowledgements:

5 We are grateful to the collaborating centres' local clinical research personnel and
6 support staff for their active participation. This study was organized and funded by
7 AOSpine International, through the AOSpine Knowledge Forum Tumor, a pathology-
8 focused working group of up to ten international spine experts acting on behalf of
9 AOSpine in the domain of scientific expertise. AOSpine is a clinical division of the
10 AO Foundation – an independent medically guided not-for-profit organization
11 dedicated to improving patient care worldwide. Study support, including data
12 management, analysis, and manuscript preparation was provided directly through
13 AOSpine's Research department and AO's Clinical Investigation and Documentation
14 unit.

15
16 **Abstract**

17 Background Context

18 Spinal osteoid osteomas are benign primary tumours arising predominantly from the
19 posterior column of the spine. These 'osteoblastic' lesions have traditionally been
20 treated with intralesional excision.

21
22 Purpose

23 The purpose of this study was to review the treatment and local recurrence rates for
24 symptomatic spinal osteoid osteomas.

25
26 Study Design/Setting

27 Multicentre ambispective cross-sectional observational cohort study.

Patient Sample

During the study period, a total of 84 patients (65 male, 19 female) were diagnosed with a spinal osteoid osteoma and received surgical treatment. The mean age at surgery was 21.8 ± 9.0 years (range: 6.7-52.4 years) and the mean follow-up was 2.7 years (13 days-14.5 years).

Outcome Measures

Local recurrence, perioperative morbidity and cross-sectional survival.

Methods

Utilizing the AOSpine Knowledge Forum Tumour (AOSKFT) multicentre ambispective database, surgically treated osteoid osteoma cases were identified. Patient demographic, clinical and diagnostic, treatment, local recurrence, perioperative morbidity, and cross-sectional survival data were collected and retrieved. Descriptive statistics were summarized using mean/standard deviation or frequency/ percentage.

Results

In our study most of the tumours were localized in the mobile spine (81/84 (96%)); all patients reported pain as a symptom. According to the postoperative assessment, 10 (12%) patients received an en bloc resection with marginal or wide margins, while two (2%) patients received en bloc resections with intralesional margins, 69 (82%) patients were treated by piecemeal 'intralesional' resection, and 3 (4%) patients were not assessed. A total of six patients (7%) experienced a local recurrence, all of which occurred in patients who had received an intralesional resection.

Conclusions

Benign bone forming tumours of the spine are uncommon. Most patients in our series underwent a piecemeal resection with intralesional margins. This remains safe with a low local recurrence rate. En bloc excision may provide more chance of complete excision of the nidus but is not mandatory. The importance of complete excision of the nidus cannot be over-emphasized.

Keywords: Spinal Osteoid Osteoma; Surgical Management; Recurrence; Primary Spinal Tumor; Cohort Study

Introduction

Osteoid osteomas are benign bone-forming tumours [1, 2]. They are relatively rare as they represent only 3% of all primary bone tumours [3]. In 10% of the cases they occur in the spine with 70-100% located in the posterior elements [2, 4-6]. The majority of osteoid osteomas occur in the 2nd and 3rd decades of life, with a male predilection of 2–3:1 [7]. Osteoid osteomas are characterized by a nidus of osteoid tissue or even mineralized immature bone, with a highly vascularised stroma and are often surrounded by sclerotic reactive bone [7]. The nidus is less than 1.5 cm in diameter by definition, with larger lesions being called osteoblastomas [7]. This is an important distinction as osteoblastomas can be aggressive and thus, careful radiological as well histological analysis is essential to distinguish these tumour types.

Bone scintigraphy is almost invariably positive and has been advocated for localizing the vertebral level in patients with clinically suspected osteoid osteoma [4]. Subsequent targeted CT is generally regarded as the preferred cross-sectional technique for the demonstration and precise localization of the nidus. Osteoid osteoma characteristically manifests as a low-attenuation nidus with central mineralization and varying degrees of perinidal sclerosis [4].

Patients typically present with pain, usually at night-time are relieved by salicylates [8]. There may be stiffness in 89% of cases [9] and scoliosis in 60-70% of adolescents [10, 11] which may become structural [12]

Neurological impairment in osteoid osteomas is rare. Often these tumours go into remission over 2 to 8 years. Surgical intervention is usually indicated in adults with pain not controlled with medication and in adolescents with a risk of developing a structural scoliotic curve [8] The surgical options include intralesional curettage, marginal or en-bloc excision [2, 13-15]. Minimally invasive options such as radio-frequency thermo-ablation [16-18] or CT guided core –excision [19] have also been reported. The purpose of this study was to review the treatment and local recurrence rates for primary symptomatic osteoid osteomas of the spine using the AOSpine Knowledge Forum Tumour (AOSKFT) multicentre database.

Material and methods:

This study was developed and funded under the umbrella of the AOSpine International Knowledge Forum Tumour – AOSpine is a non-profit, international organization of spine care professionals dedicated to delivering knowledge, experience, and evidence to improve patient care and outcomes.

Database Development:

AOSKFT investigators identified centres with experience in the treatment of primary spinal tumours (PSTs) and an established research track record. A feasibility questionnaire was sent to potential sites to gather data on infrastructure, epidemiology, tumour pathology, treatment modalities and outcomes. Centres treating ≥ 10 patients per year with multidisciplinary oncology care and prospectively collected data were identified. All centres obtained institutional review board or ethics committee approval prior to study commencement.

A PST database consisting of a secure, web-based application to support data capture (REDCap, Vanderbilt University, Nashville, TN, USA) was created. The

AOSKFT Steering Committee composed of Spine Oncology Surgeons, Oncologists, Musculoskeletal Pathologists, and Epidemiologists developed six data modules including: 1) demographic, 2) clinical, 3) diagnostic, 4) therapeutic, 5) cross-sectional survival and local recurrence, and 6) peri-operative morbidity. Data collection was performed using a combined ambispective, cross-sectional design with >75 % of cases being collected prospectively. Prospectively collected databases and clinical charts were used for retrospective data capture. When necessary, mortality information was obtained from government vital statistics databases.

Data Collection:

Between 1981 and 2012, data were collected on 1,495 surgically treated patients with specific primary benign and malignant spine tumours [20] performed at 13 spine oncology referral centres from three geographical regions. Cases were only included if they were admitted to one of the participating spine centres with a diagnosis of a PST and received surgical treatment and standardized follow-up. A study coordinator was employed to assist with data collection, data entry, and capture of cross-sectional survival data. Mandatory and quantifiable data points (i.e., dates, estimated blood loss, etc.) were screened for inconsistencies and incompleteness. Outstanding issues were reported to the centres for clarification.

The treating surgeon used the Weinstein-Boriani-Biagini (WBB) [21] classification system to classify the osteoid osteoma. This is an anatomical classification that incorporates the unique anatomic complexity of the spine with an underlying goal of identifying a safe surgical corridor for tumour excision that will ensure sparing of spinal cord without compromising surgical margins [21] (Figure 1). Postoperatively, the treating surgeon classified the surgical resection according to the following:

En bloc: wide or marginal

En bloc: intralesional (if the surgical intent was en bloc, but resulted in an intralesional resection)

Piecemeal: intra-lesional

or not assessed

En bloc resection includes the removal of the tumour nidus as a single intact whole, fully encased by healthy tissue (circumferential margin of tumour-free normal tissue). A wide margin occurs when the plane of dissection has been peripheral to the reactive zone (sclerotic area) through normal tissue; a marginal margin describes that the plane of dissection has been performed along the reactive zone; and an intralesional margin is when the tumour has been violated and the dissection is through the nidus.

Results

Patient Demographics

A total of 84 patients (65 males and 19 females) with osteoid osteoma were identified as having received surgical treatment between July 1990 and June 2012 (Table 1). The mean age at surgery was 21.8 ± 9.0 years (range: 6.7-52.4 years) where 36 were children (0-18 years) and 48 were adults (≥ 19 years). The mean follow-up was 2.7 years (range: 13 days-14.5 years). Most cases appeared in the mobile spine (N = 81, 96%) where C2, T9, T11, L2, and L4 were the most affected vertebrae. All patients reported pain as a symptom.

Figure 2 illustrates a typical example of the presentation of an osteoid osteoma. In this case, a 21-year old male presented with a new onset of painful scoliosis. Radiographs and CT confirmed the diagnosis of an osteoid osteoma at T12 in the concavity. He underwent an uncomplicated intralesional resection.

Tumour Characteristics

The mean size of the osteoid osteoma in the anterior-posterior dimension was 1.40 ± 0.86 cm (range: 0.30-7.00 cm), 1.31 ± 0.60 cm (range: 0.20-4.00 cm) in the left-right dimension, and 1.27 ± 0.53 cm (range: 0.30-3.00 cm) in the cephalad-caudal dimension (Table 2). By using the WBB classification system and grouping the sectors by pedicle (sectors 2-4 and 9-11), spinous process (sectors 12-1), and

vertebral body involvement (sectors 5-8), we found that of the WBB classified tumours (N = 79), 92% involved the pedicles, 23% spinous process, and 25% vertebral body.

Treatment

Preoperative embolization was performed in 15 (19%) patients and the treatment strategies are shown in Table 3. Most patients underwent a posterior surgical approach (N = 69, 83%) and spinal fixation was not utilised in the majority of patients (N = 64, 76%).

Based on the postoperative assessment, 10 (12%) patients received an en bloc resection with marginal or wide margins, whilst two (2%) patients received en bloc resections with intralesional margins, and 69 (82%) patients were treated by piecemeal resection with intralesional margins (Table 4). The type of surgery was not assessed in three patients (4%). Only one patient treated by intralesional resection received adjuvant radiation therapy.

Complications

Complications were observed in five different patients. There were three intra/postoperative complications observed in three different patients: one incidental durotomy, a minor problem with postoperative ventilation and one superficial wound infection. In addition, three delayed complications were observed: postoperative neuropathic pain, pneumonia, and wound dehiscence. One patient suffered both intra/postoperative and delayed complications. These complications were observed in intralesional resected patients except for postoperative neuropathic pain (en bloc: wide resection) and wound dehiscence (unknown resection).

Local Recurrence & Mortality

Six (7%) patients experienced a local recurrence (Table 4) all of which occurred in patients with tumours located in the mobile spine who received a piecemeal 'intralesional' resection. All these recurrences were subsequently treated by further

intralesional resection. Four out of the six local recurrences happened within two years of the index surgery. Information regarding the date of local recurrence was not available for one of the patients. During follow-up, one patient death was observed due to an unknown cause (and not tumour-related), which occurred 2.8 years postoperatively.

Case Presentation

As an example of a patient with a recurrence, we present this case illustration of a 14 year old girl who presented with several months history of cervical night pain. Computed tomography (CT) showed an osteoid osteoma in the arch of C1 (Figure 3). She underwent an intralesional excision performed by a transoral approach with apparent complete removal (Figure 4). A post-operative CT revealed a small degree of residual tumour (Figure 5) but as she was asymptomatic, the surgeon decided to manage her non-operatively.

She did however, represent 5 months later with increasing pain when another CT scan showed clear evidence of the recurrence on the right border of the excision corresponding to the previous surgical site (Figure 6). As a result, it was decided to offer her a further re-excision (Figure 7) following which she had complete resolution of her pain (Figure 8). She had made excellent recovery at the 2 years follow-up without any further recurrence and full cervical motion.

Discussion

This study reports on, to our knowledge, the largest multicentre series of primary symptomatic spinal osteoid osteoma undergoing surgical intervention. It represents data collected from 13 international centres under the guidance of the AOSKFT. During the study period, a total of 84 patients were surgically treated with the majority of the tumours being located in the mobile spine (81/84 (96%)). Sixty-nine patients (82%) underwent a piecemeal 'intralesional' resection, 12 (14%) had en bloc (wide/marginal/intralesional) surgery with complete excision of the nidus. A total of six patients (7%) experienced a local recurrence all of which occurred in patients who received the piecemeal intralesional resection. Four of the 6 patients experienced a local recurrence within two years postoperatively and were treated by

1 further intralesional resection. Although limited to 12 cases, en bloc excision
2 provides more chance of complete removal of the nidus than piecemeal intralesional
3 excision. Whilst conventional intralesional excision surgery remains safe with a low
4 local recurrence and complication rate, importance of excision of the nidus should be
5 the goal.

6
7 Osteoid osteomas may be difficult to detect radiologically. Whilst osteblastomas
8 cause expansile lesions with a well-defined sclerotic margin, osteoid osteomas show
9 reactive bone sclerosis around the nidus. The use of bone scans, CT and MRI has
10 been emphasised with the latter showing high signal intensity in the tumour. Despite
11 the responsiveness of osteoid osteoma to non-steroidal anti-inflammatory
12 medication, surgical treatment is preferred when these agents fail, or to avoid the
13 long term use of anti-inflammatory medication [2]. The need for stabilisation is
14 perhaps only recommended in cases where there has been extensive resection of
15 bone and there is pre-existing deformity.

16
17 In a large study of symptomatic osteoid osteomas, Boriani treated 80 cases with
18 intralesional excision [2]. The recurrence rate was 6% and these were treated
19 successfully with further surgery. In this study, there was only one additional patient
20 who underwent percutaneous radiofrequency coagulation (PRC) for a lesion in the
21 vertebral body [2]. These authors stated that there was limited use for this technique
22 in the spine due to the potential damage to neural structures. Rosenthal et al., who
23 were the early pioneers of this method, suggested that the thermal necrosis was less
24 safe if neural tissues were within 13 mm of the probe [22, 23]. Despite the promising
25 outlook for minimally invasive treatments, more data regarding outcomes and risks of
26 minimally invasive treatments, and especially the risks in PRC are necessary [2].
27 Further, several studies show differing results in small numbers, and there has not
28 been a large study on these minimally invasive treatments for spinal osteoid
29 osteoma [16, 24-26]. There is potential advantage with less tissue trauma reducing
30 post-operative pain and quicker recovery. The main disadvantages include a long
31 learning curve associated with endoscopic surgeries (if used), risks to neurological
32 structures and uncertainty around the histological verification of the specimens using

PRC [2]. A further study is underway by our group comparing treatments between symptomatic non-spinal osteoid osteoma (mostly managed with radiofrequency coagulation in the literature) versus spinal osteoid osteomas (largely treated by intralesional excision as per present study) [27].

Our results support the notion that intralesional resection can be performed safely and with a low recurrence for these benign tumours (which may be benign active) ; this would be inconsistent with Enneking's principles [28]. Goal of surgical treatment of osteoid osteoma remains complete removal of the nidus either by piecemeal or en bloc excision. Wide margin is not required for this benign condition but en bloc excision may provide more chance of complete removal of the nidus than piecemeal excision. However, it is not mandatory and is conceptually distinct from en bloc excision in the treatment of malignant tumours. Thus, complete excision of the nidus is more important than surgical margins in osteoid osteoma and 6 of the 69 patients who had a recurrence likely had this due to incomplete excision of the nidus.

Another interesting presentation of these tumors is scoliosis. Unfortunately, this was not studied in this study but would certainly be an area of future focus. Finally, embolization of the feeding arteries is not really necessary in osteoid osteomas; it was however used in cases where the lesions looked particularly aggressive and where the diagnosis was initially uncertain [29]. The main limitations of our study are the small number of patients undergoing en bloc resection with wide or marginal margins and lack of patient reported outcome measures. However, it remains an ambispective cohort study from 13 international centres in the largest series of primary symptomatic spinal osteoid osteoma treated by surgical intervention.

Conclusion

Most patients in this series underwent a piecemeal intralesional resection (69/84 (82%)) for symptomatic osteoid osteomas with low complication and recurrence rates. Whilst en bloc excision can be performed sometimes quite simply and safely (due to their location within the posterior elements), we do not recommend this form

of excision for all these benign tumours. It is important however, to excise the nidus completely which can be via intralesional resection.

References:

1. Jaffe. Osteoid-osteoma: a benign osteoblastic tumor composed of osteoid and atypical bone. Arch Surg. 1935;31:709–28.
2. Gasbarrini A, Cappuccio M, Bandiera S, Amendola L, van Urk P, Boriani S. Osteoid osteoma of the mobile spine: surgical outcomes in 81 patients. Spine (Phila Pa 1976). 2011;36(24):2089-93.
3. Huvos. Bone Tumors: Diagnosis, Treatment and Prognosis . 2 ed. . Philadelphia: W.B. Saunders; 1991.
4. Greenspan A. Radiologic evaluation of tumors and tumor-like lesions in Orthopaedic imaging: a practical approach, 4th ed, Lippincott Williams and Wilkins, 2004.
5. Jackson RP, Reckling FW, Mants FA. Osteoid osteoma and osteoblastoma. Similar histologic lesions with different natural histories. Clin Orthop Relat Res. 1977;(128):303-13.
6. Greenspan. Radiologic evaluation of tumors and tumor-like lesions. Orthopedic imaging: a practical approach 4th ed: Lippincott Williams & Wilkins; 2004.
7. Rodallec MH, Feydy A, Larousserie F, et al. Diagnostic imaging of solitary tumors of the spine: what to do and say. Radiographics. 2008;28(4):1019-41.
8. Kneisl JS, Simon MA. Medical management compared with operative treatment for osteoid-osteoma. J Bone Joint Surg Am. 1992;74(2):179-85.
9. Kirwan EO, Hutton PA, Pozo JL, Ransford AO. Osteoid osteoma and benign osteoblastoma of the spine. Clinical presentation and treatment. J Bone Joint Surg Br. 1984;66(1):21-6.
10. Keim HA, Reina EG. Osteoid-osteoma as a cause of scoliosis. J Bone Joint Surg Am. 1975;57(2):159-63.
11. Saifuddin A, White J, Sherazi Z, Shaikh MI, Natali C, Ransford AO. Osteoid osteoma and osteoblastoma of the spine. Factors associated with the presence of scoliosis. Spine (Phila Pa 1976). 1998;23(1):47-53.
12. Ransford AO, Pozo JL, Hutton PA, Kirwan EO. The behaviour pattern of the scoliosis associated with osteoid osteoma or osteoblastoma of the spine. J Bone Joint Surg Br. 1984;66(1):16-
13. Marcove RC, Heelan RT, Huvos AG, Healey J, Lindeque BG. Osteoid osteoma. Diagnosis, localization, and treatment. Clin Orthop Relat Res. 1991;(267):197-201.

14. Maiuri F, Signorelli C, Lavano A, Gambardella A, Simari R, D'Andrea F. Osteoid osteomas of the spine. *Surg Neurol.* 1986;25(4):375-80.
15. Lee DH, Malawer MM. Staging and treatment of primary and persistent (recurrent) osteoid osteoma. Evaluation of intraoperative nuclear scanning, tetracycline fluorescence, and tomography. *Clin Orthop Relat Res.* 1992;(281):229-38.
16. Cove JA, Taminiau AH, Obermann WR, Vanderschueren GM. Osteoid osteoma of the spine treated with percutaneous computed tomography-guided thermocoagulation. *Spine (Phila Pa 1976).* 2000;25(10):1283-6.
17. Lindner NJ, Ozaki T, Roedel R, Gosheger G, Winkelmann W, Wortler K. Percutaneous radiofrequency ablation in osteoid osteoma. *J Bone Joint Surg Br.* 2001;83(3):391-6.
18. Osti OL, Sebben R. High-frequency radio-wave ablation of osteoid osteoma in the lumbar spine. *Eur Spine J.* 1998;7(5):422-5.
19. Labbe JL, Clement JL, Duparc B, Poey C, Railhac JJ. Percutaneous extraction of vertebral osteoid osteoma under computed tomography guidance. *Eur Spine J.* 1995;4(6):368-71.
20. Fisher CG, Goldschlager T, Boriani S, et al. An evidence-based medicine model for rare and often neglected neoplastic conditions. *J Neurosurg Spine.* 2014;21(5):704-10.
21. Boriani S, Weinstein JN, Biagini R. Primary bone tumors of the spine. Terminology and surgical staging. *Spine (Phila Pa 1976).* 1997;22(9):1036-44.
22. Rosenthal DI, Hornicke FJ, Torriani M, Gebhardt MC, Mankin HJ. Osteoid osteoma: percutaneous treatment with radiofrequency energy. *Radiology.* 2003;229(1):171-5.
23. Rosenthal DI, Springfield DS, Gebhardt MC, Rosenberg AE, Mankin HJ. Osteoid osteoma: percutaneous radio-frequency ablation. *Radiology.* 1995;197(2):451-4.
24. Hadjipavlou AG, Lander PH, Marchesi D, Katonis PG, Gaitanis IN. Minimally invasive surgery for ablation of osteoid osteoma of the spine. *Spine (Phila Pa 1976).* 2003;28(22):E472-7.
25. Gangi A, Dietemann JL, Guth S, et al. Percutaneous laser photocoagulation of spinal osteoid osteomas under CT guidance. *AJNR Am J Neuroradiol.* 1998;19(10):1955-8.
26. Vanderschueren GM, Taminiau AH, Obermann WR, Bloem JL. Osteoid osteoma: clinical results with thermocoagulation. *Radiology.* 2002;224(1):82-6.
27. Williams BS, Boriani S, Varga A, Luzzati Z, Gokaslan M, Fehlings C, Fisher M, Dekutoski L, Rhines J, Reynolds C, Bettgowda N, Kawahara N, A Quraishi. Spinal Osteoid Osteoma: A Multicenter Retrospective Cohort Study Comparison of Results With Appendicular Osteoid Osteoma. *Spine Society of Europe* 2014.
28. Enneking WF, Spanier SS, Goodman MA. A system for the surgical staging of musculoskeletal sarcoma. 1980. *Clin Orthop Relat Res.* 2003;(415):4-18.
29. Campanacci. Osteoid osteoma. In *Bone and soft tissue tumors.* New York: Springer-Verlag; 1999.

Figure Legends

Figure 1 – The Weinstein-Boriani-Biagini classifications of spinal tumours. The location of the lesion is described using the 12 radiating zones (1 to 12) and 5 concentric layers (A to E) with layers D and E representing epidural and intra-dural involvement, respectively.

Figure 2 – Antero-posterior radiographs (a) and axial CT (b) showing osteoid osteoma in the posterior elements of a T12 vertebra

Figure 3 – Computed tomography (CT) showed an osteoid osteoma in the arch of C1 (arrow).

Figure 4 – An immediate postoperative axial CT showing apparent complete removal of the osteoid osteoma (arrow).

Figure 5 – A follow-up axial CT revealed a small degree of residual tumour (arrow).

Figure 6 – An axial CT scan at the 5 month stage, showing clear evidence of the recurrence on the right border of the excision corresponding to the previous surgical site (circled by radiologist).

Figure 7 – Intraoperative picture of the surgical exposure through the transoral route.

Figure 8 – A postoperative axial CT scan after the re-excision, showing complete excision of the tumor.

1 Table 1: Summary of Patient Demographics

Variables	
Gender	
Female	19 (23)
Male	65 (77)
Age (years)	
At diagnosis	21.3 \pm 9.0
Child (0-18 years)	14.5 \pm 3.0
Adult (\geq 19 years)	27.2 \pm 8.2
At surgery	21.8 \pm 9.0
Child (0-18 years)	14.6 \pm 3.0
Adult (\geq 19 years)	27.1 \pm 8.3
Spinal Location	
Mobile spine	
Cervical	25 (30)
Thoracic	31 (37)
Lumbar	25 (30)
Fixed spine	
Sacral	3 (3)
Clinical Presentation Pain	
Yes	84 (100)
No	0 (0)

Data are presented as N (%) or Mean \pm Standard Deviation

2

3

4

5

6 Table 2: Summary of Spinal Osteoid Osteoma Characteristics

Variables	
Tumor Size (cm)	
Anterior-Posterior	1.40 \pm 0.86
Left - Right	1.31 \pm 0.60
Cephalad-Caudad	1.27 \pm 0.53
WBB Sectors [†]	
2 - 4	42 (53)
9 -11	31 (39)
12 -1	18 (23)
5 - 8	20 (25)

Data are presented as N (%) or Mean \pm Standard Deviation

[†] WBB = Weinstein – Boriani – Biagini, Please note that numbers are higher as the tumors involve multiple segments

Table 3: Summary of Spinal Osteoid Osteoma Treatment

Variables	N (%)
<i>Preoperative Embolization</i>	
Yes	15 (19)
No	62 (81)
<i>Surgical approach</i>	
Anterior	10 (12)
Posterior	69 (83)
Anterior-Posterior	0 (0)
Posterior-Anterior	2 (2)
Other	2 (2)
<i>Method of Spinal Fixation</i>	
Anterior	2 (2)
Posterior	18 (21)
Both	0 (0)
None	64 (76)

Table 4: The Effect of Surgical Treatment on Local Recurrence

Variables	Local Recurrence		
	No	Yes	Total
	N (%)	N (%)	N (%)
Surgical Procedure (and margin)			
En bloc: wide or marginal	10 (12)	0 (0)	10 (12)
En bloc: intra-lesional	2 (2)	0 (0)	2 (2)
Piecemeal: intra-lesional	63 (75)	6 (7)	69 (82)
Piecemeal: not assessed	3 (4)	0 (0)	3 (4)