



Artificial Intelligence

Sem. Ganjil 2023/2024

10. Reasoning with Uncertainty

Lionov



**INFORMATIKA
UNPAR**

- Conditional Probability
- Bayes' Rule
- Bayesian Network

Uncertainty

Uncertainty: “a typical feature” of real world problems for the agent
cause: partial observability, non-determinism, and adversaries

Uncertainty

Uncertainty: “a typical feature” of real world problems for the agent
cause: partial observability, non-determinism, and adversaries

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2 B OK	2,2	3,2	4,2
1,1 OK	2,1 B OK	3,1	4,1

Uncertainty

Uncertainty: “a typical feature” of real world problems for the agent
cause: partial observability, non-determinism, and adversaries

Belief state is a set of all possible world states. Only suitable for small problem, because:

Uncertainty

Uncertainty: “a typical feature” of real world problems for the agent
cause: partial observability, non-determinism, and adversaries

Belief state is a set of all possible world states. Only suitable for small problem, because:

- consider every possible world states
- consider arbitrarily unlikely contingencies
- compare plans that are not guaranteed to achieve the goal

Uncertainty

Uncertainty: “a typical feature” of real world problems for the agent
cause: partial observability, non-determinism, and adversaries

Belief state is a set of all possible world states. Only suitable for small problem, because:

- consider every possible world states
- consider arbitrarily unlikely contingencies
- compare plans that are not guaranteed to achieve the goal

Rational decision (or “the right thing to do”) depends on :

Uncertainty

Uncertainty: “a typical feature” of real world problems for the agent
cause: partial observability, non-determinism, and adversaries

Belief state is a set of all possible world states. Only suitable for small problem, because:

- consider every possible world states
- consider arbitrarily unlikely contingencies
- compare plans that are not guaranteed to achieve the goal

Rational decision (or “the right thing to do”) depends on :

- the relative importance of various goals

Uncertainty

Uncertainty: “a typical feature” of real world problems for the agent
cause: partial observability, non-determinism, and adversaries

Belief state is a set of all possible world states. Only suitable for small problem, because:

- consider every possible world states
- consider arbitrarily unlikely contingencies
- compare plans that are not guaranteed to achieve the goal

Rational decision (or “the right thing to do”) depends on :

- the relative importance of various goals
- the likelihood that, and degree to which, they will be achieved

Uncertainty

Uncertainty: “a typical feature” of real world problems for the agent
cause: partial observability, non-determinism, and adversaries

Belief state is a set of all possible world states. Only suitable for small problem, because:

- consider every possible world states
- consider arbitrarily unlikely contingencies
- compare plans that are not guaranteed to achieve the goal

Rational decision (or “the right thing to do”) depends on :

- the relative importance of various goals
- the likelihood that, and degree to which, they will be achieved

Probability Theory: provides a way of summarizing the uncertainty

Uncertainty

Uncertainty: “a typical feature” of real world problems for the agent
cause: partial observability, non-determinism, and adversaries

Belief state is a set of all possible world states. Only suitable for small problem, because:

- consider every possible world states
- consider arbitrarily unlikely contingencies
- compare plans that are not guaranteed to achieve the goal

Rational decision (or “the right thing to do”) depends on :

- the relative importance of various goals
- the likelihood that, and degree to which, they will be achieved

Probability Theory: provides a way of summarizing the uncertainty

Utility Theory: every state (or state sequence) has a degree of usefulness or utility

Uncertainty

Uncertainty: “a typical feature” of real world problems for the agent
cause: partial observability, non-determinism, and adversaries

Belief state is a set of all possible world states. Only suitable for small problem, because:

- consider every possible world states
- consider arbitrarily unlikely contingencies
- compare plans that are not guaranteed to achieve the goal

Rational decision (or “the right thing to do”) depends on :

- the relative importance of various goals
- the likelihood that, and degree to which, they will be achieved

Probability Theory: provides a way of summarizing the uncertainty

Utility Theory: every state (or state sequence) has a degree of usefulness or utility

Decision Theory: an agent is rational if and only if it chooses the action that yields the highest expected utility, averaged over all the possible outcomes of the action.

Uncertainty

Uncertainty: “a typical feature” of real world problems for the agent
cause: partial observability, non-determinism, and adversaries

Belief state is a set of all possible world states. Only suitable for small problem, because:

- consider every possible world states
- consider arbitrarily unlikely contingencies
- compare plans that are not guaranteed to achieve the goal

Rational decision (or “the right thing to do”) depends on :

- the relative importance of various goals
- the likelihood that, and degree to which, they will be achieved

Probability Theory: provides a way of summarizing the uncertainty

Utility Theory: every state (or state sequence) has a degree of usefulness or utility

Decision Theory: an agent is rational if and only if it chooses the action that yields the highest expected utility, averaged over all the possible outcomes of the action.

Decision Theory = Probability Theory + Utility Theory

Decision-Theory Agent

function DT-AGENT(*percept*) **returns** an *action*

persistent: *belief_state*, probabilistic beliefs about the current state of the world
action, the agent's action

update *belief_state* based on *action* and *percept*

calculate outcome probabilities for actions,

given action descriptions and current *belief_state*

select *action* with highest expected utility

given probabilities of outcomes and utility information

return *action*

Probability Theory

Probability Theory

Ω : sample space, all possible worlds (e.g.: 36 of two dice)

ω : a particular possible world (e.g. one of 36 of two dice)

Probability Theory

Ω : sample space, all possible worlds (e.g.: 36 of two dice)

ω : a particular possible world (e.g. one of 36 of two dice)

$P(\omega)$: a probability model associates with each possible world

$0 \leq P(\omega) \leq 1$ for every ω and $\sum_{\omega \in \Omega} P(\omega) = 1$

Probability Theory

Ω : sample space, all possible worlds (e.g.: 36 of two dice)

ω : a particular possible world (e.g. one of 36 of two dice)

$P(\omega)$: a probability model associates with each possible world

$0 \leq P(\omega) \leq 1$ for every ω and $\sum_{\omega \in \Omega} P(\omega) = 1$

Events: set of possible worlds (e.g. $P(\text{Total} = 11) = P((5,6) + P(6,5)) = \frac{1}{18}$)

Probability Theory

Ω : sample space, all possible worlds (e.g.: 36 of two dice)

ω : a particular possible world (e.g. one of 36 of two dice)

$P(\omega)$: a probability model associates with each possible world

$0 \leq P(\omega) \leq 1$ for every ω and $\sum_{\omega \in \Omega} P(\omega) = 1$

Events: set of possible worlds (e.g. $P(\text{Total} = 11) = P((5,6) + P(6,5)) = \frac{1}{18}$)

Unconditional/prior probabilities: degrees of belief in the absence of other information

Probability Theory

Ω : sample space, all possible worlds (e.g.: 36 of two dice)

ω : a particular possible world (e.g. one of 36 of two dice)

$P(\omega)$: a probability model associates with each possible world

$$0 \leq P(\omega) \leq 1 \text{ for every } \omega \text{ and } \sum_{\omega \in \Omega} P(\omega) = 1$$

Events: set of possible worlds (e.g. $P(\text{Total} = 11) = P((5,6) + P(6,5)) = \frac{1}{18}$)

Unconditional/prior probabilities: degrees of belief in the absence of other information

Conditional/posterior probabilities: degrees of belief we have some **evidence**

$P(\text{doubles} | D_1 = 5)$: probability of rolling doubles **given** the first dice is a 5

Probability Theory

Ω : sample space, all possible worlds (e.g.: 36 of two dice)

ω : a particular possible world (e.g. one of 36 of two dice)

$P(\omega)$: a probability model associates with each possible world

$$0 \leq P(\omega) \leq 1 \text{ for every } \omega \text{ and } \sum_{\omega \in \Omega} P(\omega) = 1$$

Events: set of possible worlds (e.g. $P(\text{Total} = 11) = P((5,6) + P(6,5)) = \frac{1}{18}$)

Unconditional/prior probabilities: degrees of belief in the absence of other information

Conditional/posterior probabilities: degrees of belief we have some **evidence**

$P(\text{doubles} | D_1 = 5)$: probability of rolling doubles **given** the first dice is a 5

Product Rule: $P(a \wedge b) = P(a | b)P(b)$

$$P(a | b) = \frac{P(a \wedge b)}{P(b)}, \text{ holds whenever } P(b) > 0$$

Probability Notation & Assertion

Probability Notation & Assertion

Random variable: aspect of the world which we have uncertainty
e.g. *Cavity*: is there a cavity?

Probability Notation & Assertion

Random variable: aspect of the world which we have uncertainty
e.g. *Cavity*: is there a cavity?

Domain of random variable: set of possible values
e.g. *Cavity* in {true, false} (or {+cavity, \neg cavity})

Probability Notation & Assertion

Random variable: aspect of the world which we have uncertainty
e.g. *Cavity*: is there a cavity?

Domain of random variable: set of possible values
e.g. *Cavity* in {true, false} (or {+cavity, ¬cavity})

Notation: random variables using uppercase, values using lowercase (e.g. $\sum_x P(X = x)$)
Shorthand: $P(x)$, iff all domain entries are unique

Probability Notation & Assertion

Random variable: aspect of the world which we have uncertainty
e.g. *Cavity*: is there a cavity?

Domain of random variable: set of possible values
e.g. *Cavity* in {true, false} (or {+cavity, ¬cavity})

Notation: random variables using uppercase, values using lowercase (e.g. $\sum_x P(X = x)$)

Shorthand: $P(x)$, iff all domain entries are unique

Associate a **probability** with each value

Probability Notation & Assertion

Random variable: aspect of the world which we have uncertainty
e.g. *Cavity*: is there a cavity?

Domain of random variable: set of possible values
e.g. *Cavity* in {true, false} (or {+cavity, ¬cavity})

Notation: random variables using uppercase, values using lowercase (e.g. $\sum_x P(X = x)$)
Shorthand: $P(x)$, iff all domain entries are unique

Associate a **probability** with each value

$P(\text{Cavity} = \text{true}) = P(\text{cavity}) = 0.4$, $P(\text{Cavity} = \text{false}) = P(\neg \text{cavity}) = 0.6$
where $\forall x P(X = x) > 0$ and $\sum_x P(X = x) = 1$

Probability Notation & Assertion

Random variable: aspect of the world which we have uncertainty
e.g. *Cavity*: is there a cavity?

Domain of random variable: set of possible values
e.g. *Cavity* in {true, false} (or {+cavity, ¬cavity})

Notation: random variables using uppercase, values using lowercase (e.g. $\sum_x P(X = x)$)
Shorthand: $P(x)$, iff all domain entries are unique

Associate a **probability** with each value

$P(\text{Cavity} = \text{true}) = P(\text{cavity}) = 0.4$, $P(\text{Cavity} = \text{false}) = P(\neg \text{cavity}) = 0.6$
where $\forall x P(X = x) > 0$ and $\sum_x P(X = x) = 1$

Use the connectives of propositional logic:

e.g. $P(\text{cavity} | \neg \text{toothache} \wedge \text{teen}) = 0.1$ or $P(\text{cavity} | \neg \text{toothache}, \text{teen}) = 0.1$

Probability Notation & Assertion

Random variable: aspect of the world which we have uncertainty
e.g. *Cavity*: is there a cavity?

Domain of random variable: set of possible values
e.g. *Cavity* in {true, false} (or {+cavity, ¬cavity})

Notation: random variables using uppercase, values using lowercase (e.g. $\sum_x P(X = x)$)
Shorthand: $P(x)$, iff all domain entries are unique

Associate a **probability** with each value

$P(\text{Cavity} = \text{true}) = P(\text{cavity}) = 0.4$, $P(\text{Cavity} = \text{false}) = P(\neg \text{cavity}) = 0.6$
where $\forall x P(X = x) > 0$ and $\sum_x P(X = x) = 1$

Use the connectives of propositional logic:

e.g. $P(\text{cavity} | \neg \text{toothache} \wedge \text{teen}) = 0.1$ or $P(\text{cavity} | \neg \text{toothache}, \text{teen}) = 0.1$

Distribution: table of probabilities of values

Probability Notation & Assertion

Random variable: aspect of the world which we have uncertainty
e.g. *Cavity*: is there a cavity?

Domain of random variable: set of possible values
e.g. *Cavity* in {true, false} (or {+cavity, ¬cavity})

Notation: random variables using uppercase, values using lowercase (e.g. $\sum_x P(X = x)$)
Shorthand: $P(x)$, iff all domain entries are unique

Associate a **probability** with each value

$P(\text{Cavity} = \text{true}) = P(\text{cavity}) = 0.4$, $P(\text{Cavity} = \text{false}) = P(\neg \text{cavity}) = 0.6$
where $\forall x P(X = x) > 0$ and $\sum_x P(X = x) = 1$

Use the connectives of propositional logic:

e.g. $P(\text{cavity} | \neg \text{toothache} \wedge \text{teen}) = 0.1$ or $P(\text{cavity} | \neg \text{toothache}, \text{teen}) = 0.1$

Distribution: table of probabilities of values

Probability Distribution: for a random variable X , $P(X)$ gives the values of $P(X = x_i)$ for each possible i . E.g.: $\mathcal{P}(\text{Weather}) = \langle 0.6, 0.1, 0.28, 0.01 \rangle$ for sun, rain, cloud, snow.

Probability Notation & Assertion

Random variable: aspect of the world which we have uncertainty
e.g. *Cavity*: is there a cavity?

Domain of random variable: set of possible values
e.g. *Cavity* in {true, false} (or {+cavity, ¬cavity})

Notation: random variables using uppercase, values using lowercase (e.g. $\sum_x P(X = x)$)
Shorthand: $P(x)$, iff all domain entries are unique

Associate a **probability** with each value

$P(\text{Cavity} = \text{true}) = P(\text{cavity}) = 0.4$, $P(\text{Cavity} = \text{false}) = P(\neg \text{cavity}) = 0.6$
where $\forall x P(X = x) > 0$ and $\sum_x P(X = x) = 1$

Use the connectives of propositional logic:

e.g. $P(\text{cavity} | \neg \text{toothache} \wedge \text{teen}) = 0.1$ or $P(\text{cavity} | \neg \text{toothache}, \text{teen}) = 0.1$

Distribution: table of probabilities of values

Probability Distribution: for a random variable X , $P(X)$ gives the values of $P(X = x_i)$ for each possible i . E.g.: $\mathcal{P}(\text{Weather}) = \langle 0.6, 0.1, 0.28, 0.01 \rangle$ for sun, rain, cloud, snow.

Conditional Distribution: $\mathcal{P}(X|Y)$ gives the values of $P(X = x_i | Y = y_j)$ for each possible i, j pair

Full Joint Distribution

Needs notation for distributions on multiple random variables

Full Joint Distribution

Needs notation for distributions on multiple random variables

The **joint probability distribution** $\mathcal{P}(X, Y)$ denotes probabilities of all combinations of the values of X and Y . E.g. $\mathcal{P}(\text{Weather}, \text{Cavity})$ is a 4×2 table

Full Joint Distribution

Needs notation for distributions on multiple random variables

The **joint probability distribution** $\mathcal{P}(X, Y)$ denotes probabilities of all combinations of the values of X and Y . E.g. $\mathcal{P}(\text{Weather}, \text{Cavity})$ is a 4×2 table

Concise: the product rule for all possible values (of Weather and Cavity) can be written as a single equation $\mathcal{P}(\text{Weather}, \text{Cavity}) = \mathcal{P}(\text{Weather}|\text{Cavity})\mathcal{P}(\text{Cavity})$

Full Joint Distribution

Needs notation for distributions on multiple random variables

The **joint probability distribution** $\mathcal{P}(X, Y)$ denotes probabilities of all combinations of the values of X and Y . E.g. $\mathcal{P}(\text{Weather}, \text{Cavity})$ is a 4×2 table

Concise: the product rule for all possible values (of Weather and Cavity) can be written as a single equation $\mathcal{P}(\text{Weather}, \text{Cavity}) = \mathcal{P}(\text{Weather}|\text{Cavity})\mathcal{P}(\text{Cavity})$

Full Joint (probability) Distribution is the joint distribution for *all* of the random variables

Full Joint Distribution

Needs notation for distributions on multiple random variables

The **joint probability distribution** $\mathcal{P}(X, Y)$ denotes probabilities of all combinations of the values of X and Y . E.g. $\mathcal{P}(\text{Weather}, \text{Cavity})$ is a 4×2 table

Concise: the product rule for all possible values (of Weather and Cavity) can be written as a single equation $\mathcal{P}(\text{Weather}, \text{Cavity}) = \mathcal{P}(\text{Weather}|\text{Cavity})\mathcal{P}(\text{Cavity})$

Full Joint (probability) Distribution is the joint distribution for *all* of the random variables
E.g. given Toothache , Cavity , and Catch , the full joint distribution is a $2 \times 2 \times 2$ table

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	0.108	0.012	0.072	0.008
\neg <i>cavity</i>	0.016	0.064	0.144	0.576

Full Joint Distribution

Needs notation for distributions on multiple random variables

The **joint probability distribution** $\mathcal{P}(X, Y)$ denotes probabilities of all combinations of the values of X and Y . E.g. $\mathcal{P}(\text{Weather}, \text{Cavity})$ is a 4×2 table

Concise: the product rule for all possible values (of Weather and Cavity) can be written as a single equation $\mathcal{P}(\text{Weather}, \text{Cavity}) = \mathcal{P}(\text{Weather}|\text{Cavity})\mathcal{P}(\text{Cavity})$

Full Joint (probability) Distribution is the joint distribution for *all* of the random variables
E.g. given Toothache , Cavity , and Catch , the full joint distribution is a $2 \times 2 \times 2$ table

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	0.108	0.012	0.072	0.008
\neg <i>cavity</i>	0.016	0.064	0.144	0.576

For a set of random variables X_1, X_2, \dots, X_n , $\mathcal{P}(X_1, X_2, \dots, X_n)$ must obey:

- $\mathcal{P}(X_1, X_2, \dots, X_n) > 0$
- $\sum_{(X_1, X_2, \dots, X_n)} \mathcal{P}(X_1, X_2, \dots, X_n) = 1$

Probabilistic Inference

Probabilistic Inference

Computation of posterior probabilities given observed evidence
with full joint distribution as \mathcal{KB}

Probabilistic Inference

Computation of posterior probabilities given observed evidence
with full joint distribution as \mathcal{KB}

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	0.108	0.012	0.072	0.008
\neg <i>cavity</i>	0.016	0.064	0.144	0.576

Probabilistic Inference

Computation of posterior probabilities given observed evidence
with full joint distribution as \mathcal{KB}

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	0.108	0.012	0.072	0.008
\neg <i>cavity</i>	0.016	0.064	0.144	0.576

Example: $P(\text{cavity} \vee \text{toothache}) = 0.28$

Probabilistic Inference

Computation of posterior probabilities given observed evidence
with full joint distribution as \mathcal{KB}

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	0.108	0.012	0.072	0.008
\neg <i>cavity</i>	0.016	0.064	0.144	0.576

Example: $P(\text{cavity} \vee \text{toothache}) = 0.28$

Extract the distribution over some subset of variables or a single variable:

Probabilistic Inference

Computation of posterior probabilities given observed evidence
with full joint distribution as \mathcal{KB}

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	0.108	0.012	0.072	0.008
\neg <i>cavity</i>	0.016	0.064	0.144	0.576

Example: $P(\text{cavity} \vee \text{toothache}) = 0.28$

Extract the distribution over some subset of variables or a single variable:

Marginalization: summing out, combined by adding and remove other variables

Marginal distributions are sub-tables which eliminate variables

$$\mathcal{P}(Y) = \sum_z \mathcal{P}(Y, Z = z)$$

Probabilistic Inference

Computation of posterior probabilities given observed evidence
with full joint distribution as \mathcal{KB}

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	0.108	0.012	0.072	0.008
\neg <i>cavity</i>	0.016	0.064	0.144	0.576

Example: $P(\text{cavity} \vee \text{toothache}) = 0.28$

Extract the distribution over some subset of variables or a single variable:

Marginalization: summing out, combined by adding and remove other variables

Marginal distributions are sub-tables which eliminate variables

$$\mathcal{P}(Y) = \sum_z \mathcal{P}(Y, Z = z)$$

E.g. $\mathcal{P}(\text{cavity}) = \langle 0.2, 0.8 \rangle$

Probabilistic Inference

Computation of posterior probabilities given observed evidence
with full joint distribution as \mathcal{KB}

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	0.108	0.012	0.072	0.008
\neg <i>cavity</i>	0.016	0.064	0.144	0.576

Example: $P(\text{cavity} \vee \text{toothache}) = 0.28$

Extract the distribution over some subset of variables or a single variable:

Marginalization: summing out, combined by adding and remove other variables

Marginal distributions are sub-tables which eliminate variables

$$\mathcal{P}(Y) = \sum_z \mathcal{P}(Y, Z = z)$$

E.g. $\mathcal{P}(\text{cavity}) = \langle 0.2, 0.8 \rangle$

Using the product rule, we obtain a rule called **conditioning**:

$$\mathcal{P}(Y) = \sum_z \mathcal{P}(Y|z)P(z)$$

Probabilistic Inference

Computation of posterior probabilities given observed evidence
with full joint distribution as \mathcal{KB}

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	0.108	0.012	0.072	0.008
\neg <i>cavity</i>	0.016	0.064	0.144	0.576

We mostly want conditional probabilities of some variables, given evidence about others

Probabilistic Inference

Computation of posterior probabilities given observed evidence
with full joint distribution as \mathcal{KB}

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	0.108	0.012	0.072	0.008
\neg <i>cavity</i>	0.016	0.064	0.144	0.576

We mostly want conditional probabilities of some variables, given evidence about others

E.g.: compute the probability of a cavity, given evidence of a toothache

$$P(\text{cavity} \mid \text{toothache}) = 0.6 \text{ and } P(\neg \text{cavity} \mid \text{toothache}) = 0.4$$

Probabilistic Inference

Computation of posterior probabilities given observed evidence
with full joint distribution as \mathcal{KB}

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	0.108	0.012	0.072	0.008
\neg <i>cavity</i>	0.016	0.064	0.144	0.576

We mostly want conditional probabilities of some variables, given evidence about others

E.g.: compute the probability of a cavity, given evidence of a toothache

$$P(\text{cavity} \mid \text{toothache}) = 0.6 \text{ and } P(\neg \text{cavity} \mid \text{toothache}) = 0.4$$

Normalization: ensuring the distribution of \mathcal{P} adds up to 1 (using α)

$$\begin{aligned} P(\text{Cavity} \mid \text{toothache}) &= \alpha P(\text{Cavity}, \text{toothache}) \\ &= \alpha \langle 0.12, 0.08 \rangle = \langle 0.6, 0.4 \rangle \end{aligned}$$

Note that $P(\text{toothache})$ is not even needed!

Probabilistic Inference

Computation of posterior probabilities given observed evidence
with full joint distribution as \mathcal{KB}

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	0.108	0.012	0.072	0.008
\neg <i>cavity</i>	0.016	0.064	0.144	0.576

General inference procedure:

Probabilistic Inference

Computation of posterior probabilities given observed evidence
with full joint distribution as \mathcal{KB}

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	0.108	0.012	0.072	0.008
\neg <i>cavity</i>	0.016	0.064	0.144	0.576

General inference procedure:

- query involves a single variable X (e.g. Cavity)
- E is the list of evidence variables (e.g. Toothache)
- e is the list of observed values for E
- Y the remaining unobserved variables (e.g. Catch)

Probabilistic Inference

Computation of posterior probabilities given observed evidence
with full joint distribution as \mathcal{KB}

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	0.108	0.012	0.072	0.008
\neg <i>cavity</i>	0.016	0.064	0.144	0.576

General inference procedure:

- query involves a single variable X (e.g. Cavity)
- E is the list of evidence variables (e.g. Toothache)
- e is the list of observed values for E
- Y the remaining unobserved variables (e.g. Catch)

$$P(X | e) = \alpha P(X, e) = \alpha \sum_y P(X, e, y)$$

Probabilistic Inference

Computation of posterior probabilities given observed evidence
with full joint distribution as \mathcal{KB}

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	0.108	0.012	0.072	0.008
\neg <i>cavity</i>	0.016	0.064	0.144	0.576

General inference procedure:

- query involves a single variable X (e.g. Cavity)
- E is the list of evidence variables (e.g. Toothache)
- e is the list of observed values for E
- Y the remaining unobserved variables (e.g. Catch)

$$P(X | e) = \alpha P(X, e) = \alpha \sum_y P(X, e, y)$$

where summation is over all possible combinations of values of Y

Probabilistic Inference

Computation of posterior probabilities given observed evidence
with full joint distribution as \mathcal{KB}

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	0.108	0.012	0.072	0.008
\neg <i>cavity</i>	0.016	0.064	0.144	0.576

General inference procedure:

- query involves a single variable X (e.g. Cavity)
- E is the list of evidence variables (e.g. Toothache)
- e is the list of observed values for E
- Y the remaining unobserved variables (e.g. Catch)

$$P(X | e) = \alpha P(X, e) = \alpha \sum_y P(X, e, y)$$

where summation is over all possible combinations of values of Y

Problems: needs $O(2^n)$ space and $O(2^n)$ time to process

Independence

Expand the full joint distribution: add **Weather**

Independence

Expand the full joint distribution: add *Weather*

$$P(\text{toothache}, \text{catch}, \text{cavity}, \text{cloud}) = \\ P(\text{cloud} \mid \text{toothache}, \text{catch}, \text{cavity})P(\text{toothache}, \text{catch}, \text{cavity})$$

Independence

Expand the full joint distribution: add *Weather*

$$P(\text{toothache}, \text{catch}, \text{cavity}, \text{cloud}) = \\ P(\text{cloud} \mid \text{toothache}, \text{catch}, \text{cavity})P(\text{toothache}, \text{catch}, \text{cavity})$$

However, the weather does not influence the dental variables:

Independence

Expand the full joint distribution: add **Weather**

$$P(\text{toothache}, \text{catch}, \text{cavity}, \text{cloud}) = \\ P(\text{cloud} \mid \text{toothache}, \text{catch}, \text{cavity})P(\text{toothache}, \text{catch}, \text{cavity})$$

However, the weather does not influence the dental variables:

Independence property: $P(\text{cloud} \mid \text{toothache}, \text{catch}, \text{cavity}) = P(\text{cloud})$

Independence

Expand the full joint distribution: add **Weather**

$$P(\text{toothache, catch, cavity, cloud}) = \\ P(\text{cloud} \mid \text{toothache, catch, cavity})P(\text{toothache, catch, cavity})$$

However, the weather does not influence the dental variables:

Independence property: $P(\text{cloud} \mid \text{toothache, catch, cavity}) = P(\text{cloud})$

$$\begin{aligned} \text{Therefore, } P(\text{Toothache, Catch, Cavity, Weather}) &= \\ &= P(\text{Toothache, Catch, Cavity})P(\text{Weather}) \end{aligned}$$

Independence

Expand the full joint distribution: add **Weather**

$$P(\text{toothache}, \text{catch}, \text{cavity}, \text{cloud}) = \\ P(\text{cloud} \mid \text{toothache}, \text{catch}, \text{cavity})P(\text{toothache}, \text{catch}, \text{cavity})$$

However, the weather does not influence the dental variables:

Independence property: $P(\text{cloud} \mid \text{toothache}, \text{catch}, \text{cavity}) = P(\text{cloud})$

$$\begin{aligned} \text{Therefore, } P(\text{Toothache}, \text{Catch}, \text{Cavity}, \text{Weather}) &= \\ &= P(\text{Toothache}, \text{Catch}, \text{Cavity})P(\text{Weather}) \end{aligned}$$

From 32-entries table to 8-entries table and 4-entries table!

Independence

Expand the full joint distribution: add **Weather**

$$P(\text{toothache}, \text{catch}, \text{cavity}, \text{cloud}) = \\ P(\text{cloud} \mid \text{toothache}, \text{catch}, \text{cavity})P(\text{toothache}, \text{catch}, \text{cavity})$$

However, the weather does not influence the dental variables:

Independence property: $P(\text{cloud} \mid \text{toothache}, \text{catch}, \text{cavity}) = P(\text{cloud})$

$$\begin{aligned} \text{Therefore, } \mathcal{P}(\text{Toothache}, \text{Catch}, \text{Cavity}, \text{Weather}) &= \\ &= \mathcal{P}(\text{Toothache}, \text{Catch}, \text{Cavity})\mathcal{P}(\text{Weather}) \end{aligned}$$

From 32-entries table to 8-entries table and 4-entries table!

Independence between variables **X** and **Y**:

$$\mathcal{P}(X \mid Y) = \mathcal{P}(X) \text{ or } \mathcal{P}(Y \mid X) = \mathcal{P}(Y) \text{ or } \mathcal{P}(X, Y) = \mathcal{P}(X)\mathcal{P}(Y)$$

Independence

Expand the full joint distribution: add *Weather*

$$P(\text{toothache}, \text{catch}, \text{cavity}, \text{cloud}) = \\ P(\text{cloud} \mid \text{toothache}, \text{catch}, \text{cavity})P(\text{toothache}, \text{catch}, \text{cavity})$$

However, the weather does not influence the dental variables:

Independence property: $P(\text{cloud} \mid \text{toothache}, \text{catch}, \text{cavity}) = P(\text{cloud})$

$$\text{Therefore, } \mathcal{P}(\text{Toothache}, \text{Catch}, \text{Cavity}, \text{Weather}) = \\ = \mathcal{P}(\text{Toothache}, \text{Catch}, \text{Cavity})\mathcal{P}(\text{Weather})$$

From 32-entries table to 8-entries table and 4-entries table!

Independence between variables *X* and *Y*:

$$\mathcal{P}(X \mid Y) = \mathcal{P}(X) \text{ or } \mathcal{P}(Y \mid X) = \mathcal{P}(Y) \text{ or } \mathcal{P}(X, Y) = \mathcal{P}(X)\mathcal{P}(Y)$$

Independence assertions can help in reducing the size of the domain representation and the complexity of the inference problem

Independence

Expand the full joint distribution: add **Weather**

$$P(\text{toothache}, \text{catch}, \text{cavity}, \text{cloud}) = \\ P(\text{cloud} \mid \text{toothache}, \text{catch}, \text{cavity})P(\text{toothache}, \text{catch}, \text{cavity})$$

However, the weather does not influence the dental variables:

Independence property: $P(\text{cloud} \mid \text{toothache}, \text{catch}, \text{cavity}) = P(\text{cloud})$

$$\text{Therefore, } \mathcal{P}(\text{Toothache}, \text{Catch}, \text{Cavity}, \text{Weather}) = \\ = \mathcal{P}(\text{Toothache}, \text{Catch}, \text{Cavity})\mathcal{P}(\text{Weather})$$

From 32-entries table to 8-entries table and 4-entries table!

Independence between variables **X** and **Y**:

$$\mathcal{P}(X \mid Y) = \mathcal{P}(X) \text{ or } \mathcal{P}(Y \mid X) = \mathcal{P}(Y) \text{ or } \mathcal{P}(X, Y) = \mathcal{P}(X)\mathcal{P}(Y)$$

Independence assertions can help in reducing the size of the domain representation and the complexity of the inference problem

Unfortunately, clean separation of entire sets of variables is quite rare because it based on knowledge of the domain. We need more subtle methods!!!

Bayes' Rule

Bayes' Rule

Product rule can be written in two forms

$$P(a \wedge b) = P(a | b)P(b) = P(b | a)P(a)$$

Bayes' Rule

Product rule can be written in two forms

$$P(a \wedge b) = P(a | b)P(b) = P(b | a)P(a)$$

We get “the most important AI equation”, **Bayes' Rule**:

$$P(b | a) = \frac{P(a | b)P(b)}{P(a)}$$

Bayes' Rule

Product rule can be written in two forms

$$P(a \wedge b) = P(a | b)P(b) = P(b | a)P(a)$$

We get “the most important AI equation”, **Bayes' Rule**:

$$P(b | a) = \frac{P(a | b)P(b)}{P(a)}$$

- It lets us build one conditional from its reverse!
- Often one conditional is tricky but the other one is simple
- Describes an “update” step from prior $P(b)$ to posterior $P(b | a)$

More general version conditionalized on some background evidence e

$$P(Y | X, e) = \frac{P(X | Y, e)P(Y | e)}{P(X | e)}$$

Bayes' Rule

Product rule can be written in two forms

$$P(a \wedge b) = P(a | b)P(b) = P(b | a)P(a)$$

We get “the most important AI equation”, **Bayes' Rule**:

$$P(b | a) = \frac{P(a | b)P(b)}{P(a)}$$

- It lets us build one conditional from its reverse!
- Often one conditional is tricky but the other one is simple
- Describes an “update” step from prior $P(b)$ to posterior $P(b | a)$

More general version conditionalized on some background evidence e

$$P(Y | X, e) = \frac{P(X | Y, e)P(Y | e)}{P(X | e)}$$

The general form of Bayes' Rule with normalization is

$$P(Y | X) = \alpha P(X | Y)P(Y)$$

Applying Bayes' Rule

Applying Bayes' Rule

We perceive, as evidence, the **effect** of some unknown **cause** and we would like to determine that cause:

Applying Bayes' Rule

We perceive, as evidence, the **effect** of some unknown **cause** and we would like to determine that cause:

$$P(\text{cause} \mid \text{effect}) = \frac{P(\text{effect} \mid \text{cause})P(\text{cause})}{P(\text{effect})}$$

Applying Bayes' Rule

We perceive, as evidence, the **effect** of some unknown **cause** and we would like to determine that cause:

$$P(\text{cause} \mid \text{effect}) = \frac{P(\text{effect} \mid \text{cause})P(\text{cause})}{P(\text{effect})}$$

$P(\text{cause} \mid \text{effect})$: **diagnostic** direction

$P(\text{effect} \mid \text{cause})$: **causal** direction

Applying Bayes' Rule

We perceive, as evidence, the **effect** of some unknown **cause** and we would like to determine that cause:

$$P(\text{cause} \mid \text{effect}) = \frac{P(\text{effect} \mid \text{cause})P(\text{cause})}{P(\text{effect})}$$

$P(\text{cause} \mid \text{effect})$: **diagnostic** direction

$P(\text{effect} \mid \text{cause})$: **causal** direction

Medical diagnosis: conditional probabilities on causal relationship, the doctor knows

$P(\text{symptoms} \mid \text{disease})$ and want to derive $P(\text{disease} \mid \text{symptoms})$

Applying Bayes' Rule

We perceive, as evidence, the **effect** of some unknown **cause** and we would like to determine that cause:

$$P(\text{cause} \mid \text{effect}) = \frac{P(\text{effect} \mid \text{cause})P(\text{cause})}{P(\text{effect})}$$

$P(\text{cause} \mid \text{effect})$: **diagnostic** direction

$P(\text{effect} \mid \text{cause})$: **causal** direction

Medical diagnosis: conditional probabilities on causal relationship, the doctor knows

$P(\text{symptoms} \mid \text{disease})$ and want to derive $P(\text{disease} \mid \text{symptoms})$

The doctor doesn't need Bayes if she has information on diagnostic direction

Applying Bayes' Rule

We perceive, as evidence, the **effect** of some unknown **cause** and we would like to determine that cause:

$$P(\text{cause} \mid \text{effect}) = \frac{P(\text{effect} \mid \text{cause})P(\text{cause})}{P(\text{effect})}$$

$P(\text{cause} \mid \text{effect})$: **diagnostic** direction

$P(\text{effect} \mid \text{cause})$: **causal** direction

Medical diagnosis: conditional probabilities on causal relationship, the doctor knows

$P(\text{symptoms} \mid \text{disease})$ and want to derive $P(\text{disease} \mid \text{symptoms})$

The doctor doesn't need Bayes if she has information on diagnostic direction

- However, if there is sudden epidemic, $P(\text{cause})$ will go up and $P(\text{cause} \mid \text{effect})$ increase proportionately.

Applying Bayes' Rule

We perceive, as evidence, the **effect** of some unknown **cause** and we would like to determine that cause:

$$P(\text{cause} \mid \text{effect}) = \frac{P(\text{effect} \mid \text{cause})P(\text{cause})}{P(\text{effect})}$$

$P(\text{cause} \mid \text{effect})$: **diagnostic** direction

$P(\text{effect} \mid \text{cause})$: **causal** direction

Medical diagnosis: conditional probabilities on causal relationship, the doctor knows

$P(\text{symptoms} \mid \text{disease})$ and want to derive $P(\text{disease} \mid \text{symptoms})$

The doctor doesn't need Bayes if she has information on diagnostic direction

- However, if there is sudden epidemic, $P(\text{cause})$ will go up and $P(\text{cause} \mid \text{effect})$ increase proportionately.
- She doesn't have any idea how to update the information on $P(\text{cause} \mid \text{effect})$

Applying Bayes' Rule

We perceive, as evidence, the **effect** of some unknown **cause** and we would like to determine that cause:

$$P(\text{cause} \mid \text{effect}) = \frac{P(\text{effect} \mid \text{cause})P(\text{cause})}{P(\text{effect})}$$

$P(\text{cause} \mid \text{effect})$: **diagnostic** direction

$P(\text{effect} \mid \text{cause})$: **causal** direction

Medical diagnosis: conditional probabilities on causal relationship, the doctor knows

$P(\text{symptoms} \mid \text{disease})$ and want to derive $P(\text{disease} \mid \text{symptoms})$

The doctor doesn't need Bayes if she has information on diagnostic direction

- However, if there is sudden epidemic, $P(\text{cause})$ will go up and $P(\text{cause} \mid \text{effect})$ increase proportionately.
- She doesn't have any idea how to update the information on $P(\text{cause} \mid \text{effect})$

Diagnostic knowledge is often more fragile than causal knowledge!

Combining Evidence

What happens if there are two or more pieces of evidence?

Combining Evidence

What happens if there are two or more pieces of evidence?

$\mathcal{P}(\text{Cavity} \mid \text{toothache} \wedge \text{catch}) = \alpha\langle 0.108, 0.016 \rangle$ (from full joint distribution)

Combining Evidence

What happens if there are two or more pieces of evidence?

$\mathcal{P}(\text{Cavity} \mid \text{toothache} \wedge \text{catch}) = \alpha \langle 0.108, 0.016 \rangle$ (from full joint distribution)

Try Bayes' rule:

$\mathcal{P}(\text{Cavity} \mid \text{toothache} \wedge \text{catch}) = \alpha \mathcal{P}(\text{toothache} \wedge \text{catch} \mid \text{Cavity}) \mathcal{P}(\text{Cavity})$

Combining Evidence

What happens if there are two or more pieces of evidence?

$\mathcal{P}(\text{Cavity} \mid \text{toothache} \wedge \text{catch}) = \alpha \langle 0.108, 0.016 \rangle$ (from full joint distribution)

Try Bayes' rule:

$\mathcal{P}(\text{Cavity} \mid \text{toothache} \wedge \text{catch}) = \alpha \mathcal{P}(\text{toothache} \wedge \text{catch} \mid \text{Cavity}) \mathcal{P}(\text{Cavity})$

Unfortunately, no better than full joint distribution

Combining Evidence

What happens if there are two or more pieces of evidence?

$\mathcal{P}(\text{Cavity} \mid \text{toothache} \wedge \text{catch}) = \alpha \langle 0.108, 0.016 \rangle$ (from full joint distribution)

Try Bayes' rule:

$\mathcal{P}(\text{Cavity} \mid \text{toothache} \wedge \text{catch}) = \alpha \mathcal{P}(\text{toothache} \wedge \text{catch} \mid \text{Cavity}) \mathcal{P}(\text{Cavity})$

Unfortunately, no better than full joint distribution

Independence: unfortunately, toothache and catch are not independent!

Combining Evidence

What happens if there are two or more pieces of evidence?

$\mathcal{P}(\text{Cavity} \mid \text{toothache} \wedge \text{catch}) = \alpha \langle 0.108, 0.016 \rangle$ (from full joint distribution)

Try Bayes' rule:

$\mathcal{P}(\text{Cavity} \mid \text{toothache} \wedge \text{catch}) = \alpha \mathcal{P}(\text{toothache} \wedge \text{catch} \mid \text{Cavity}) \mathcal{P}(\text{Cavity})$

Unfortunately, no better than full joint distribution

Independence: unfortunately, toothache and catch are not independent!

However, they are independent **given** the presence/absence of a cavity

$\mathcal{P}(\text{toothache} \wedge \text{catch} \mid \text{Cavity}) = \alpha \mathcal{P}(\text{toothache} \mid \text{Cavity}) \mathcal{P}(\text{catch} \mid \text{Cavity})$

Combining Evidence

What happens if there are two or more pieces of evidence?

$\mathcal{P}(\text{Cavity} \mid \text{toothache} \wedge \text{catch}) = \alpha \langle 0.108, 0.016 \rangle$ (from full joint distribution)

Try Bayes' rule:

$\mathcal{P}(\text{Cavity} \mid \text{toothache} \wedge \text{catch}) = \alpha \mathcal{P}(\text{toothache} \wedge \text{catch} \mid \text{Cavity}) \mathcal{P}(\text{Cavity})$

Unfortunately, no better than full joint distribution

Independence: unfortunately, toothache and catch are not independent!

However, they are independent **given** the presence/absence of a cavity

$\mathcal{P}(\text{toothache} \wedge \text{catch} \mid \text{Cavity}) = \alpha \mathcal{P}(\text{toothache} \mid \text{Cavity}) \mathcal{P}(\text{catch} \mid \text{Cavity})$

Conditional Independence of two variables X and Y , given a third variable Z , is

$$\mathcal{P}(X, Y \mid Z) = \mathcal{P}(X \mid Z) \mathcal{P}(Y \mid Z)$$

Combining Evidence

What happens if there are two or more pieces of evidence?

$\mathcal{P}(\text{Cavity} \mid \text{toothache} \wedge \text{catch}) = \alpha \langle 0.108, 0.016 \rangle$ (from full joint distribution)

Try Bayes' rule:

$\mathcal{P}(\text{Cavity} \mid \text{toothache} \wedge \text{catch}) = \alpha \mathcal{P}(\text{toothache} \wedge \text{catch} \mid \text{Cavity}) \mathcal{P}(\text{Cavity})$

Unfortunately, no better than full joint distribution

Independence: unfortunately, toothache and catch are not independent!

However, they are independent **given** the presence/absence of a cavity

$\mathcal{P}(\text{toothache} \wedge \text{catch} \mid \text{Cavity}) = \alpha \mathcal{P}(\text{toothache} \mid \text{Cavity}) \mathcal{P}(\text{catch} \mid \text{Cavity})$

Conditional Independence of two variables X and Y , given a third variable Z , is

$$\mathcal{P}(X, Y \mid Z) = \mathcal{P}(X \mid Z) \mathcal{P}(Y \mid Z)$$

With this, the size of the representation grows as $O(n)$

Combining Evidence

What happens if there are two or more pieces of evidence?

$\mathcal{P}(\text{Cavity} \mid \text{toothache} \wedge \text{catch}) = \alpha \langle 0.108, 0.016 \rangle$ (from full joint distribution)

Try Bayes' rule:

$\mathcal{P}(\text{Cavity} \mid \text{toothache} \wedge \text{catch}) = \alpha \mathcal{P}(\text{toothache} \wedge \text{catch} \mid \text{Cavity}) \mathcal{P}(\text{Cavity})$

Unfortunately, no better than full joint distribution

Independence: unfortunately, toothache and catch are not independent!

However, they are independent **given** the presence/absence of a cavity

$\mathcal{P}(\text{toothache} \wedge \text{catch} \mid \text{Cavity}) = \alpha \mathcal{P}(\text{toothache} \mid \text{Cavity}) \mathcal{P}(\text{catch} \mid \text{Cavity})$

Conditional Independence of two variables X and Y , given a third variable Z , is

$$\mathcal{P}(X, Y \mid Z) = \mathcal{P}(X \mid Z) \mathcal{P}(Y \mid Z)$$

With this, the size of the representation grows as $O(n)$

The decomposition of probabilistic domains through conditional independence is one of the most important developments in AI

Naive Bayes Models

Naive Bayes Models

A single cause influences a number of effects, all of which are conditionally independent:

Naive Bayes Models

A single cause influences a number of effects, all of which are conditionally independent:

$$\mathcal{P}(\text{Cause}, \text{Effect}_1, \dots, \text{Effect}_n) = \mathcal{P}(\text{Cause}) \prod_i \mathcal{P}(\text{Effect}_i | \text{cause})$$

Naive Bayes Models

A single cause influences a number of effects, all of which are conditionally independent:

$$\mathcal{P}(\text{Cause}, \text{Effect}_1, \dots, \text{Effect}_n) = \mathcal{P}(\text{Cause}) \prod_i \mathcal{P}(\text{Effect}_i | \text{cause})$$

It is called **naive** Bayes models (sometimes Bayesian Classifier) because it is often used where the effects are not independent

Naive Bayes Models

A single cause influences a number of effects, all of which are conditionally independent:

$$\mathcal{P}(\text{Cause}, \text{Effect}_1, \dots, \text{Effect}_n) = \mathcal{P}(\text{Cause}) \prod_i \mathcal{P}(\text{Effect}_i | \text{cause})$$

It is called **naive** Bayes models (sometimes Bayesian Classifier) because it is often used where the effects are not independent

Let $\mathbf{E} = \mathbf{e}$ are the observed effects and \mathbf{Y} are unobserved

$$\begin{aligned}\mathcal{P}(\text{Cause} | \mathbf{e}) &= \alpha \sum_{\mathbf{y}} \mathcal{P}(\text{Cause}, \mathbf{e}, \mathbf{y}) \\ &= \alpha \mathcal{P}(\text{Cause}) \prod_j \mathcal{P}(e_j | \text{Cause})\end{aligned}$$

Naive Bayes Models

A single cause influences a number of effects, all of which are conditionally independent:

$$\mathcal{P}(\text{Cause}, \text{Effect}_1, \dots, \text{Effect}_n) = \mathcal{P}(\text{Cause}) \prod_i \mathcal{P}(\text{Effect}_i | \text{cause})$$

It is called **naive** Bayes models (sometimes Bayesian Classifier) because it is often used where the effects are not independent

Let $\mathbf{E} = \mathbf{e}$ are the observed effects and \mathbf{Y} are unobserved

$$\begin{aligned}\mathcal{P}(\text{Cause} | \mathbf{e}) &= \alpha \sum_{\mathbf{y}} \mathcal{P}(\text{Cause}, \mathbf{e}, \mathbf{y}) \\ &= \alpha \mathcal{P}(\text{Cause}) \prod_j \mathcal{P}(e_j | \text{Cause})\end{aligned}$$

Text classification using the naive Bayes models:

Naive Bayes Models

A single cause influences a number of effects, all of which are conditionally independent:

$$\mathcal{P}(\text{Cause}, \text{Effect}_1, \dots, \text{Effect}_n) = \mathcal{P}(\text{Cause}) \prod_i \mathcal{P}(\text{Effect}_i | \text{cause})$$

It is called **naive** Bayes models (sometimes Bayesian Classifier) because it is often used where the effects are not independent

Let $\mathbf{E} = \mathbf{e}$ are the observed effects and \mathbf{Y} are unobserved

$$\begin{aligned} \mathcal{P}(\text{Cause} | \mathbf{e}) &= \alpha \sum_{\mathbf{y}} \mathcal{P}(\text{Cause}, \mathbf{e}, \mathbf{y}) \\ &= \alpha \mathcal{P}(\text{Cause}) \prod_j \mathcal{P}(e_j | \text{Cause}) \end{aligned}$$

Text classification using the naive Bayes models:

- given a text, decide on which (predefined) categories it belongs

Naive Bayes Models

A single cause influences a number of effects, all of which are conditionally independent:

$$\mathcal{P}(\text{Cause}, \text{Effect}_1, \dots, \text{Effect}_n) = \mathcal{P}(\text{Cause}) \prod_i \mathcal{P}(\text{Effect}_i | \text{cause})$$

It is called **naive** Bayes models (sometimes Bayesian Classifier) because it is often used where the effects are not independent

Let $\mathbf{E} = \mathbf{e}$ are the observed effects and \mathbf{Y} are unobserved

$$\begin{aligned}\mathcal{P}(\text{Cause} | \mathbf{e}) &= \alpha \sum_{\mathbf{y}} \mathcal{P}(\text{Cause}, \mathbf{e}, \mathbf{y}) \\ &= \alpha \mathcal{P}(\text{Cause}) \prod_j \mathcal{P}(e_j | \text{Cause})\end{aligned}$$

Text classification using the naive Bayes models:

- given a text, decide on which (predefined) categories it belongs
- The naive Bayes models consist of $\mathcal{P}(\text{Category})$ and the conditional probabilities $\mathcal{P}(\text{HasWord}_i | \text{Category})$

Naive Bayes Models

A single cause influences a number of effects, all of which are conditionally independent:

$$\mathcal{P}(\text{Cause}, \text{Effect}_1, \dots, \text{Effect}_n) = \mathcal{P}(\text{Cause}) \prod_i \mathcal{P}(\text{Effect}_i | \text{cause})$$

It is called **naive** Bayes models (sometimes Bayesian Classifier) because it is often used where the effects are not independent

Let $\mathbf{E} = \mathbf{e}$ are the observed effects and \mathbf{Y} are unobserved

$$\begin{aligned}\mathcal{P}(\text{Cause} | \mathbf{e}) &= \alpha \sum_{\mathbf{y}} \mathcal{P}(\text{Cause}, \mathbf{e}, \mathbf{y}) \\ &= \alpha \mathcal{P}(\text{Cause}) \prod_j \mathcal{P}(e_j | \text{Cause})\end{aligned}$$

Text classification using the naive Bayes models:

- given a text, decide on which (predefined) categories it belongs
- The naive Bayes models consist of $\mathcal{P}(\text{Category})$ and the conditional probabilities $\mathcal{P}(\text{HasWord}_i | \text{Category})$
- Categorize a new document: check which keywords appear in the document and then apply above equation to obtain the posterior probability distribution

Probabilistic Reasoning in Wumpus World

The sensor of an agent gets “Breeze” in [1,2] and [2,1]

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2 B OK	2,2	3,2	4,2
1,1 OK	2,1 B OK	3,1	4,1

Probabilistic Reasoning in Wumpus World

The sensor of an agent gets “Breeze” in [1,2] and [2,1]

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2 B OK	2,2	3,2	4,2
1,1 OK	2,1 B OK	3,1	4,1

Three unvisited but reachable rooms ([1,3],[2,2],[3,1]) may contain a pit

Probabilistic Reasoning in Wumpus World

The sensor of an agent gets “Breeze” in [1,2] and [2,1]

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2 B OK	2,2	3,2	4,2
1,1 OK	2,1 B OK	3,1	4,1

Three unvisited but reachable rooms ([1,3],[2,2],[3,1]) may contain a pit

The agent *must* move to one of these rooms

Probabilistic Reasoning in Wumpus World

The sensor of an agent gets “Breeze” in [1,2] and [2,1]

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2 B OK	2,2	3,2	4,2
1,1 OK	2,1 B OK	3,1	4,1

Three unvisited but reachable rooms ([1,3],[2,2],[3,1]) may contain a pit

The agent *must* move to one of these rooms
Logical agent cannot conclude which square most likely to be safe and it chooses randomly
Can probabilistic agent do better?

Probabilistic Reasoning in Wumpus World

The sensor of an agent gets “Breeze” in [1,2] and [2,1]

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2 B OK	2,2	3,2	4,2
1,1 OK	2,1 B OK	3,1	4,1

Three unvisited but reachable rooms ([1,3],[2,2],[3,1]) may contain a pit

The agent *must* move to one of these rooms
Logical agent cannot conclude which square most likely to be safe and it chooses randomly

Can probabilistic agent do better?

We know:

- a pit causes breezes in all neighboring rooms
- each room (except [1,1]) contains a pit with probability 0.2

Probabilistic Reasoning in Wumpus World

The sensor of an agent gets “Breeze” in [1,2] and [2,1]

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2 B OK	2,2	3,2	4,2
1,1 OK	2,1 B OK	3,1	4,1

Three unvisited but reachable rooms ([1,3],[2,2],[3,1]) may contain a pit

The agent *must* move to one of these rooms

Logical agent cannot conclude which square most likely to be safe and it chooses randomly

Can probabilistic agent do better?

We know:

- a pit causes breezes in all neighboring rooms
- each room (except [1,1]) contains a pit with probability 0.2

- For each square, $P[i,j]$ is TRUE iff room [i,j] contains a pit
- For each square, $B[i,j]$ is TRUE iff room [i,j] is breezy and we know the value of B_{11} , B_{12} , and B_{21} ,

Probabilistic Reasoning in Wumpus World

Specify the full joint distribution and apply the product rule:

Probabilistic Reasoning in Wumpus World

Specify the full joint distribution and apply the product rule:

$$\mathcal{P}(P_{11}, \dots, P_{44}, B_{11}, B_{12}, B_{21}) = \mathcal{P}(B_{11}, B_{12}, B_{21} \mid P_{11}, \dots, P_{44}) \mathcal{P}(P_{11}, \dots, P_{44})$$

Probabilistic Reasoning in Wumpus World

Specify the full joint distribution and apply the product rule:

$$\mathcal{P}(P_{11}, \dots, P_{44}, B_{11}, B_{12}, B_{21}) = P(\mathcal{B}_{11}, B_{12}, B_{21} \mid P_{11}, \dots, P_{44}) \mathcal{P}(P_{11}, \dots, P_{44})$$

First term: conditional probability distribution of a breeze, given a pit configuration.
One if all the breezy rooms adjacent to the pits, 0 otherwise.

Probabilistic Reasoning in Wumpus World

Specify the full joint distribution and apply the product rule:

$$\mathcal{P}(P_{11}, \dots, P_{44}, B_{11}, B_{12}, B_{21}) = \mathcal{P}(B_{11}, B_{12}, B_{21} \mid P_{11}, \dots, P_{44}) \mathcal{P}(P_{11}, \dots, P_{44})$$

First term: conditional probability distribution of a breeze, given a pit configuration. One if all the breezy rooms adjacent to the pits, 0 otherwise.

Second term: prior probability of a pit configuration. Each square contains a pit with probability 0.2 independent to each other

Probabilistic Reasoning in Wumpus World

Specify the full joint distribution and apply the product rule:

$$\mathcal{P}(P_{11}, \dots, P_{44}, B_{11}, B_{12}, B_{21}) = \mathcal{P}(B_{11}, B_{12}, B_{21} \mid P_{11}, \dots, P_{44}) \mathcal{P}(P_{11}, \dots, P_{44})$$

First term: conditional probability distribution of a breeze, given a pit configuration. One if all the breezy rooms adjacent to the pits, 0 otherwise.

Second term: prior probability of a pit configuration. Each square contains a pit with probability 0.2 independent to each other

$$\mathcal{P}(P_{11}, \dots, P_{44}) = \prod_{i,j=1,1}^{4,4} \mathcal{P}(P_{ij}). \text{ For world with } n \text{ pits, it is } 0.2^n \times 0.8^{16-n}.$$

Probabilistic Reasoning in Wumpus World

Specify the full joint distribution and apply the product rule:

$$\mathcal{P}(P_{11}, \dots, P_{44}, B_{11}, B_{12}, B_{21}) = \mathcal{P}(B_{11}, B_{12}, B_{21} \mid P_{11}, \dots, P_{44}) \mathcal{P}(P_{11}, \dots, P_{44})$$

First term: conditional probability distribution of a breeze, given a pit configuration. One if all the breezy rooms adjacent to the pits, 0 otherwise.

Second term: prior probability of a pit configuration. Each square contains a pit with probability 0.2 independent to each other

$$\mathcal{P}(P_{11}, \dots, P_{44}) = \prod_{i,j=1,1}^{4,4} \mathcal{P}(P_{ij}). \text{ For world with } n \text{ pits, it is } 0.2^n \times 0.8^{16-n}.$$

The evidence:

- the observed of breeze in each square that is visited, $b = \neg B_{11} \wedge B_{12} \wedge B_{21}$
- each visited square contains no pit, $\text{known} = \neg P_{11} \wedge \neg P_{12} \wedge \neg P_{21}$

Probabilistic Reasoning in Wumpus World

Specify the full joint distribution and apply the product rule:

$$\mathcal{P}(P_{11}, \dots, P_{44}, B_{11}, B_{12}, B_{21}) = \mathcal{P}(B_{11}, B_{12}, B_{21} \mid P_{11}, \dots, P_{44}) \mathcal{P}(P_{11}, \dots, P_{44})$$

First term: conditional probability distribution of a breeze, given a pit configuration. One if all the breezy rooms adjacent to the pits, 0 otherwise.

Second term: prior probability of a pit configuration. Each square contains a pit with probability 0.2 independent to each other

$$\mathcal{P}(P_{11}, \dots, P_{44}) = \prod_{i,j=1,1}^{4,4} \mathcal{P}(P_{ij}). \text{ For world with } n \text{ pits, it is } 0.2^n \times 0.8^{16-n}.$$

The evidence:

- the observed of breeze in each square that is visited, $b = \neg B_{11} \wedge B_{12} \wedge B_{21}$
- each visited square contains no pit, $\text{known} = \neg P_{11} \wedge \neg P_{12} \wedge \neg P_{21}$

Query: $\mathcal{P}(P_{13} \mid \text{known}, b)$ (How likely it is that [1,3] contains a pit?)

Probabilistic Reasoning in Wumpus World

Specify the full joint distribution and apply the product rule:

$$\mathcal{P}(P_{11}, \dots, P_{44}, B_{11}, B_{12}, B_{21}) = \mathcal{P}(B_{11}, B_{12}, B_{21} \mid P_{11}, \dots, P_{44}) \mathcal{P}(P_{11}, \dots, P_{44})$$

First term: conditional probability distribution of a breeze, given a pit configuration. One if all the breezy rooms adjacent to the pits, 0 otherwise.

Second term: prior probability of a pit configuration. Each square contains a pit with probability 0.2 independent to each other

$$\mathcal{P}(P_{11}, \dots, P_{44}) = \prod_{i,j=1,1}^{4,4} \mathcal{P}(P_{ij}). \text{ For world with } n \text{ pits, it is } 0.2^n \times 0.8^{16-n}.$$

The evidence:

- the observed of breeze in each square that is visited, $b = \neg B_{11} \wedge B_{12} \wedge B_{21}$
- each visited square contains no pit, $\text{known} = \neg P_{11} \wedge \neg P_{12} \wedge \neg P_{21}$

Query: $\mathcal{P}(P_{13} \mid \text{known}, b)$ (How likely it is that [1,3] contains a pit?)

$$\text{We have: } \mathcal{P}(P_{13} \mid \text{known}, b) = \alpha \sum_{\text{unknown}} \mathcal{P}(P_{13}, \text{known}, b, \text{unknown})$$

Probabilistic Reasoning in Wumpus World

$$\mathcal{P}(P_{13} \mid \text{known}, b) = \alpha \sum_{\text{unknown}} \mathcal{P}(P_{13}, \text{known}, b, \text{unknown})$$

The evidence:

- the observed of breeze in each square that is visited, $b = \neg B_{11} \wedge B_{12} \wedge B_{21}$
- each visited square contains no pit, $\text{known} = \neg P_{11} \wedge \neg P_{12} \wedge \neg P_{21}$

Query: $\mathcal{P}(P_{13} \mid \text{known}, b)$ (How likely it is that [1,3] contains a pit?)

We have: $\mathcal{P}(P_{13} \mid \text{known}, b) = \alpha \sum_{\text{unknown}} \mathcal{P}(P_{13}, \text{known}, b, \text{unknown})$

Probabilistic Reasoning in Wumpus World

$$\mathcal{P}(P_{13} \mid \text{known}, b) = \alpha \sum_{\text{unknown}} \mathcal{P}(P_{13}, \text{known}, b, \text{unknown})$$

There are 12 unknown room, hence the summation contains 2^{12} terms

Probabilistic Reasoning in Wumpus World

$$\mathcal{P}(P_{13} \mid \text{known}, b) = \alpha \sum_{\text{unknown}} \mathcal{P}(P_{13}, \text{known}, b, \text{unknown})$$

There are 12 unknown room, hence the summation contains 2^{12} terms


Fortunately, other (more distant) rooms are irrelevant!

Probabilistic Reasoning in Wumpus World

$$\mathcal{P}(P_{13} \mid \text{known}, b) = \alpha \sum_{\text{unknown}} \mathcal{P}(P_{13}, \text{known}, b, \text{unknown})$$

There are 12 unknown room, hence the summation contains 2^{12} terms

Fortunately, other (more distant) rooms are irrelevant!




Probabilistic Reasoning in Wumpus World

$$\mathcal{P}(P_{13} \mid \text{known}, b) = \alpha \sum_{\text{unknown}} \mathcal{P}(P_{13}, \text{known}, b, \text{unknown})$$

There are 12 unknown room, hence the summation contains 2^{12} terms

Fortunately, other (more distant) rooms are irrelevant!



Frontier: pit variables adjacent to visited rooms

Other: pit variables for other rooms


Unknown = Frontier \cup Other

Probabilistic Reasoning in Wumpus World

$$\mathcal{P}(P_{13} \mid \text{known}, b) = \alpha \sum_{\text{unknown}} \mathcal{P}(P_{13}, \text{known}, b, \text{unknown})$$

There are 12 unknown room, hence the summation contains 2^{12} terms

Fortunately, other (more distant) rooms are irrelevant!



Frontier: pit variables adjacent to visited rooms

Other: pit variables for other rooms

Unknown = Frontier \cup Other

Insight: the breezes are conditionally independent of the other variables, given the frontier and query variables

Probabilistic Reasoning in Wumpus World

$$\mathcal{P}(P_{13} \mid \text{known}, b) = \alpha \sum_{\text{unknown}} \mathcal{P}(P_{13}, \text{known}, b, \text{unknown})$$

Apply conditional independence:

$$\mathcal{P}(P_{13} \mid \text{known}, b) = \alpha \frac{\sum_{\text{frontier}} \mathcal{P}(b \mid \text{known}, P_{13}, \text{frontier})}{\sum_{\text{other}} \mathcal{P}(P_{13}, \text{known}, \text{frontier}, \text{other})}$$

Probabilistic Reasoning in Wumpus World

$$\mathcal{P}(P_{13} \mid \text{known}, b) = \alpha \sum_{\text{unknown}} \mathcal{P}(P_{13}, \text{known}, b, \text{unknown})$$

Apply conditional independence:

$$\mathcal{P}(P_{13} \mid \text{known}, b) = \alpha \frac{\sum_{\text{frontier}} \mathcal{P}(b \mid \text{known}, P_{13}, \text{frontier})}{\sum_{\text{other}} \mathcal{P}(P_{13}, \text{known}, \text{frontier}, \text{other})}$$

$$\mathcal{P}(P_{13} \mid \text{known}, b) = \alpha' \sum_{\text{frontier}} \mathcal{P}(b \mid \text{known}, P_{13}, \text{frontier}) \mathcal{P}(\text{frontier})$$

Probabilistic Reasoning in Wumpus World

$$\mathcal{P}(P_{13} \mid \text{known}, b) = \alpha \sum_{\text{unknown}} \mathcal{P}(P_{13}, \text{known}, b, \text{unknown})$$

Apply conditional independence:

$$\mathcal{P}(P_{13} \mid \text{known}, b) = \alpha \frac{\sum_{\text{frontier}} \mathcal{P}(b \mid \text{known}, P_{13}, \text{frontier})}{\sum_{\text{other}} \mathcal{P}(P_{13}, \text{known}, \text{frontier}, \text{other})}$$

$$\mathcal{P}(P_{13} \mid \text{known}, b) = \alpha' \sum_{\text{frontier}} \mathcal{P}(b \mid \text{known}, P_{13}, \text{frontier}) \mathcal{P}(\text{frontier})$$

The probabilities $\mathcal{P}(b \mid \text{known}, P_{13}, \text{frontier})$ are 1 when the breeze are consistent with other variables. For each value of P_{13} , we sum over the logical models for frontier that are consistent with the fact

Probabilistic Reasoning in Wumpus World


$$\mathcal{P}(P_{13} \mid \text{known}, b) = \alpha \sum_{\text{unknown}} \mathcal{P}(P_{13}, \text{known}, b, \text{unknown})$$

Apply conditional independence:

$$\mathcal{P}(P_{13} \mid \text{known}, b) = \alpha \frac{\sum_{\text{frontier}} \mathcal{P}(b \mid \text{known}, P_{13}, \text{frontier})}{\sum_{\text{other}} \mathcal{P}(P_{13}, \text{known}, \text{frontier}, \text{other})}$$

$$\mathcal{P}(P_{13} \mid \text{known}, b) = \alpha' \sum_{\text{frontier}} \mathcal{P}(b \mid \text{known}, P_{13}, \text{frontier}) \mathcal{P}(\text{frontier})$$

The probabilities $\mathcal{P}(b \mid \text{known}, P_{13}, \text{frontier})$ are 1 when the breeze are consistent with other variables. For each value of P_{13} , we sum over the logical models for frontier that are consistent with the fact



Probabilistic Reasoning in Wumpus World


$$\mathcal{P}(P_{13} \mid \text{known}, b) = \alpha \sum_{\text{unknown}} \mathcal{P}(P_{13}, \text{known}, b, \text{unknown})$$

Apply conditional independence:

$$\mathcal{P}(P_{13} \mid \text{known}, b) = \alpha \frac{\sum_{\text{frontier}} \mathcal{P}(b \mid \text{known}, P_{13}, \text{frontier})}{\sum_{\text{other}} \mathcal{P}(P_{13}, \text{known}, \text{frontier}, \text{other})}$$

$$\mathcal{P}(P_{13} \mid \text{known}, b) = \alpha' \sum_{\text{frontier}} \mathcal{P}(b \mid \text{known}, P_{13}, \text{frontier}) \mathcal{P}(\text{frontier})$$

The probabilities $\mathcal{P}(b \mid \text{known}, P_{13}, \text{frontier})$ are 1 when the breeze are consistent with other variables. For each value of P_{13} , we sum over the logical models for frontier that are consistent with the fact



$$P_{13} = P_{31} = \langle 0.31, 0.69 \rangle$$

$$P_{22} = \langle 0.86, 0.14 \rangle$$



**INFORMATIKA
UNPAR**



informatika.unpar.ac.id



informatika@unpar.ac.id



[if.unpar](https://www.facebook.com/if.unpar)



[if.unpar](https://www.instagram.com/if.unpar)