
2 ELEMENTS, COMPOUNDS AND MIXTURES

Most of the substances that we are familiar with from everyday life are mixtures. For example, the air that we breathe is a mixture containing elements such as nitrogen and oxygen, and compounds such as carbon dioxide and nitrogen oxides. The food that we eat and the drinks that we drink are mixtures. This chapter looks at the properties of elements, compounds and mixtures, and also how to separate the components of a mixture. Separation of mixtures is very important in the analysis of substances, such as in forensics.



▲ Figure 2.1 Gold is an element, but a gold ring made from 18-carat gold only contains 75% gold. The metal is a mixture of gold and, usually, copper.



▲ Figure 2.2 Pure water is a compound, but the water we drink is a mixture of water and other dissolved substances.

LEARNING OBJECTIVES

- Understand how to classify a substance as an element, compound or mixture
- Understand that a pure substance has a fixed melting and boiling point, but that a mixture may melt or boil over a range of temperatures
- Describe these experimental techniques for the separation of mixtures:
  - simple distillation
  - fractional distillation
  - filtration
  - crystallisation
  - paper chromatography
- Understand how a chromatogram provides information about the composition of a mixture
- Understand how to use the calculation of R_f values to identify the components of a mixture
- Practical: Investigate paper chromatography using inks/food colourings

REMINDER


You might want to look at Chapter 3 if you do not already know the term 'atom'.

KEY POINT


It isn't completely true to say that elements consist of only one type of atom. A better way of saying it would be that *all the atoms in an element have the same atomic number*. Most elements consist of mixtures of isotopes, which have the same atomic number, but different mass numbers (due to different numbers of neutrons). When we draw diagrams or make models, we aren't usually interested in the differences between the isotopes. Isotopes will be discussed in Chapter 3.

ELEMENTS

Elements are *substances that can't be split into anything simpler by chemical means*. An element contains only one type of atom (but see the key point in the margin). In models or diagrams they are shown as atoms of a single colour or size.



a pure metal such as magnesium



oxygen gas



diamond (a form of carbon)

▲ Figure 2.3 Elements contain only one type of atom.

There are 118 elements and these are shown in the Periodic Table. Most of the elements occur naturally, such as hydrogen, helium and sulfur. Some others have to be made artificially, such as einsteinium.

COMPOUNDS

Compounds are formed when *two or more elements chemically combine*. The elements always combine in fixed proportions. For example, hydrogen and fluorine always combine to form hydrogen fluoride, with formula HF, whereas magnesium and fluorine always combine to form magnesium fluoride, with formula MgF_2 – the elements must combine in these ratios. Examples of other compounds are carbon dioxide (CO_2) and methane (CH_4). Diagrams of compounds show more than one type of atom bonded together.




water | silicon dioxide | sodium chloride

▲ Figure 2.4 Some compounds

MIXTURES

In a **mixture**, the various substances are mixed together and no chemical reaction occurs. Mixtures can be made from elements and/or compounds. The various components can be in any proportion, for example you can put any amount of sugar into your cup of tea or coffee (until it becomes saturated).



mixture of elements – nitrogen and oxygen | mixture of compounds – carbon dioxide and water (vapour) | mixture of an element with a compound – carbon dioxide and nitrogen

▲ Figure 2.5 Some mixtures

SIMPLE DIFFERENCES BETWEEN MIXTURES AND COMPOUNDS

PROPORTIONS

In water (a compound), every single water molecule has two hydrogen atoms combined with one oxygen atom. It never varies. In a mixture of hydrogen and oxygen gases, the two could be mixed together in any proportion.

If you had some iron metal and some sulfur, you could mix them in any proportion you wanted to. In iron sulfide (FeS), a compound, the proportion of iron to sulfur is always exactly the same.

PROPERTIES

REMINDER

You can find out about the reactions of metals with dilute acids on pages 174–175. The reaction between iron sulfide and acids isn't needed for exam purposes at International GCSE.

In a mixture of elements, each element keeps its own properties, but the properties of the compound are quite different. For example, in a mixture of iron and sulfur, the iron is grey and the sulfur is yellow. The iron reacts with dilute acids such as hydrochloric acid to produce hydrogen; the sulfur doesn't react with the acid. However, the compound iron sulfide (FeS) reacts quite differently with acids to produce poisonous hydrogen sulfide gas, which smells of bad eggs.

A mixture of hydrogen and oxygen is a colourless gas which explodes when you put a flame to it. The compound, water, is a colourless liquid which just puts out a flame.

EASE OF SEPARATION

Mixtures can be separated by **physical means**. Physical means are things like changing the temperature or dissolving part of the mixture in a solvent such as water; in other words, methods that don't involve any chemical reactions.

For example, a mixture of iron and sulfur is quite easy to separate into the two elements using a magnet. The iron sticks to the magnet and the sulfur doesn't. The elements in a compound cannot be separated by physical means. To convert iron sulfide into separate samples of iron and sulfur requires chemical reactions.

You can cool a mixture of hydrogen and oxygen gases to separate it by a physical process. Oxygen condenses into a liquid at a much higher temperature than hydrogen (–183 °C as opposed to –253 °C). This would leave you with liquid oxygen and hydrogen gas, which are easy to separate. But to separate water into hydrogen and oxygen, you have to change it chemically using electrolysis. Electrolysis is explained in Chapter 10.

MELTING POINT AND BOILING POINT

Pure substances, such as elements and pure compounds, melt and boil at fixed temperatures. For example, the melting point of water is 0 °C and the boiling point 100 °C. However, mixtures usually melt or boil over a *range of temperatures*.

KEY POINT

A mixture is not a pure substance. If a sample contains only a small amount of an unwanted substance, the unwanted substance might be called an *impurity*.


The presence of impurities lowers the melting point of a substance and raises the boiling point. For instance, dissolving 10 g of common (table) salt (sodium chloride) in 1 litre of water lowers the melting point to about –0.6 °C and raises the boiling point to about 100.2 °C.

The melting point can be very useful in determining whether or not a substance is pure. If you continue to study chemistry you might carry out a practical experiment to make some aspirin. In order to determine whether your sample is pure or not you can measure the melting point. You would record the temperature at which your sample starts to melt, and then you would record the temperature at which it has fully melted to completely form a liquid. Aspirin is a white powder that melts at 138 °C. If the melting point of the sample you made is 128–134 °C you can see that it is quite impure because it melts over a wide range of temperature (below the melting point of pure aspirin).

SEPARATION OF MIXTURES

Separating mixtures is extremely important in chemistry. For example, we can see this in the processing of crude oil, in producing fresh water from salt water and in the enrichment of uranium. In forensic science, the components of a mixture usually have to be separated before they can be analysed.

FILTRATION



▲ Figure 2.6 Filtration can be used to separate a mixture of sand and water.

Filtration can be used to separate a solid from a liquid.

For example, sand can be separated from water by filtration. The apparatus for filtration is shown in Figure 2.6.


The substance left in the filter paper is called the **residue** and the liquid that comes through is called the **filtrate**.

Filtration can also be used to separate two solids from each other if only one of them is soluble in water (see below – rock salt).

CRYSTALLISATION

Crystallisation can be used to separate a solute from a solution. For example, it could be used to separate sodium chloride from a sodium chloride solution. The solution is heated in an evaporating basin to boil off some of the water until an almost saturated solution is formed. This can be tested by dipping a glass rod into the solution and seeing if crystals form quickly on its surface when it is removed. The Bunsen burner is then turned off and the crystals allowed to form as more water evaporates and the solution cools. The crystals can now be removed from the mixture by filtration.

The apparatus for crystallisation is shown in Figure 2.7.



▲ Figure 2.7 Crystallisation can be used to separate a solute from a solution.

MAKING PURE SALT FROM ROCK SALT



▲ Figure 2.8 Rock salt

We can use filtration and crystallisation to obtain pure salt from rock salt.

Rock salt consists of salt contaminated by various earthy or rocky impurities. These impurities aren't soluble in water.


If you crush the rock salt and mix it with hot water, the salt dissolves, but the impurities don't. The impurities can be filtered off, and remain on the filter paper. The filtrate is then a salt solution. The solid salt can be obtained from the solution by crystallisation.

This is typical of the way you can separate any mixture of two solids, one of which is soluble in water and one of which isn't.

SIMPLE DISTILLATION

Simple distillation can be used to separate the components of a solution. Although we can use crystallisation to separate sodium chloride from a sodium chloride solution, we can also collect the water if we use simple distillation.

The water boils and is condensed back to a liquid by the condenser. The salt remains in the flask.



▲ Figure 2.9 Distilling pure water from sodium chloride solution.

KEY POINT

Notice that water is always fed into the condenser at the lower end. That way it fills the condenser jacket better and if the flow of water stops for any reason the condenser jacket remains full of water.


You could, of course, collect the salt from the solution as well as collecting pure water. The sodium chloride solution eventually becomes so concentrated that the salt will crystallise out.

FRACTIONAL DISTILLATION

Fractional distillation is used to separate a mixture of liquids such as ethanol (alcohol) and water. Ethanol and water are completely miscible with each other. That means you can mix them together in any proportion and they will form a single liquid layer. You can separate them by taking advantage of their different boiling points: water boils at 100 °C, ethanol at 78 °C.

EXTENSION WORK

The fractionating column is often packed with glass beads or something similar, although the separation of ethanol and water in the lab works perfectly well just with an empty column. For reasons that are beyond International GCSE, a high surface area in the column helps separation of the two vapours. The ethanol produced by this experiment is about 96% pure. For complicated reasons, again beyond International GCSE, it is impossible to remove the last 4% of water by distillation.



▲ Figure 2.10 Fractional distillation

Both liquids boil, but by careful heating you can control the temperature of the column so that all the water condenses in the column and trickles back into the flask. Only the ethanol remains as a vapour all the way to the top of the fractionating column and out into the condenser.

PAPER CHROMATOGRAPHY

Paper chromatography can be used to separate a variety of mixtures. However, at International GCSE level we will usually use it to separate mixtures of coloured inks or food colourings. Most inks and food colourings are not just made up of one colour but contain a mixture of dyes.

Paper chromatography can also be used to separate a mixture of colourless substances such as sugars, but then some method must be used to make the spots visible on the paper.

ACTIVITY 2

▼ PRACTICAL: INVESTIGATING THE COMPOSITION OF DYE WITH PAPER CHROMATOGRAPHY

Safety Note: Avoid skin contact with the solvents and dyes, especially if you have sensitive skin.

We can investigate the composition of a mixture of coloured dyes using paper chromatography. To do this we carry out the following steps.

1. Draw a line with a pencil across a piece of chromatography paper; this line should be about 1 cm from the bottom of the paper. Do not use a pen as the colours in the ink may move up the chromatography paper with the solvent.
2. Put a spot (use a teat pipette or a capillary tube) of the mixture of dyes on the pencil line and allow it to dry.
3. Suspend the chromatography paper in a beaker that contains a small amount of solvent so that the bottom of the paper goes into the solvent. It is important that the solvent is below the pencil line so that the inks/colourings don't just dissolve in the solvent.
4. Put a lid (such as a watch glass) on the beaker so that the atmosphere becomes saturated with the solvent. This is to stop evaporation of the solvent from the surface of the paper.
5. When the solvent has moved up the paper to about 1 cm from the top, remove the paper from the beaker and draw a pencil line to show where the solvent got to. The highest level of the solvent on the paper at any time is called the *solvent front*.
6. Leave the paper to dry so that all the solvent evaporates.

For the solvent you can use water or a non-aqueous solvent (a solvent other than water). Which solvent you use depends on what substances are present in the mixture. A suitable solvent is usually found by experimenting with different ones.


The dyes that make up the mixture will be different in two important ways:

- the affinity they have for the paper (how well they 'stick' to the paper)
- how soluble they are in the solvent which moves up the paper.

In Figure 2.11 spot C has hardly moved. Either it was not very soluble in the solvent or it has a very high affinity for the paper (or both). On the other hand, spot A has moved almost as far as the solvent. It must be very soluble in the solvent and not have much affinity for the paper. The pattern you get is called a **chromatogram**.

KEY POINT

If the dye does not move from the pencil line during an experiment, then the dye is not at all soluble in the solvent you are using. In this case, you need to find a different solvent. If the dye moves up the paper with the solvent front, the dye is too soluble in that solvent and, again, you have to try a different solvent.



◀ Figure 2.11 Paper chromatography

In this example, the mixture must have contained a minimum of three different dyes. We say a *minimum* of three dyes because there could be more – it is possible that one of the spots is made up of two coloured dyes that by concidence moved the same distance. You could only confirm this by doing the experiment again with a different solvent.




▲ Figure 2.12 A paper chromatography experiment

USING PAPER CHROMATOGRAPHY IN ANALYSIS

You can use paper chromatography to identify the particular dyes in a mixture. If you think that your mixture (m) could contain dyes d1, d2, d3 and d4, you can carry out an experiment to determine this.

A pencil line is drawn on a larger sheet of paper and pencil marks are drawn along the line to show the original positions of the various dyes placed on the line (see Figure 2.13). One spot is your unknown mixture; the others are single, known dyes. The chromatogram is then allowed to develop as before.



▲ Figure 2.13 Paper chromatography can be used to analyse a mixture. Lines will not be present on your paper, but they have been added here to help you measure the distances.

The mixture (m) has spots corresponding to dyes d1, d3 and d4. They have the same colour as spots in the mixture, and have travelled the same distance on the paper. Although dye d2 is the same colour as one of the spots in the mixture, it has travelled a different distance and so must be a different compound.

Instead of just saying the spots move different distances we can use the R_f value to describe how far the spots move. R_f stands for **retardation factor**. Each time we do a chromatography experiment the solvent (and therefore the spots) will move different distances along the paper. This means we can't just report the distance moved by a particular spot so we have to work out a ratio instead.

$$R_f = \frac{\text{distance moved by a spot (from the pencil line)}}{\text{distance moved by the solvent front (from the pencil line)}}$$

In Figure 2.13 $R_f = \frac{x}{y}$.

So in Figure 2.13 the R_f value for dye d3 is:

$$R_f = \frac{2.9\,\text{cm}}{3.6\,\text{cm}} = 0.81$$

HINT

Measure to the centre of the spot.

The R_f values of the dyes in mixture m are:

blue spot: $R_f = \frac{0.9}{3.6} = 0.25$

orange spot: $R_f = \frac{2.0}{3.6} = 0.56$

green spot: $R_f = \frac{2.9}{3.6} = 0.81$

The R_f values of dyes d1 to d4 are:

d1: $R_f = 0.56$

d2: $R_f = 0.36$

d3: $R_f = 0.81$

d4: $R_f = 0.25$

Because the spots in mixture m have the same R_f values as d1, d3 and d4, we can conclude that the mixture contains these dyes.

An R_f value must be between 0 and 1. If you get a number bigger than 1 you have probably divided the numbers the wrong way round. An R_f value has no units.

You have to be careful when using R_f values as they depend on the solvent used and on the type of paper. There was no problem in the experiment described above because the mixture and the individual dyes were all put on the same piece of paper. However, if the mixture was put on one piece of chromatography paper and the individual dyes on a separate piece, you can still compare R_f values as long as you use the same type of paper and the same solvent.

CHAPTER QUESTIONS

SKILLS CRITICAL THINKING


1 Classify each of the following substances as an element, compound or mixture:

sea water	hydrogen	honey
magnesium oxide	copper(II) sulfate	blood
calcium	mud	potassium iodide solution


SKILLS ANALYSIS

2 Look at the diagrams below and classify each one as an element, compound or mixture.


a




b




c




d



e



f



SKILLS REASONING, PROBLEM SOLVING

3 A teacher has found two white powders on a desk in the chemistry laboratory. She wants to test to see if they are pure substances, so she measures the melting points. Substance X melts at 122 °C and substance Y melts between 87 and 93 °C. Explain which one is the pure substance.

SKILLS DECISION MAKING

4 State which separation method you would use to carry out the following separations:

a Potassium iodide from a potassium iodide solution.

b Water from a potassium iodide solution.

c Ethanol from a mixture of ethanol and water.

d Red dye from a mixture of red and blue dyes.


e Calcium carbonate (insoluble in water) from a mixture of calcium carbonate and water.

SKILLS CREATIVITY, DECISION MAKING

5 Suppose you had a valuable collection of small diamonds, which you kept safe from thieves by mixing them with white sugar crystals. You store the mixture in a jar labelled 'sugar'. Now you want to sell the diamonds. Describe how you would separate all the diamonds from the sugar.

SKILLS ANALYSIS 5

6 In order to identify the writer of an anonymous letter, a sample of ink from the letter was dissolved in a solvent and then placed on some chromatography paper. Spots of ink from the pens of five possible writers, **G**, **M**, **P**, **R** and **T**, were placed next to the sample on the chromatography paper. The final chromatogram looked like this:



a Which of the five writers is using ink that matches the sample from the letter?

b Which of the writers is using a pen that contains ink made from a single dye?

SKILLS PROBLEM SOLVING 6

c What is the R_f value of the blue dye in suspect **P**'s pen?

SKILLS ANALYSIS 5

d Which two of the five writers are using pens containing the same ink?

6

e Whose pen contained the dye that was most soluble in the solvent?