8 COVALENT BONDING

We met ionic compounds in Chapter 7 and in this chapter we will understand what covalent bonding is. There are a lot more covalent compounds than ionic compounds so it is important that you understand how the bonding works.



▲ Figure 8.1 Water is a covalent compound but the salt dissolved in sea water is an ionic compound.

LEARNING OBJECTIVES

- Know that a covalent bond is formed between atoms by the sharing of a pair of electrons
- Understand covalent bonds in terms of electrostatic attractions
- Understand how to use dot-and-cross diagrams to represent covalent bonds in:
  - diatomic molecules, including hydrogen, oxygen, nitrogen, halogens and hydrogen halides
  - inorganic molecules, including water, ammonia and carbon dioxide
  - organic molecules containing up to two carbon atoms, including methane, ethane, ethene and those containing halogen atoms
- Explain why substances with a simple molecular structure are gases or liquids, or solids with low melting and boiling points
- Understand that the term intermolecular forces of attraction can be used to represent all forces between molecules
- Explain why the melting and boiling points of substances with simple molecular structures increase, in general, with increasing relative molecular mass
- Explain why substances with giant covalent structures are solids with high melting and boiling points
- Explain how the structures of diamond, graphite and C_{60} fullerene influence their physical properties, including electrical conductivity and hardness
- Know that covalent compounds do not usually conduct electricity


COVALENT BONDING

WHAT IS A COVALENT BOND?

In any bond, particles are held together by electrostatic attractions between something positively charged and something negatively charged. In a **covalent bond**, *a pair of electrons is shared between two atoms*. What holds the atoms together is *the strong electrostatic attraction between the nuclei (positively charged) of the atoms that make up the bond, and the shared pair of electrons (negatively charged).*

EXTENSION WORK

In most of the simple examples you will meet at International GCSE, each atom in a covalent bond supplies one electron to the shared pair of electrons. That doesn't have to be the case. Both electrons may come from the same atom.



▲ Figure 8.2 The shared pair of electrons is attracted to the nuclei of both atoms.


COVALENT BONDING IN A HYDROGEN MOLECULE

HINT

Remember that, although the electrons are drawn as dots or as crosses, there is absolutely no difference between them in reality; the dots and the crosses simply show that the electrons have come from two different atoms.

Covalent bonds are often shown using dot-and-cross diagrams.

Both hydrogen nuclei in Figure 8.3 are strongly attracted to the shared pair of electrons.



▲ Figure 8.3 A dot-and-cross diagram for H_2

Hydrogen atoms form diatomic molecules with the formula H_2. The atoms in an H_2 molecule are joined together by a covalent bond. The covalent bond between two hydrogen atoms is very strong.

Molecules contain a certain fixed number of atoms, which are joined together by covalent bonds. Hydrogen molecules are said to be **diatomic** because they contain two atoms. Other sorts of molecule may have as many as thousands of atoms joined together, for example proteins and DNA.

THE SIGNIFICANCE OF NOBLE GAS STRUCTURES IN COVALENT BONDING

In H_2, each hydrogen atom has only one electron to share, so it can only form one covalent bond. The shared pair of electrons is in the outer shell of both, therefore each atom has the same number of electrons as a noble gas atom (helium in this case).

Does that mean that the hydrogen has turned into helium? No. The number of protons in the nucleus hasn't changed, it is the number of protons that defines what an atom is.

REMINDER

If atoms have 8 electrons in their outer shell they have the same number of electrons as a noble gas atom – they are isoelectronic with a noble gas atom.

In virtually all of the molecules you will meet at this level, electrons will be shared so that H atoms have a total of 2 electrons in their outer shell and all other atoms will have 8 electrons in their outer shell. Some people talk about the '**octet** rule', referring to this 8. Remember, we are counting shared electrons as belonging to the outer shells of both atoms.

When there is one atom in the middle and other atoms are joined to it (as in CH_4 or PCl_3) the *outer* atoms will virtually always have 8 electrons in their outer shell (or 2 if they are H). In fact, it is very difficult to think of an example where the outer atoms do not have 8 electrons. There are some molecules where the *central* atom does not have 8 electrons in the outer shell, and we will look at a couple of examples of those later on.

KEY POINT

In chemistry, we talk about things being more stable or less stable. When we do this we are usually talking about how much energy something has. Generally, the lower the energy something has, the more stable it is. Chemical reactions usually occur so that something becomes more stable. Think about holding a book: if you let go of the book, it will fall to the floor, where it has less (potential) energy, therefore it becomes more stable. When bonds form energy is given out and so the substance formed has less energy, therefore it is more stable.

WHY DOES HYDROGEN FORM MOLECULES?


Whenever a bond is formed (of whatever kind), energy is released, and that makes the things involved more stable than they were before. The more bonds an atom can form, the more energy is released and the more stable the system becomes. The H_2 molecule is much more stable than two separate hydrogen atoms.

COVALENT BONDING IN A HYDROGEN CHLORIDE MOLECULE

A chlorine atom has 7 electrons in its outer shell. By sharing 1 electron with a hydrogen atom, both atoms will have the same number of electrons as the nearest noble gas atom. If you look at the arrangement of electrons around the chlorine atom in the covalently bonded molecule of HCl (Figure 8.4), you will see that its electronic configuration is now [2, 8, 8]. That is the same as an argon atom. Similarly, the hydrogen now has 2 electrons in its outer shell – the same as helium.

Notice in Figure 8.4 that only the electrons in the outer shell of the chlorine are used in bonding. In the examples you will meet at International GCSE, the inner electrons never get used. In fact, the inner electrons are often left out of bonding diagrams. But be careful! In an exam, only leave out the inner electrons if the question tells you to. Another way of representing the covalent bonding in HCl is shown in Figure 8.5.


We also use lines to represent the covalent bonds between atoms, but be careful, the diagram shown in Figure 8.6 is not a dot-and-cross diagram.



▲ Figure 8.4 A dot-and-cross diagram for HCl.




▲ Figure 8.5 The covalent bonding in HCl showing outer shell electrons only.



▲ Figure 8.6 The line between the atoms represents a covalent bond.


COVALENT BONDING IN A CHLORINE MOLECULE (Cl_2)

A chlorine atom has 7 electrons in its outer shell. Each Cl shares 1 electron so that both Cl atoms will have 8 electrons in their outer shell.



▲ Figure 8.7 Two ways of showing the covalent bonding in Cl_2


COVALENT BONDING IN METHANE, AMMONIA AND WATER



▲ Figure 8.8 The electronic configurations of C, N and O atoms

COVALENT BONDING IN METHANE

A carbon atom has 4 electrons in its outer shell. By sharing 1 electron with each of 4 hydrogen atoms the C will have 8 electrons in its outer shell and each H will have 2 electrons in its outer shell. Therefore C forms 4 covalent bonds, 1 with each H atom. Methane has the formula CH_4.




▲ Figure 8.9 Dot-and-cross diagrams for methane

COVALENT BONDING IN AMMONIA

A nitrogen atom has 5 electrons in its outer shell. It will therefore share 3 other electrons to have 8 electrons in its outer shell. In ammonia, a nitrogen atom forms 3 covalent bonds, 1 with each H atom.


The formula of ammonia is NH_3.



▲ Figure 8.10 Dot-and-cross diagrams for ammonia

COVALENT BONDING IN WATER

An oxygen atom has 6 electrons in its outer shell. It will therefore share 2 other electrons to have 8 electrons in its outer shell. In water, an oxygen atom forms 2 covalent bonds – 1 with each H atom. The formula of water is H_2O.



▲ Figure 8.11 Dot-and-cross diagrams for water

LOOKING AHEAD

Shapes of molecules



▲ Figure 8.12 (a) A methane molecule is tetrahedral and (b) a water molecule is bent.

By understanding the bonding in a covalent molecule it is possible to work out the shape of the molecule. Pairs of electrons in the outer shell of the central atom repel each other and will therefore tend to get as far apart as possible. For example, in a methane molecule there are four pairs of electrons around the central C atom and for these to be as far away from each other as possible, they must be arranged in a tetrahedral shape. A tetrahedron (adjective: tetrahedral) is a triangular pyramid.

There are also four pairs of electrons around the central atom in water: two of these are pairs of electrons involved in covalent bonds and two are pairs of electrons in the outer shell of the oxygen which are not involved in bonding (these are often called *lone pairs of electrons* or just *lone pairs*). These four pairs of electrons are also arranged in a tetrahedral arrangement so the actual shape of a water molecule (how the atoms are arranged) is described as 'bent' or 'V-shaped'. The fact that a water molecule is bent and the electrons are attracted to a different extent by the oxygen and hydrogen atoms means that a water molecule is polar (has a slightly negative and a slightly positive end) and that a stream of water can be bent by an electrically charged object.



▲ Figure 8.13 Water being bent by an electrically charged comb.


COVALENT BONDING IN A SLIGHTLY MORE COMPLICATED MOLECULE: ETHANE

Ethane has the formula C_2H_6. The bonding is similar to methane (Figure 8.9), except that there is a carbon–carbon covalent bond as well as the carbon–hydrogen bonds.

This is called an organic compound. You will learn more about molecules such as this in Unit 4.

HINT

When drawing molecules containing carbon and hydrogen it is useful to remember that carbon always forms 4 bonds by sharing 4 electrons, and hydrogen always forms 1 bond by sharing 1 electron. The hydrogen atoms always go on the outside, never in the middle.




▲ Figure 8.14 A dot-and-cross diagram for ethane

MULTIPLE COVALENT BONDING

COVALENT BONDING IN AN OXYGEN MOLECULE: DOUBLE BONDING


An oxygen atom has 6 electrons in its outer shell and so if two oxygen atoms combine, they will both share 2 electrons each; this means that each atom will have 8 electrons in its outer shell. There are therefore two shared pairs of electrons between the oxygen atoms; this is called a double covalent bond or, usually, just a **double bond**.



▲ Figure 8.15 (a) A dot-and-cross diagram for O_2 and (b) the double covalent bond.

THE TRIPLE BOND IN A NITROGEN MOLECULE

A nitrogen atom has 5 electrons in its outer shell and so if two nitrogen atoms combine they will both share 3 electrons each; this means that each atom will have 8 electrons in its outer shell. There are, therefore, three shared pairs of electrons between the nitrogen atoms; this is called a triple covalent bond or, usually, just a **triple bond**.




▲ Figure 8.16 N_2 has a triple bond

Nitrogen gas consists of nitrogen molecules bonded like this. The triple bond from the sharing of three pairs of electrons between the two nitrogen atoms is very strong and needs a lot of energy to break. That is why nitrogen is relatively unreactive.

COVALENT DOUBLE BONDING IN CARBON DIOXIDE, CO_2

An oxygen atom has 6 electrons in its outer shell and a carbon atom has 4. Each oxygen atom will share 2 electrons with the carbon atom. Two double bonds are formed between the carbon and the two oxygens (Figure 8.17). All atoms have 8 electrons in their outer shells.




▲ Figure 8.17 CO_2 has two double bonds between atoms.

THE DOUBLE BOND IN ETHENE, C_2H_4

Ethene is rather like ethane in Figure 8.14, except that it only has two hydrogen atoms attached to each carbon atom and a double bond between the carbon atoms.

HINT


With organic compounds such as ethane and ethene you have to look at their names very carefully, even one different letter in the name can matter. Here, for example, ethane and ethene are completely different compounds. You will find out more about this in Unit 4.




▲ Figure 8.18 Ethene has a double bond between the C atoms.

ORGANIC MOLECULES CONTAINING HALOGEN ATOMS

Bromomethane has the formula CH_3Br: the 3 H atoms and the Br atom are joined to the central C atom. Br has 35 electrons and we have not learned how to work out the electronic configuration of an atom with 35 electrons, but if




▲ Figure 8.19 A dot-and-cross diagram for CH_3Br showing outer shells only. Although we have used different colours here for clarity, you do not need to use different colours in the exam.



▲ Figure 8.21 The structure of chloroethene showing the covalent bonds. This is not, however, a dot-and-cross diagram and you must not draw this if you are asked to draw a dot-and-cross diagram.

you look at the Periodic Table you will see that Br is in Group 7. Because it is in Group 7 we know that it has 7 electrons in its outer shell, and will share 1 electron so that it has 8 in its outer shell. Therefore, we know that Br will form just 1 covalent bond. When you draw a dot-and-cross diagram for CH_3Br you will only be asked to show the outer electrons (Figure 8.19).

Probably the most complicated molecule for which you could be asked to draw a dot-and-cross diagram would be something like chloroethene (CH_2CHCl). When drawing chloroethene, remember that C will form 4 covalent bonds, H will form 1 and Cl will form 1. The dot-and-cross diagram (outer shells only) is shown in Figure 8.20.



▲ Figure 8.20 There is a double bond between the 2 C atoms. It does not matter where you put the H and Cl atoms relative to the C, and you do not have to use different colours.


The structure of chloroethene can also be shown as illustrated in Figure 8.21.

SOME MORE DIFFICULT MOLECULES WHERE THE CENTRAL ATOM DOES NOT HAVE 8 ELECTRONS IN ITS OUTER SHELL

Although the *outer* atoms in a molecule will always have 8 (or 2 in the case of hydrogen) electrons in their outer shell, there are a few examples you may encounter where the *central* atom has more or fewer than 8 electrons in the outer shell.

BF_3

Boron is the central atom and the Fs are the outer atoms. Each F will share 1 other electron to have 8 electrons in its outer shell. This means that B only has a total of 6 electrons in its outer shell. Another way of thinking about this is that a B atom only has 3 electrons in its outer shell, and so this is the maximum number that it can share.



▲ Figure 8.22 A dot-and-cross diagram for BF_3

SO_2

This is called sulfur dioxide or sulfur(IV) oxide. The central atom is S and the outer atoms are O. Each O atom has 6 electrons in its outer shell and so will need to share 2 other electrons to have 8 electrons in the outer shell. So the S atom shares 2 electrons with each of the O atoms to form two double bonds. A sulfur atom originally had 6 electrons in its outer shell and so now, if it shares 4 electrons, it has 10 electrons in its outer shell.




▲ Figure 8.23 There are two double bonds in SO_2.

EXTENSION WORK

Atoms in Periods 3 and below (4, 5, 6, 7) can have more than 8 electrons in their outer shells. The maximum number of electrons that the central atom in a molecule can share is equal to the number of electrons in its outer shell. So, sulfur can form up to 6 bonds (in SF_6) and chlorine 7 (in $HClO_4$).

SIMPLE MOLECULAR STRUCTURES


Molecules contain fixed numbers of atoms joined by strong covalent bonds. If we look closely at liquid water, there are individual water molecules, where the H and O atoms are joined together with strong covalent bonds. But there must also be some forces between water molecules which keep them in the liquid state. These forces are **intermolecular forces**.



▲ Figure 8.24 Water is a simple molecular compound.

DID YOU KNOW?

Intermolecular literally means between molecules.



▲ Figure 8.25 Only intermolecular forces are broken when water evaporates/boils.

These intermolecular forces between molecules are much weaker than covalent bonds. When we boil water it is only these weak intermolecular forces of attraction that are broken; covalent bonds are not broken. Looking at Figure 8.25 you can see that there are H_2O molecules in liquid water and in gaseous water. The covalent bonds between the H and O atoms in the molecules have not changed in any way. All that has changed in gaseous water is that there are no intermolecular forces, they have been broken.

When a substance consists of molecules with intermolecular forces of attraction between them, we say that it has a *simple molecular structure*. Examples of things that have simple molecular structures are H_2O, CO_2, CH_4, NH_3 and C_2H_4. These are all the things that we have drawn dot-and-cross diagrams for above. Virtually all the compounds you will encounter that have covalent bonding will have simple molecular structures.

Substances with simple molecular structures tend to be gases or liquids or solids with low melting points and boiling points. The reason for this is that not much energy is required to break the weak intermolecular forces of attractions between molecules. Remember, no covalent bonds are broken, covalent bonds are strong.

EXTENSION WORK

If you continue with chemistry you will learn that there are different types of intermolecular forces. You will come across terms like van der Waals' forces, London dispersion forces and hydrogen bonds. There is a special type of intermolecular force between water molecules called **hydrogen bonds**. Hydrogen bonding gives water some of its very special properties, for example the solid form (ice) is less dense than the liquid form.

MELTING AND BOILING POINTS INCREASE AS RELATIVE MOLECULAR MASS INCREASES

The halogens (Group 7 in the Periodic Table) all have a simple molecular structure consisting of diatomic molecules with intermolecular forces between them.

You can see from Table 8.1 that the melting points and boiling points increase as the relative molecular mass increases.

Table 8.1 The melting points and boiling points of the halogens.

Halogen	Formula	Relative molecular mass/M_r	Melting point/°C	Boiling point/°C
fluorine	F_2	38	−220	−188
chlorine	Cl_2	71	−101	−34
bromine	Br_2	160	−7	59
iodine	I_2	254	114	184

EXTENSION WORK

It is not always the case that melting and boiling points increase as the M_r increases and really the rule only applies to sets of very similar substances, such as the halogens or the alkanes (see Chapter 23). Some examples where the rule does not work are water (M_r = 18, boiling point = 100 °C), ethane (M_r = 30, boiling point = –89 °C), NH_3 (M_r = 17, boiling point = –33 °C) and PH_3 (M_r = 34, boiling point = –88 °C).

Remember, as we melt or boil these substances we are only breaking the intermolecular forces of attraction between molecules. The boiling points increase down this group, which means we have to put in more energy to break the intermolecular forces as the relative molecular mass increases. This means that the *intermolecular forces of attraction must become stronger as relative molecular mass increases*.

This is something that we see quite often in chemistry, for example boiling points increase along the series CH_4, C_2H_6, C_3H_8 as relative molecular mass increases. This is discussed further in Chapter 23.

SOME OTHER PHYSICAL PROPERTIES OF COVALENT COMPOUNDS

KEY POINT

We are using the term covalent molecular compounds to mean covalent compounds with a simple molecular structure.

Covalent molecular compounds do not conduct electricity. This is because the molecules don't have any overall electrical charge (there are no ions) and all the electrons are held tightly in the atoms or in covalent bonds and so are not able to move from molecule to molecule.

Covalent molecular substances tend to be insoluble in water. There are some exceptions to this, for example ethanol (C_2H_5OH) and substances such as NH_3 and HCl that react with water as they dissolve.

Covalent molecular substances are often soluble in organic solvents.

GIANT COVALENT STRUCTURES


DIAMOND

KEY POINT

A tetrahedron is a triangular-based pyramid. In a tetrahedral arrangement, one atom is at the centre of the tetrahedron, and the atoms it is attached to are at the four corners. Look carefully at the top five atoms in Figure 8.26 to see what this looks like. You will find other similar arrangements in this diagram.
In Figure 8.26 some carbon atoms seem to be forming only two bonds (or even one bond), but that's not really the case. We are only showing a small part of the whole structure. The structure continues in three dimensions, and each of the atoms drawn here is attached to four others. Each of the lines in this diagram represents a covalent bond.

Diamond is a form of pure carbon.

Each carbon atom has four electrons in its outer shell and it therefore forms four covalent bonds. In diamond, each carbon bonds strongly to four other carbon atoms in a *tetrahedral* arrangement. Figure 8.26 shows enough of the structure to see what is happening.



▲ Figure 8.26 Part of the structure of diamond

This is a giant covalent structure; it continues on and on in three dimensions. It is *not* a molecule because the number of atoms joined up in a real diamond is completely variable and depends on the size of the crystal. Molecules always contain *fixed numbers* of atoms joined by covalent bonds.

Diamond has a very high melting and boiling point. This is because of the very strong carbon–carbon covalent bonds, which extend throughout the whole crystal in three dimensions. A lot of energy has to be supplied to break these strong covalent bonds, therefore diamond has very high melting and boiling points. It is important to realise how this is different from the simple molecular structures that we saw above. In order to melt or boil a substance with a simple molecular structure, such as CH_4, we only had to supply enough



▲ Figure 8.27 Diamond (a form of carbon) is crystalline, and is the hardest naturally occurring substance.

energy to break the relatively weak intermolecular forces of attraction. In diamond there are no intermolecular forces (it has a giant structure, there are no molecules). Covalent bonds, which are very strong, must be broken in order to melt or boil it.

In general, *all substances with giant covalent structures are solids with high melting and boiling points* because a lot of energy has to be supplied to break all the strong covalent bonds throughout the giant structure. Other substances with giant covalent structures include graphite (discussed below) and silicon dioxide (SiO_2).

Diamond is very hard. Again, a lot of energy has to be supplied to break the strong covalent bonds in the giant structure. Drill bits can be tipped with diamonds for drills used on stone and rock.

Diamond doesn't conduct electricity. All the electrons in the outer shells of the carbon atoms are tightly held in covalent bonds between the atoms. None are free to move around.

EXTENSION WORK

Diamond doesn't dissolve in water or in any other solvent. This is again because of the strong covalent bonds between the carbon atoms. If the diamond dissolved, these bonds would have to be broken.
Diamond conducts heat very well (better than any other element). As one end of the crystal is heated the atoms vibrate more. The strong bonds throughout the giant structure mean that these vibrations are quickly transmitted from one end of the crystal to the other.

GRAPHITE


Graphite is also a form of carbon, but the atoms are arranged differently, although it still has a giant structure. Graphite has a *layer structure*, rather like a pack of playing cards. In a pack of cards, each card is strong but the individual cards are easily separated. The same is true in graphite.



▲ Figure 8.28 Graphite has a structure rather like a pack of playing cards.



▲ Figure 8.29 Graphite is soft with a layer structure.



▲ Figure 8.30 The structure of graphite. Some forces of attraction between layers have been shown. These are not bonds and could have been drawn anywhere between the layers.

EXTENSION WORK

Actually, the reason that the layers slide over each other fairly easily is more complicated than this and graphite is not a lubricant in a vacuum. Graphite being a lubricant relies on water molecules sticking to the surface; this does not happen in a vacuum.

Graphite is a soft material. Although the forces holding the atoms together in each layer are very strong, the attractions between the layers are much weaker and not much energy is needed to overcome them. Layers slide over each other and can easily be flaked off.

Graphite (mixed with clay to make it harder) is used in pencils. When you write with a pencil, you are leaving a trail of graphite layers behind on the paper. Pure graphite is so slippery that it is used as a dry lubricant, for example powdered graphite is used to lubricate locks.

Graphite has high melting and boiling points. To melt/boil graphite, you don't just have to separate the layers, you have to break up the whole structure, including the covalent bonds. That needs very large amounts of energy because the covalent bonds are so strong.



▲ Figure 8.31 Three graphite electrodes glow red hot after their removal from an electric arc furnace used to produce steel.


Graphite conducts electricity. If you look back at Figure 8.30, you will see that each carbon atom is joined to only three others. Each carbon atom uses three of its outer shell electrons to form three single covalent bonds. The fourth electron in the outer shell of each atom is free to move around throughout the whole of the layer. The electrons that are free to move throughout the layers are called **delocalised electrons**. The movement of these delocalised electrons allows graphite to conduct electricity.

EXTENSION WORK

Some other properties of graphite are:

- *Graphite is insoluble in all solvents* because it would take too much energy to break all the strong covalent bonds.
- *Graphite is less dense than diamond* because the layers in graphite are relatively far apart. The distance between the graphite layers is more than twice the distance between atoms in each layer. In a sense, a graphite crystal contains a lot of wasted space, which isn't there in a diamond crystal.

C_{60} FULLERENE




▲ Figure 8.32 A C_{60} fullerene molecule. Remember that molecules consist of fixed number of atoms; each C_{60} fullerene molecule contains 60 carbon atoms joined by covalent bonds.

Diamond and graphite are two **allotropes** of carbon. Allotropes are different forms of the same element. Another allotrope of carbon is C_{60} fullerene. Diamond and graphite both have giant structures but C_{60} fullerene has a simple molecular structure.

In solid or liquid C_{60} fullerene there are C_{60} molecules with weak intermolecular forces between them. The fact that C_{60} has a simple molecular structure has a big influence on its physical properties.

C_{60} fullerene has lower melting and boiling points than diamond and graphite. When fullerene is melted, only the relatively weak intermolecular forces of attraction must be broken. This does not require as much energy as breaking all the strong covalent bonds when diamond and graphite are melted.



▲ Figure 8.33 C_{60} fullerene has a simple molecular structure.

KEY POINT

There are different fullerenes, where the molecules contain different numbers of carbon atoms. This is why we include the C_{60} in the name.

EXTENSION WORK

Unlike diamond and graphite C_{60} fullerene does dissolve in some solvents; only relatively weak intermolecular forces of attraction have to be broken for it to dissolve.

C_{60} fullerene is not as hard as diamond. It does not take as much energy to break the intermolecular forces of attraction in C_{60} fullerene compared to breaking the strong covalent bonds in diamond.

C_{60} fullerene does not conduct electricity. Although all the carbon atoms in C_{60} only form three bonds, the fourth electron on each atom can only move around *within* each C_{60} molecule; the electrons cannot jump from molecule to molecule.

CHAPTER QUESTIONS

You will need to use the Periodic Table in Appendix A on page 320.

SKILLS CRITICAL THINKING 

1 State whether each of the following compounds is ionic or covalent:

a MgO b CH_3Br c H_2O_2

d $FeCl_2$ e NaF f HCN

SKILLS INTERPRETATION 

2 a What is meant by a covalent bond? How does this bond hold two atoms together?

b Draw dot-and-cross diagrams (showing outer shell electrons only) to show the covalent bonding in

i methane (CH_4)

ii hydrogen sulfide (H_2S)

iii phosphine (PH_3)

iv silicon tetrachloride ($SiCl_4$)

c Draw dot-and-cross diagrams (showing outer shell electrons only) to show the covalent bonding in

i O_2

ii N_2

3 Draw dot-and-cross diagrams (showing outer shell electrons only) to show the covalent bonding in

a ethane (C_2H_6) b ethene (C_2H_4) c chloroethane (CH_3CH_2Cl)

SKILLS REASONING

4 Explain why carbon dioxide sublimes at –78.5 °C but diamond sublimes at around 4000 °C.

5 Hexane has the formula C_6H_{14}. It is a liquid at room temperature.

a Explain whether hexane has a simple molecular or giant structure.

b Explain whether you would expect pentane (C_5H_{12}) to have a higher or lower boiling point than hexane.

c Explain whether or not you would expect hexane to conduct electricity.



6 Explain the following in terms of structure and bonding:

a diamond is harder than graphite

b C_{60} fullerene has a lower melting point than graphite

c graphite conducts electricity

d diamond does not conduct electricity.

SKILLS ANALYSIS

8


7 The table below gives details of the boiling temperatures of some substances made of covalent molecules. Arrange these substances in increasing order of the strength of their intermolecular forces and explain your order.

Substance	Boiling point/°C
ammonia	–33
ethanamide	221
ethanol	78.5
hydrogen	–253
phosphorus trifluoride	–101
water	100

(Don't panic if you don't recognise some of the names. The substances could also have been labelled A, B, C, D, E and F.)

SKILLS REASONING

8 The compound N_2F_2 has the structure:



a Explain whether you would expect the bond between the nitrogen atoms to be a single, double or triple bond.

SKILLS INTERPRETATION

b Draw a dot-and-cross diagram for N_2F_2 showing outer electrons only.

SKILLS REASONING

9 a Draw a dot-and-cross diagram (showing outer electrons only) to show the covalent bonding in BCl_3.

9

b BCl_3 is sometimes described as an *electron-deficient* compound. Explain what you think that means.