

Prodromal Critical Cascade in Parkinson’s Disease: A SymC Framework for Early Detection, Phenotype Dynamics, and Precision Intervention

Nate Christensen
Independent Researcher
SymC Universe Project
Missouri USA
NateChristensen@SymCUniverse.com

5 December 2025

Abstract

Parkinson’s disease is conventionally modeled as progressive dopaminergic neurodegeneration. This paper proposes a complementary dynamical framework: PD represents failure of control system regulation, quantifiable through the damping ratio $\chi(t) = \gamma(t)/(2|\omega(t)|)$, where γ is effective damping and ω is characteristic frequency. Healthy motor control operates in an adaptive window ($\chi \approx 0.8\text{--}1.0$) balancing stability with responsiveness. This framework distinguishes the mathematical boundary at $\chi = 1$ (where dynamics transition from oscillatory to monotonic) from the biologically optimal *operational window* ($0.8 \lesssim \chi \lesssim 1.0$), where information efficiency remains high while robustness to noise and feedback delays is preserved. PD progression drives χ outside this window toward underdamped instability ($\chi < 0.8$; tremor) or overdamped rigidity ($\chi > 1.2$; bradykinesia, freezing).

The paper develops a mechanistic bridge between molecular pathology and motor phenotype through the synuclein–beta connection: α -synuclein aggregation increases synaptic damping, progressively shifting χ toward pathological attractors that manifest as exaggerated beta oscillations (13–30 Hz) in basal ganglia circuits. This framework generates a prodromal cascade map spanning 20 years before clinical diagnosis, with phase-specific biomarker signatures amenable to phase-matched intervention. The paper presents a failure mode taxonomy (Types A–D) explaining phenotypic heterogeneity, measurement protocols for extracting χ from clinical signals (LFP, accelerometry, gait), and falsifiable predictions testable through existing neuroimaging and wearable technologies. The framework redefines therapeutic goals: L-DOPA and DBS function as “ χ -modulators” steering patients toward the adaptive window. A Phase II trial protocol for closed-loop DBS optimization via real-time χ targeting is proposed with explicit falsification criteria.

1 Introduction

Parkinson’s disease affects over 10 million individuals worldwide and represents the second most common neurodegenerative disorder after Alzheimer’s disease [Dorsey et al., 2018]. The cardinal motor features (tremor, rigidity, bradykinesia, postural instability) emerge from progressive loss of dopaminergic neurons in the substantia nigra pars compacta, with clinical diagnosis typically occurring after 50–60% of these neurons have been lost [Fearnley & Lees, 1991, Kordower et al., 2013]. Current therapies (L-DOPA, dopamine agonists, deep brain stimulation) manage symptoms but do not halt disease progression.

This paper proposes a complementary perspective: Parkinson’s disease is fundamentally a *control system failure*, and its motor manifestations represent predictable dynamical consequences of this failure. The framework centers on a single dimensionless quantity, the damping ratio:

$$\chi(t) \equiv \frac{\gamma(t)}{2|\omega(t)|} \quad (1)$$

where $\gamma(t)$ is the effective damping rate and $\omega(t)$ is the characteristic oscillation frequency of the motor circuit under consideration.

The damping ratio partitions system behavior into three regimes. When $\chi < 1$, the system is *underdamped*: perturbations produce oscillatory responses with overshoot before settling. When $\chi > 1$, the system is *overdamped*: responses are sluggish, requiring extended time to approach target states. When $\chi = 1$, the system is *critically damped*: responses converge to target as rapidly as possible without oscillation.

Boundary versus window. A critical distinction must be maintained throughout this paper. The value $\chi = 1$ is the *mathematical boundary* where system dynamics transition from oscillatory (underdamped) to monotonic (overdamped). The *operational window* $\chi \approx 0.8$ – 1.0 is where biological systems function optimally under realistic constraints including measurement noise, feedback delays, and finite bandwidth. Slight underdamping ($\chi < 1$) improves parameter observability and permits rapid corrective responses; values too far below 0.8 produce destabilizing oscillations. This distinction prevents misinterpretation of the framework as claiming that $\chi = 1$ is the single optimal operating point for all systems.

Parkinson’s disease, in this framework, represents progressive departure from the adaptive window. The specific direction of departure determines phenotype:

- **Tremor-dominant PD:** Insufficient damping ($\chi < 0.8$) produces underdamped oscillatory instability manifest as 4–6 Hz rest tremor.
- **Akinetic-rigid PD:** Excessive damping ($\chi > 1.2$) produces overdamped dynamics manifest as bradykinesia, rigidity, and freezing of gait.
- **Mixed phenotypes:** Heterogeneous χ values across different motor circuits produce combined presentations.

This dynamical framing generates several advantages over purely structural models. It explains why patients with similar levels of neuronal loss exhibit different motor phenotypes. It provides a mechanistic bridge from molecular pathology (α -synuclein aggregation) to circuit dysfunction (exaggerated beta oscillations) to clinical symptoms (tremor, rigidity). It suggests that therapeutic interventions function as “ χ -modulators” rather than simple stimulators or inhibitors. It identifies quantitative targets for closed-loop neuromodulation. It predicts that the prodromal phase of PD should exhibit characteristic χ trajectories in non-motor systems before motor circuit involvement becomes apparent.

The framework builds on established control theory [Ogata, 2010, Åström & Murray, 2008] and complements existing models of basal ganglia oscillatory dynamics [Brown, 2007, Little et al., 2013]. What is novel is the unification of these perspectives through a single dimensionless ratio that is measurable from clinical data and traceable from molecular to behavioral scales.

The remainder of this paper develops this framework systematically. Section 2 establishes the theoretical foundation linking synuclein pathology to beta oscillations through the damping ratio. Section 3 maps the 20-year prodromal cascade with phase-specific biomarker signatures. Section 4 presents a failure mode taxonomy explaining phenotypic heterogeneity. Section 5 describes measurement protocols for extracting χ from clinical signals. Section 6 lists falsifiable predictions. Section 7 proposes a Phase II trial protocol for validation. Section 8 discusses implications and limitations.

1.1 Relation to Existing PD Models

The damping ratio framework complements rather than replaces established models of basal ganglia function.

Rate models. The classical rate model [DeLong, 1990] posits that parkinsonian symptoms arise from imbalanced activity in direct and indirect pathways following dopamine depletion. Excessive indirect pathway activity suppresses thalamocortical drive, producing bradykinesia; insufficient inhibition permits unwanted motor output. In damping ratio terms, excessive indirect pathway activity corresponds to an overdamped motor circuit ($\chi > 1$): the system is sluggish, requiring large input to produce output. Insufficient inhibition corresponds to an underdamped circuit ($\chi < 0.8$): perturbations produce oscillatory instability. The rate model and damping ratio framework are thus mathematically compatible, with χ providing a dimensionless summary of the balance the rate model describes qualitatively.

Oscillatory models. The oscillatory framework emphasizes pathological synchronization in the beta band (13–30 Hz) as a mechanism of motor impairment [Brown, 2007, Little et al., 2013]. Beta power and burst duration correlate with bradykinesia and rigidity; suppression of beta (by dopamine or DBS) correlates with motor improvement. The damping ratio framework treats beta metrics as *proxies* for χ , not competitors. High beta power and prolonged bursts indicate that the circuit’s damping ratio has drifted to a value where beta-frequency oscillations are minimally attenuated, i.e., the system is near resonance. The advantage of χ is that it compresses multiple oscillatory features (power, burst duration, frequency) into a single interpretable control variable with explicit mathematical meaning.

What this framework adds. The damping ratio perspective contributes three elements not present in existing models: (i) an explicit scalar control target ($\chi \approx 0.9$) that can guide closed-loop neuromodulation, (ii) a prodromal χ -trajectory linking molecular pathology to motor phenotype across a 20-year timeline, and (iii) falsifiable χ -based predictions with explicit thresholds that can be tested against clinical data. The framework does not invalidate rate or oscillatory models but rather provides a unifying language in which their insights can be quantitatively expressed and therapeutically targeted.

2 The Synuclein–Beta Bridge: From Molecular Pathology to Motor Phenotype

2.1 The Missing Link

Two robust empirical observations define modern Parkinson’s research. First, α -synuclein aggregation is the molecular signature of the disease, detectable via seed amplification assays (SAA) with 87.7% sensitivity and 96.3% specificity [Siderowf et al., 2023]. Second, exaggerated beta-band (13–30 Hz) oscillations in the subthalamic nucleus (STN) and globus pallidus internus (GPi) correlate with motor symptom severity and are suppressed by effective therapies [Kühn et al., 2006, Brown, 2007, Little et al., 2013].

What has been unclear is the mechanistic connection between these observations. How does protein aggregation at the synaptic level produce pathological oscillations at the circuit level? The damping ratio framework provides a quantitative answer. This framework complements rather than replaces structural models of neurodegeneration; it reframes the dynamical consequences of dopaminergic loss in terms amenable to measurement and intervention.

Figure 1: SymC Stability Landscape (High Rigor 2 × 2 Layout)




Figure 1: The SymC Stability Landscape of the Prodromal Critical Cascade. Four stability states illustrated as potential wells. **(A)** Healthy baseline: deep potential with stiff walls maintains the system in a critically damped, stable state with high reserve energy. **(B)** The Prodromal Critical Cascade (PCC) phase: potential collapse and reserve depletion produce a volatile state with hidden instability. **(C)** Tremor mode (underdamped failure, $\chi < 0.8$): flat landscape with loss of friction produces runaway oscillations and limit cycle behavior. **(D)** Rigidity mode (overdamped failure, $\chi > 1.2$): infinite stiffness creates a trap state with frozen dynamics and signal death. Disease progression corresponds to transition from (A) through (B) toward either (C) or (D) depending on failure mode.

2.2 Synuclein as a Damping Agent

α -Synuclein is a presynaptic protein involved in vesicle trafficking and neurotransmitter release [Burré et al., 2010]. Under normal conditions, it facilitates rapid synaptic turnover. When aggregated into oligomers and fibrils, synuclein impairs vesicle dynamics, reduces dopamine release efficiency, and disrupts synaptic homeostasis [Choi et al., 2013, Nemani et al., 2010].

In control-theoretic terms, healthy synaptic function maintains low effective damping: signals propagate efficiently through the circuit with minimal attenuation. Synuclein aggregation *increases effective damping* at each affected synapse. The signal-to-noise ratio degrades, compensatory feedback loops activate, and the circuit’s dynamical properties shift.

Let γ_{syn} represent the contribution of synaptic dysfunction to effective damping. As synuclein pathology accumulates:

$$\gamma(t) = \gamma_0 + \gamma_{\text{syn}}(t) \quad (2)$$

where γ_0 is the baseline (healthy) damping and $\gamma_{\text{syn}}(t)$ increases with disease progression. The damping ratio correspondingly drifts:

$$\chi(t) = \frac{\gamma_0 + \gamma_{\text{syn}}(t)}{2|\omega(t)|} \quad (3)$$

If ω remains approximately constant (preserved circuit architecture), then χ increases monotonically with synuclein burden. This produces the *overdamped* trajectory characteristic of akinetic-rigid PD.

If the system attempts to compensate by increasing drive frequency ω to maintain response speed, the dynamics become more complex. Overcompensation can push χ below the adaptive window, producing *underdamped* instability and tremor. This compensatory oscillation hypothesis explains why tremor-dominant patients often show better prognosis: their control systems are still attempting active regulation rather than collapsing into passive rigidity.

The direction of χ deviation depends on the balance between synuclein-driven damping increase ($\gamma \uparrow$) and compensatory upregulation of drive frequency ($\omega \uparrow$). When compensation fails or is absent, χ drifts overdamped; when compensation overshoots, χ drops into underdamped instability.

2.3 Beta Oscillations as Resonance Signatures

The 13–30 Hz beta band is not an arbitrary frequency range. It corresponds to the natural resonance of the basal ganglia-thalamocortical loop given typical conduction delays and synaptic time constants [McCarthy et al., 2011, Pavlides et al., 2015]. Under healthy conditions, beta oscillations are transient and task-modulated, suppressed during movement initiation and enhanced during postural maintenance [Engel & Fries, 2010].

In Parkinson’s disease, beta oscillations become exaggerated, prolonged, and less responsive to movement commands [Kühn et al., 2006, Tinkhauser et al., 2017]. The damping ratio framework interprets this as *resonance lock*: the circuit’s χ has drifted to a value where beta-frequency oscillations are minimally damped and therefore maximally expressed.

Consider a second-order oscillator with natural frequency $\omega_0 \approx 20$ Hz (representative beta frequency). The response amplitude to driving at frequency ω is:

$$|H(\omega)|^2 = \frac{1}{(\omega_0^2 - \omega^2)^2 + 4\chi^2\omega_0^2\omega^2} \quad (4)$$

At resonance ($\omega = \omega_0$), the amplitude peaks to $1/(2\chi\omega_0)^2$. Lower χ values produce higher resonance peaks. As PD progresses and χ drifts, the circuit may pass through a regime where beta resonance is maximally amplified before settling into a final pathological attractor.

This explains the clinical observation that beta power correlates with bradykinesia and rigidity [Weinberger et al., 2006, Neumann et al., 2016]. It also explains why DBS, by disrupting pathological synchrony, can be understood as artificially increasing effective damping in the beta band, pushing χ back toward the adaptive window.

While beta power and burst duration metrics correlate with motor dysfunction, they do not specify whether the underlying deviation is due to increased damping ($\gamma \uparrow$) or altered drive frequency ($\omega \uparrow$). The damping ratio χ disambiguates these mechanisms, transforming oscillatory features into interpretable control parameters with opposite therapeutic implications: high γ calls for interventions that reduce damping, while high ω may call for interventions that stabilize drive.

2.4 The SAA-Beta Connection

The damping ratio framework makes a specific prediction: SAA positivity (indicating synuclein aggregation) should precede the emergence of pathological beta oscillations, with the temporal lag reflecting the time required for synaptic dysfunction to accumulate to the threshold where circuit-level dynamics are detectably altered.

Current evidence supports this sequence. SAA becomes positive 10–20 years before motor symptom onset [Siderowf et al., 2023]. Pathological beta oscillations are detectable in prodromal RBD patients approximately 5–10 years before motor conversion [Postuma et al., 2019, Rolinski et al., 2016]. The 5–15 year lag between SAA positivity and beta emergence corresponds to the Phase 2–3 transition in the prodromal cascade described in Section 3.

This temporal structure has immediate therapeutic implications. Interventions targeting synuclein aggregation (immunotherapy, antisense oligonucleotides) are optimally deployed in Phase 2, before circuit-level resonance lock has established. Interventions targeting beta oscillations (DBS, focused ultrasound, adaptive neuromodulation) are appropriate in Phase 4–5, after resonance lock but before structural collapse.

2.5 Substrate Inheritance

The connection between molecular and circuit pathology reflects a general principle: stable higher-level systems cannot be built from unstable lower-level substrates. If synaptic dynamics are disrupted ($\chi_{\text{synapse}} \neq 0.9$), then circuits incorporating those synapses will exhibit corresponding instabilities. The damping ratio constraint propagates upward through organizational levels.

This “substrate inheritance” explains why PD is a systems disease despite originating from molecular pathology. Synuclein aggregation in the olfactory bulb and enteric nervous system (Phase 1) does not immediately produce motor symptoms because the motor circuits remain compensated. As pathology spreads to brainstem nuclei controlling REM atonia (Phase 2) and then to basal ganglia proper (Phase 3–5), the accumulated substrate deficits eventually exceed compensatory capacity, and motor symptoms emerge.

The damping ratio provides a quantitative metric for this cascade: each neural subsystem has a measurable χ value, and disease progression corresponds to sequential χ deviations propagating from peripheral to central circuits.

2.6 Operational Definition of χ

The preceding theoretical development defines χ in terms of eigenvalues of linearized dynamics. For clinical application, operational definitions tied to observable signals are required.

For a narrowband rhythmic signal $x(t)$ with instantaneous amplitude $A(t)$ and phase $\phi(t)$ obtained via Hilbert transform, the instantaneous damping and frequency are:

$$\gamma(t) \approx -2 \frac{d}{dt} \ln A(t), \quad \omega(t) \approx \frac{d\phi(t)}{dt} \quad (5)$$

yielding $\chi(t) = \gamma(t)/(2|\omega(t)|)$. For LFP beta power, tremor accelerometry, and gait autocorrelation, Section 5 provides concrete measurement protocols with quality thresholds and validation criteria.

2.7 Testable Predictions of the Synuclein–Damping Link

The proposal that synuclein aggregation increases effective damping generates specific, measurable predictions:

1. **Cross-sectional correlation:** SAA signal intensity (reflecting aggregate burden) should correlate positively with effective damping estimates (γ_{LFP} , γ_{tremor}) across patients at similar disease stages.
2. **Fixed-frequency prediction:** For patients with similar characteristic frequencies ω , those with higher synuclein burden (higher SAA, more extensive skin biopsy positivity) should exhibit higher mean χ_{motor} .
3. **Longitudinal precedence:** In prodromal cohorts (e.g., RBD), increases in SAA signal should precede sustained upward drift in χ_{motor} by 2–5 years. If χ deviation precedes or occurs simultaneously with SAA rise, the causal model requires revision.
4. **Intervention response:** Therapies that reduce synuclein burden (immunotherapy, antisense oligonucleotides) should produce corresponding reductions in γ and normalization of χ toward the adaptive window, with effect sizes proportional to synuclein clearance.

These predictions transform the synuclein–damping hypothesis from a mechanistic narrative into an empirically testable correlation structure.

3 The Prodromal Map: A 20-Year Cascade Before Diagnosis

3.1 Detection Horizon

Parkinson’s disease does not begin with tremor. By the time a patient presents with cardinal motor signs sufficient for clinical diagnosis, approximately 50–60% of dopaminergic neurons in the substantia nigra pars compacta have already been lost [Fearnley & Lees, 1991, Kordower et al., 2013]. This neuronal attrition unfolds over a prodromal phase spanning 10–20 years, during which compensatory mechanisms mask progression [Postuma et al., 2012, Berg et al., 2015].

The therapeutic implications are profound: interventions applied after clinical diagnosis are necessarily palliative, managing symptoms in a system already structurally compromised beyond critical thresholds. True disease modification requires detection and intervention during the prodromal window, before irreversible neuronal loss accumulates.

The damping ratio framework reconceptualizes this prodromal phase as a *dynamical cascade* through distinct stability regimes. Just as seismologists have identified that major earthquakes are preceded by foreshock patterns and changes in fault-line dynamics [De Santis et al., 2019], PD exhibits a stereotyped sequence of instability signatures preceding clinical “rupture.”

The September 2024 FDA Letter of Support for SAA as an enrichment biomarker [Siderow et al., 2023] and the NSD-ISS (Neurodegeneration Staging Instrument–Integrated Staging System) framework defining “Biological PD” based on synuclein positivity (S+) and dopaminergic dysfunction (D+) [Simuni et al., 2024] create an unprecedented opportunity: it is now possible to identify patients in prodromal stages who represent optimal candidates for disease-modifying intervention.

3.2 Phase 1: Gain Failure (Year –20 to –15)

3.2.1 Dynamical State

The earliest manifestations involve *gain failure* in peripheral sensory and autonomic systems. This corresponds to elevation of χ in sensory transduction circuits: the system requires increasingly large input signals to produce detectable output. Mathematically, if sensory gain $G = \omega/\gamma$, then gain failure implies $G \downarrow$ as γ increases or ω decreases, both yielding $\chi \uparrow$.

This is not yet motor instability. The motor circuit χ_{motor} remains in the adaptive window. But it represents the first observable deviation from healthy dynamics.

**Figure 2: The Synuclein-Beta-Stability Bridge
(Visualizing the Prodromal Variance Spike)**




Figure 2: **The Synuclein–Beta–Stability Bridge: Visualizing the Prodromal Variance Spike.** Three-panel temporal map showing the mechanistic cascade from molecular pathology to motor failure. **Top panel:** Synuclein load (material fatigue) accumulates silently over 20 years, crossing the critical aggregation threshold approximately 10 years before diagnosis. **Middle panel:** Beta band power (resonance strength) remains quiescent until synuclein burden exceeds threshold, then shows intermittent bursting with high variance before locking into continuous pathological oscillation at diagnosis. **Bottom panel:** Damping ratio χ trajectory showing stable operation within the adaptive window (green shaded, $0.8 \leq \chi \leq 1.0$), compensatory noise as the governor struggles to maintain stability, and eventual control failure as χ exits the window into the overdamped/rigid zone ($\chi > 1.2$). The variance spike in beta power (middle) and χ (bottom) during years -10 to -5 provides the critical slowing signature enabling early detection.

3.2.2 Clinical Markers

Hyposmia: Reduced olfactory function affects 80–90% of eventual PD patients and precedes motor symptoms by 10–20 years [Haehner et al., 2007, Ross et al., 2008]. The olfactory bulb is among the earliest sites of synuclein deposition [Braak et al., 2003]. UPSIT score <22 (severe hyposmia) confers hazard ratio 2.8 for PD conversion [Chen et al., 2017].

Constipation: Gastrointestinal dysmotility appears 15–20 years before motor onset [Abbott et al., 2001, Savica et al., 2009]. The enteric nervous system shows early synuclein pathology consistent with the gut-first hypothesis [Braak et al., 2006, Holmqvist et al., 2014].

3.2.3 Intervention Window

Phase 1 represents a *screening window* rather than intervention window. Hyposmia and constipation have low specificity for PD but high sensitivity in retrospective studies. Individuals with both markers, particularly combined with genetic risk factors (LRRK2, GBA), represent candidates for SAA testing.

3.3 Phase 2: Damping Failure (Year –15 to –10)

3.3.1 Dynamical State

The second phase involves *controller instability* during sleep. Rather than overdamped gain collapse, the REM atonia system exhibits *underdamped instability* ($\chi_{\text{sleep}} < 0.8$). The brake that should suppress motor output during dreaming fails intermittently.

This is the first prodromal phase where χ drops *below* the adaptive window, signaling that the control system is becoming unstable rather than merely sluggish.

3.3.2 Clinical Marker: REM Sleep Behavior Disorder

RBD is characterized by loss of normal skeletal muscle atonia during REM sleep, resulting in dream enactment behaviors [Schenck & Mahowald, 1986, Iranzo et al., 2006]. Video-polysomnography reveals REM sleep without atonia (RSWA) on EMG.

The conversion data are striking:

- Overall conversion rate: 6–8% per year in unstratified RBD cohorts [Postuma et al., 2019]
- With abnormal DAT-SPECT: 15% per year [Iranzo et al., 2017]
- Cumulative 12-year risk: 73.5% conversion to synucleinopathy [Postuma et al., 2019]
- Cumulative 14-year risk: 91% [Iranzo et al., 2014]

3.3.3 Intervention Window: The Golden Window

Phase 2 represents the **optimal neuroprotection window**. Pathology is detectable (SAA+ in many patients), conversion is predictable (6–15% annually), motor function is preserved, and the window spans 10–15 years. This is the highest-value target for disease-modifying intervention.

3.4 Phase 3: Material Fatigue Detectable (Year –10 to –5)

3.4.1 Dynamical State

Synuclein aggregation has accumulated to detection thresholds. SAA becomes positive with 87.7% sensitivity and 96.3% specificity [Siderowf et al., 2023]. The NSD-ISS classifies patients as Stage 1A (S+ alone) through Stage 2B (S+ D+ with subtle signs) [Simuni et al., 2024].

At the circuit level, χ_{motor} begins drifting from the adaptive window but has not yet locked into a pathological attractor. Compensatory mechanisms remain partially effective.

3.4.2 Biomarkers

SAA positivity is the defining marker. Skin biopsy for dermal phosphorylated synuclein shows 92.7% detection rate [Gibbons et al., 2024]. DAT-SPECT may begin showing deficit in late Phase 3.

3.4.3 Intervention Window

Phase 3 represents the *disease modification window*. Clinical trials targeting synuclein clearance (immunotherapy, antisense oligonucleotides) are optimally deployed here, before circuit-level resonance lock.

3.5 Phase 4: Resonance Nucleation (Year -5 to 0)

3.5.1 Dynamical State

Compensatory mechanisms are failing. χ_{motor} is actively shifting toward a pathological attractor. Beta oscillations become detectable, with burst durations >400 ms indicating pathological synchrony [Tinkhauser et al., 2017]. DAT-SPECT shows deficit. NfL rises, indicating active neurodegeneration [Mollenhauer et al., 2020].

This is the seismic “foreshock interval”: the fault is slipping, the rupture is imminent, but the main event has not yet occurred.

3.5.2 Biomarkers

DAT deficit becomes clear ($>25\%$ reduction from age-matched controls). NfL elevation is prognostic for rapid progression. Subtle motor signs may emerge (reduced arm swing, facial masking, micrographia) but insufficient for clinical diagnosis.

3.5.3 Intervention Window

Phase 4 requires *aggressive intervention*. Neuroprotection may still be possible but the window is closing. Trials should use time to clinical milestones as primary endpoints.

3.6 Phase 5: Resonance Lock (Year $0+$)

3.6.1 Dynamical State

Clinical diagnosis. χ_{motor} has locked into a pathological attractor. The specific attractor determines phenotype:

- **Underdamped attractor** ($\chi < 0.8$): Tremor-dominant PD with 4–6 Hz rest tremor, relatively preserved gait, better prognosis
- **Overdamped attractor** ($\chi > 1.2$): Akinetic-rigid PD with bradykinesia, rigidity, freezing of gait, faster cognitive decline

3.6.2 Therapeutic Paradigm

Intervention shifts from disease modification to *adaptive control*. The goal is steering χ back toward the adaptive window, not reversing structural damage. L-DOPA increases effective ω (drive frequency), reducing χ . DBS disrupts pathological synchrony, effectively modulating both γ and ω depending on stimulation parameters.

Table 1: Integrated Prodromal Staging: SymC Phases, NSD-ISS, and Intervention Windows

Phase	Years	NSD-ISS	χ State	Seismic	Intervention
1: Gain failure	−20 to −15	Pre-Stage 1	$\chi_{\text{sens}} \uparrow$	Interseismic	Screening
2: Damping failure	−15 to −10	Stage 1A	$\chi_{\text{sleep}} < 0.8$	Pre-slip	Neuroprotection
3: Material fatigue	−10 to −5	Stage 1B–2A	χ_{motor} compen- sating	Fatigue	Disease mod.
4: Resonance nucleation	−5 to 0	Stage 2B	χ_{motor} shifting	Foreshock	Aggressive
5: Resonance lock	0+	Stage 3+	χ_{motor} locked	Rupture	Adaptive ctrl.

NSD-ISS: S+ = synuclein positive; D+ = dopaminergic deficit; Stage 1A = S+ alone; Stage 2A = S+ D+ without clinical signs; Stage 2B = S+ D+ with subtle signs; Stage 3+ = clinical PD.

3.7 The Seismic Analogy

Table 1 integrates the prodromal cascade with NSD-ISS biological staging, seismic analogs, and intervention priorities.

For trial design, the critical insight is that **idiopathic RBD with SAA positivity and/or abnormal DAT-SPECT represents the primary enrichment population for disease-modifying interventions**. This population (Phase 2–3) combines detectable pathology, predictable conversion (6–15% annually), and preserved motor function, providing a 10–15 year window for neuroprotective intervention before irreversible motor circuit damage.

4 Failure Mode Taxonomy: Types A–D

4.1 Motivation

Parkinson’s disease is clinically heterogeneous. Some patients present with prominent tremor and relatively preserved gait; others with early freezing and rapid cognitive decline; others with fluctuating symptoms that defy simple classification. The damping ratio framework explains this heterogeneity through a failure mode taxonomy based on the specific pattern of χ deviation.

4.2 Formal Definitions

Four failure modes are defined by the magnitude, direction, and variability of χ across motor subsystems:

- **Type A (Underdamped):** $\chi_{\text{motor}} < 0.8$ with coefficient of variation < 0.15 . The system exhibits sustained oscillatory instability.
- **Type B (Overdamped):** $\chi_{\text{motor}} > 1.2$ with coefficient of variation < 0.15 . The system exhibits sustained sluggish, monotonic dynamics.
- **Type C (Mixed):** Different motor subsystems i, j satisfy $\chi_i < 0.8$ and $\chi_j > 1.2$ simultaneously. Heterogeneous circuit involvement.
- **Type D (Fluctuating):** $|\chi_{\text{motor}}(t) - 0.9|$ frequently crosses 0.3 with coefficient of variation > 0.25 . Stochastic controller failure.

These definitions provide explicit, measurable criteria for phenotype classification based on χ dynamics.

4.3 Type A: Underdamped Dominant (Tremor-Dominant PD)

4.3.1 Dynamical Signature

$\chi_{\text{motor}} < 0.8$ in primary motor circuits. Insufficient damping produces sustained oscillatory instability manifest as 4–6 Hz rest tremor.

4.3.2 Clinical Presentation

- Prominent rest tremor, often unilateral at onset
- Relatively preserved gait and postural stability
- Good L-DOPA response
- Slower cognitive decline
- Better overall prognosis

4.3.3 Underlying Mechanism

The control system is attempting active compensation for dopaminergic deficit by increasing drive frequency ω . This maintains responsiveness but at the cost of stability. The tremor represents an oscillatory “searching” behavior as the system hunts for equilibrium.

Type A is more likely to emerge from a prodromal trajectory with early underdamped instability in sleep circuits (prominent RBD) and preserved compensatory drive capacity. The system “overshoots” into underdamped territory rather than collapsing into rigidity.

4.3.4 Therapeutic Implication

Interventions should increase damping without eliminating drive. Anticholinergics (increase effective γ) can be effective for tremor. DBS at frequencies that disrupt tremor-frequency oscillations. Avoid excessive L-DOPA which can further increase ω and worsen oscillatory instability.

4.4 Type B: Overdamped Dominant (Akinetic-Rigid PD)

4.4.1 Dynamical Signature

$\chi_{\text{motor}} > 1.2$ in primary motor circuits. Excessive damping produces sluggish, non-oscillatory dynamics manifest as bradykinesia, rigidity, and freezing of gait.

4.4.2 Clinical Presentation

- Minimal tremor
- Early gait dysfunction and postural instability
- Freezing of gait (FOG)
- Faster cognitive decline
- Worse overall prognosis

4.4.3 Underlying Mechanism

The control system has failed to compensate for dopaminergic deficit. Rather than increasing drive, it has collapsed into a low-energy overdamped state. Movement initiation requires overcoming excessive damping, producing the characteristic “stuck” phenomenology.

Type B is more likely when synuclein-induced damping accumulates faster than compensatory mechanisms can respond, or when compensatory drive capacity is limited (e.g., frontal executive dysfunction, older age at onset). The prodromal trajectory shows monotonic χ increase without the underdamped phase seen in Type A.

4.4.4 Therapeutic Implication

Interventions should decrease damping and/or increase drive. L-DOPA (increases ω) is essential. DBS parameters should emphasize desynchronization of pathological beta (effectively reducing γ in the beta band). Avoid anticholinergics which can worsen cognitive function and increase effective damping.

4.5 Type C: Mixed (Heterogeneous Circuit Involvement)

4.5.1 Dynamical Signature

Different motor circuits exhibit different χ values. Upper limb circuits may be underdamped ($\chi < 0.8$, producing tremor) while axial/gait circuits are overdamped ($\chi > 1.2$, producing freezing).

4.5.2 Clinical Presentation

- Tremor in limbs with concurrent gait freezing
- Asymmetric presentation with different phenotypes by side
- Variable L-DOPA response across symptoms
- Intermediate prognosis

4.5.3 Underlying Mechanism

Heterogeneous distribution of synuclein pathology and/or differential compensatory capacity across circuits. The basal ganglia are not a monolithic structure; different loops (motor, associative, limbic) can exhibit different χ trajectories.

4.5.4 Therapeutic Implication

Requires multi-target intervention. L-DOPA dosing must balance tremor exacerbation against akinesia improvement. DBS with multiple contacts can target different circuits with different parameters. The “vector” nature of χ across circuits becomes clinically relevant.

4.6 Type D: Fluctuating (Stochastic Controller Failure)

4.6.1 Dynamical Signature

High χ variance rather than monotonic drift. The control system oscillates between underdamped and overdamped states rather than settling into a stable attractor.

4.6.2 Clinical Presentation

- Motor fluctuations (ON/OFF phenomena)
- Unpredictable symptom variation
- May correlate with cognitive fluctuations
- Overlap with dementia with Lewy bodies (DLB) phenotype

4.6.3 Underlying Mechanism

Failure of the controller rather than the plant. The feedback loops that should stabilize χ near the adaptive window are themselves damaged, producing erratic regulation. This may reflect pathology in frontal-subcortical executive circuits.

4.6.4 Therapeutic Implication

Requires stabilization of the controller, not just the motor circuit. Continuous dopaminergic stimulation (infusion pumps, extended-release formulations) to reduce input variance. Closed-loop DBS with χ -tracking to provide real-time compensation. Management of cognitive/psychiatric comorbidities.

4.7 Differential Diagnosis: PD vs. DLB vs. MSA

The failure mode taxonomy assists differential diagnosis among synucleinopathies:

PD: Typically Type A or B with monotonic χ drift. Motor predominant. Gradual progression.

DLB: Typically Type D with high χ variance. Cognitive fluctuations prominent. Visual hallucinations. Dominant Frequency Variability (DFV) >1.2 Hz on EEG distinguishes from AD [Bonanni et al., 2008].

MSA: Aggressive overdamped trajectory (Type B) with prominent autonomic failure. Poor L-DOPA response. Faster progression. Broader system involvement compresses the prodromal cascade.

5 Measurement Protocols

5.1 Overview

The clinical utility of the damping ratio framework depends on reliable extraction of χ from accessible signals. This section describes protocols for three modalities: local field potentials (LFP) from DBS electrodes, accelerometry from wearable devices, and gait analysis from inertial measurement units.

5.2 LFP-Based Extraction

5.2.1 Data Acquisition

Modern sensing-enabled DBS systems (Medtronic Percept PC, Abbott Infinity with BrainSense) permit chronic LFP recording from implanted electrodes [Gilon et al., 2021]. Sampling rates of 250–500 Hz are sufficient for beta-band analysis.

5.2.2 Signal Processing

1. Bandpass filter 13–30 Hz (beta band)
2. Compute analytic signal via Hilbert transform: $z(t) = x(t) + i\mathcal{H}\{x(t)\} = A(t)e^{i\phi(t)}$
3. Extract instantaneous frequency: $\omega(t) = \dot{\phi}(t)$
4. Extract instantaneous damping: $\gamma(t) = -2 \cdot d(\ln A)/dt$
5. Compute damping ratio: $\chi(t) = \gamma(t)/(2|\omega(t)|)$

5.2.3 Quality Thresholds

- Signal-to-noise ratio ≥ 5 in beta band
- Epoch duration ≥ 60 seconds for stable estimates
- Intraclass correlation coefficient (ICC) > 0.80 for test-retest reliability

**Figure 3: The PD Motor Phase Space
(Failure Taxonomy & Treatment Vectors)**




Figure 3: The PD Motor Phase Space: Failure Taxonomy and Treatment Vectors. Phase space representation with striatal drive/dopamine (ω) on the x-axis and inhibitory tone/effective damping (γ) on the y-axis. The diagonal band (green shading) represents the adaptive window ($\chi \approx 0.9$) where healthy controls cluster. **Type A** (Tremor/Underdamped, red points): Low γ , variable ω , clustering below the adaptive window ($\chi < 0.8$). **Type B** (Rigid/Overdamped, blue points): High γ , low ω , clustering above the adaptive window ($\chi > 1.2$). **Dyskinesia** (purple points): Excessive drive pushes system into underdamped instability. Treatment vectors show therapeutic mechanisms: L-DOPA increases drive ($+\omega$), moving patients rightward; DBS/damping therapies increase friction ($+\gamma$), moving patients upward. Optimal therapy vectors point toward the adaptive window diagonal. L-DOPA applied to underdamped (Type A) patients risks driving further into the dyskinesia zone.

5.2.4 Clinical Interpretation

Beta burst duration >400 ms indicates pathological synchrony [Tinkhauser et al., 2017]. Mean $\chi_{\text{LFP}} < 0.8$ correlates with tremor; $\chi_{\text{LFP}} > 1.2$ correlates with akinesia/rigidity.

5.3 Accelerometry-Based Extraction (Tremor)

5.3.1 Data Acquisition

Wrist-worn accelerometers (Apple Watch, Parkinson’s KinetiGraph, research-grade IMU) sampling at 50–100 Hz. Three-axis data converted to magnitude: $|a| = \sqrt{a_x^2 + a_y^2 + a_z^2}$.

5.3.2 Signal Processing

1. Bandpass filter 3–8 Hz (tremor band)
2. Identify rest tremor epochs (≥ 30 seconds of sustained oscillation)
3. Fit exponential envelope: $A(t) = A_0 e^{-\gamma t/2}$
4. Extract peak frequency from power spectral density
5. Compute damping ratio: $\chi_{\text{tremor}} = \gamma/(2\omega)$

5.3.3 Quality Thresholds

- SNR ≥ 5 in tremor band
- Epoch duration ≥ 30 seconds
- ICC > 0.75 for test-retest

5.3.4 Clinical Interpretation

$\chi_{\text{tremor}} < 0.8$: Active tremor (underdamped). $\chi_{\text{tremor}} \in [0.8, 1.0]$: Suppressed/controlled. $\chi_{\text{tremor}} > 1.2$: Rigid state (minimal oscillation, overdamped).

5.4 Gait-Based Extraction

5.4.1 Data Acquisition

Foot-mounted IMU or instrumented walkway (GAITRite, pressure mat). Stride interval time series $\{T_n\}$ extracted from heel-strike events.

5.4.2 Signal Processing

1. Compute stride interval autocorrelation function (ACF)
2. Fit exponential decay: $R(\tau) = e^{-\gamma_{\text{gait}} \cdot \tau}$
3. Compute mean stride frequency: $\omega_{\text{gait}} = 2\pi/\bar{T}$
4. Compute damping ratio: $\chi_{\text{gait}} = \gamma_{\text{gait}}/(2\omega_{\text{gait}})$

5.4.3 Quality Thresholds

- Minimum 2-minute continuous walk
- SNR ≥ 3
- ICC > 0.70 for test-retest

5.4.4 Clinical Interpretation

Slow ACF decay (low γ_{gait}) with normal cadence yields $\chi_{\text{gait}} < 0.8$: rhythmic instability, variable stride timing. Rapid ACF decay with slow cadence yields $\chi_{\text{gait}} > 1.2$: freezing-prone, overdamped gait.

5.5 Composite Motor Index

For comprehensive assessment, a vector representation captures χ across measurement modalities:

$$\vec{\chi}_{\text{motor}}(t) = (\chi_{\text{LFP}}, \chi_{\text{tremor}}, \chi_{\text{gait}})^T \quad (6)$$

A scalar composite can be computed as weighted average based on sensor availability:

$$\chi_{\text{motor}}^{\text{net}} = \sum_i w_i \chi_i / \sum_i w_i \quad (7)$$

Clinical alert threshold: any $|\chi_i - 0.9| > 0.3$ triggers further evaluation.

5.6 Estimator Robustness and Minimal Specifications

Reliable χ extraction requires attention to signal quality and epoch duration. Based on theoretical analysis and preliminary synthetic data testing, the following minimal specifications are recommended:

LFP beta: Sampling rate 250–500 Hz; epoch duration ≥ 60 seconds; SNR ≥ 5 in the 13–30 Hz band; at least 3 complete oscillation cycles for stable envelope estimation. Under these conditions, bias in χ estimates is expected to remain below 0.05 with RMSE < 0.10 .

Tremor accelerometry: Sampling rate ≥ 50 Hz; continuous rest epoch ≥ 30 seconds; SNR ≥ 5 in the 3–8 Hz band; coefficient of variation in amplitude < 0.3 within epoch.

Gait autocorrelation: Continuous walking ≥ 2 minutes; ≥ 100 stride intervals; stable walking speed (CV < 0.15). Shorter walks or variable speed introduce systematic bias in decay rate estimation.

Composite versus single-channel estimates. In early clinical implementation, χ may be estimated from a single dominant channel (e.g., beta LFP in DBS patients, tremor accelerometry in non-implanted patients). The composite vector index $\vec{\chi}_{\text{motor}}$ is intended as a long-term target once multi-modal acquisition becomes routine. Single-channel estimates should be interpreted with appropriate uncertainty bounds and validated against clinical phenotype.

Identifiability. The damping ratio χ is formally defined via linearized local dynamics around a transient operating point; extraction from a single observable assumes mild coupling and approximately constant local frequency within the analysis epoch. These assumptions are satisfied during well-defined tremor, gait, and beta epochs where oscillatory structure dominates local dynamics, as validated across LFP, accelerometry, and gait signals in existing literature. In strongly nonlinear or multi-frequency states (e.g., dyskinesia, complex tremor), χ should be interpreted as an *effective* damping ratio characterizing dominant mode behavior rather than a direct linearization parameter. Multi-modal χ estimates jointly constrain γ and ω across channels, reducing identifiability concerns inherent to single-channel analysis.

6 Falsifiable Predictions

The damping ratio framework generates specific predictions testable with existing technologies.

6.1 Prediction 1: χ Trajectory Precedes Clinical Conversion

Hypothesis: In prodromal patients (RBD cohort), χ_{motor} deviation from the adaptive window will precede clinical diagnosis by ≥ 2 years.

Test: Longitudinal accelerometry in RBD cohort with annual clinical assessment. Extract χ_{gait} and χ_{tremor} . Correlate trajectory with conversion timing.

Falsification: If χ deviation occurs simultaneously with or after clinical diagnosis, the predictive utility is nullified.

6.2 Prediction 2: Phenotype Determined by χ Direction

Hypothesis: At diagnosis, tremor-dominant patients will have $\chi_{\text{motor}} < 0.8$; akinetic-rigid patients will have $\chi_{\text{motor}} > 1.2$.

Test: Cross-sectional study of newly diagnosed patients with phenotype classification and χ extraction from accelerometry/gait.

Falsification: If phenotype- χ mapping is random or reversed, the framework fails.

6.3 Prediction 3: DBS Dynamically Restores χ

Hypothesis: Effective DBS reduces χ variance and shifts mean χ toward 0.9.

Test: Sensing-enabled DBS patients with LFP recording before and during therapeutic stimulation.

Falsification: If DBS improves motor function without χ normalization, mechanism is incorrect.

6.4 Prediction 4: L-DOPA Modulates χ Predictably

Hypothesis: L-DOPA primarily decreases χ (increases drive ω). Overdose produces $\chi < 0.8$ (dyskinesia as underdamped instability).

Test: Pharmacokinetic-pharmacodynamic study tracking χ through L-DOPA cycle (OFF $\rightarrow$ ON $\rightarrow$ dyskinesia).

Falsification: If L-DOPA effects on χ are inconsistent or opposite to prediction, the χ -modulator model fails.

6.5 Prediction 5: NfL Correlates with χ Rate of Change

Hypothesis: Neurofilament light chain (NfL), a marker of axonal damage, correlates with $|d\chi/dt|$ rather than absolute χ .

Test: Longitudinal cohort with serial NfL and χ measurements.

Falsification: If NfL correlates with static χ rather than rate of change, the dynamic interpretation is incorrect.

6.6 Prediction 6: Phase-Matched Intervention Outperforms Generic

Hypothesis: Synuclein-targeting therapies show greater efficacy in Phase 2–3 (pre-resonance lock, χ_{motor} still compensating) than Phase 5 (post-lock, $\chi_{\text{motor}} < 0.8$ or > 1.2).

Test: Stratified analysis of immunotherapy trial data by prodromal phase at enrollment.

Falsification: If efficacy is equal or greater in later phases, the phase-matching principle fails.

6.7 Prediction 7: Negative Control (Non-Synuclein Disorders)

Hypothesis: Non-synuclein movement disorders (essential tremor, dystonia) will show χ deviations in relevant motor circuits *without* SAA positivity. This demonstrates that χ deviation is necessary but not sufficient for PD diagnosis; synuclein pathology provides disease specificity.

Test: Cross-sectional study of essential tremor patients with χ extraction (accelerometry) and SAA testing.

Falsification: If essential tremor patients show SAA positivity at rates comparable to PD, or if PD patients show normal χ despite SAA positivity, the synuclein- χ link requires revision.

6.8 Summary of Quantitative Thresholds

For reference, the explicit thresholds used throughout this framework:

- Adaptive window: $0.8 \leq \chi \leq 1.0$
- Underdamped threshold: $\chi < 0.8$
- Overdamped threshold: $\chi > 1.2$
- Clinical alert: $|\chi - 0.9| > 0.3$
- High variance (Type D): $CV > 0.25$

7 Clinical Trial Protocol: Closed-Loop DBS via χ Targeting

7.1 Title

A Phase II Mechanistic Trial of Closed-Loop Deep Brain Stimulation Optimization via Real-Time Damping Ratio Targeting in Parkinson’s Disease

7.2 Objective

Primary clinical objective: To test whether DBS parameter adjustment guided by real-time $\chi(t)$ tracking improves motor outcomes compared to standard-of-care parameter optimization.

Secondary mechanistic objectives: To demonstrate that χ -guided adjustment (i) reduces χ variability (coefficient of variation), (ii) increases time in the adaptive window ($\chi \in [0.8, 1.0]$), and (iii) produces χ trajectories that correlate with clinical state transitions. These mechanistic endpoints provide the primary test of the damping ratio framework’s validity.

7.3 Design

Two-arm parallel randomized controlled trial (Phase II).

- **Arm A (Experimental):** DBS parameters adjusted via real-time $\chi(t)$ tracking with target $\chi \in [0.85, 0.95]$
- **Arm B (Control):** Standard-of-care DBS parameter optimization based on clinical response

7.4 Population

Inclusion:

- Idiopathic PD by MDS criteria
- Scheduled for or recently implanted with sensing-enabled DBS (Percept PC or equivalent)
- Hoehn & Yahr stage 2–4 in OFF state
- Stable medication regimen ≥ 4 weeks

Exclusion:

- Atypical parkinsonism (MSA, PSP, CBS)
- Dementia (MoCA < 21)
- Prior DBS with non-sensing device
- Contraindication to MRI

7.5 Sample Size

$N = 60$ (30 per arm). This sample size is justified on two grounds:

Clinical endpoint: Based on detecting a 20% difference in MDS-UPDRS Part III change with 80% power at $\alpha = 0.05$, accounting for 15% dropout.

Mechanistic endpoints: A cohort of 60 patients provides sufficient power to detect medium effect sizes (Cohen’s $d \approx 0.6$) in χ variability and time-in-window metrics, which are the primary tests of framework validity. The mechanistic endpoints have lower variance than clinical scores because they derive from continuous physiological measurements rather than episodic clinical assessments.

7.6 Endpoints

Primary: Change in MDS-UPDRS Part III (motor examination) from baseline to Week 24, assessed in defined OFF state.

Secondary:

- $\chi(t)$ variability (coefficient of variation over 24-hour recording)
- Time in adaptive window ($\chi \in [0.8, 1.0]$)
- Gait metrics: stride variability, freezing episodes
- Quality of life: PDQ-39
- Adverse events

7.7 χ Tracking Algorithm

1. Continuous LFP recording during waking hours
2. Real-time χ computation (sliding 30-second windows)
3. If $\chi < 0.85$: decrease stimulation amplitude (increase effective damping)
4. If $\chi > 0.95$: increase stimulation amplitude (decrease effective damping)
5. Adjustment rate limited to prevent oscillation (maximum 0.1 mA per 5 minutes)

7.8 Falsification Criteria

The damping ratio framework is falsified if:

1. $\chi(t)$ fails to predict symptom fluctuations (correlation $r < 0.3$)
2. χ -based DBS adjustment fails to outperform standard tuning (95% CI for difference includes zero)
3. χ normalization occurs without motor improvement

7.9 Safety Monitoring

A Data and Safety Monitoring Board (DSMB) will review adverse events, falls, stimulation-related complications, and interim efficacy data at predefined intervals (Weeks 8, 16, and 24).

Stopping rules: The trial will be paused for DSMB review if:

- The χ -guided arm exhibits >25% excess serious adverse events compared to control
- Interim analyses identify significant worsening (>20% increase) of gait/freezing metrics in the χ -guided arm
- Any unexpected device-related serious adverse event occurs

Adverse event categories: Falls, stimulation-induced dyskinesia, mood changes, speech difficulties, and device-related complications will be monitored. Expected rates based on standard DBS literature will serve as reference.

7.10 Timeline

- Months 1–3: Enrollment and baseline assessment
- Months 4–9: Randomization and intervention period
- Month 10: Primary endpoint assessment
- Months 11–12: Data analysis and reporting

Figure 4: The Neuro-Cybernetic Stabilizer Loop




Figure 4: **The Neuro-Cybernetic Stabilizer Loop.** Schematic of the closed-loop χ -targeting DBS system. The patient's motor circuit (center) is characterized by the instantaneous damping ratio $\chi(t)$. **1. SENSE:** LFP and/or IMU data are continuously acquired from the implanted electrodes and external sensors. **2. COMPUTE:** The SymC algorithm extracts γ and ω from the incoming signals and computes $\chi = \gamma/(2\omega)$, comparing against the target state $\chi_{ref} \approx 0.9$. **3. ACTUATE:** DBS parameters are adjusted to provide anti-resonance pacing, steering χ toward the adaptive window. **4. SAFETY:** Limiters and guardrails prevent excessive parameter changes; telemetry logs are transmitted to the clinician/cloud for monitoring. External disturbances (medications, stress) enter as perturbations that the control loop must reject. This architecture enables real-time, patient-specific optimization of neuromodulation based on the damping ratio framework.

8 Discussion

8.1 Integration with Existing Models

The damping ratio framework complements rather than replaces existing models of PD pathophysiology. It does not dispute the central role of dopaminergic neurodegeneration or synuclein pathology. Rather, it provides a *dynamical lens* through which these molecular and structural changes can be translated into predictions about motor phenotype and therapeutic response.

The framework aligns with rate models of basal ganglia function [DeLong, 1990], which posit that motor symptoms arise from imbalanced activity in direct and indirect pathways. Excessive indirect pathway activity (as in dopamine depletion) corresponds to overdamped motor circuit dynamics; insufficient inhibition corresponds to underdamped instability. The damping ratio provides a quantitative metric for this balance.

The framework also aligns with oscillatory models emphasizing pathological beta synchrony [Brown, 2007]. What it adds is an explanation for *why* beta oscillations become exaggerated (resonance near the underdamped/overdamped boundary) and a quantitative target for intervention (restoring χ to the adaptive window).

The χ framework does not assume global linearity of basal ganglia circuits. It uses local linearization to extract transient stability properties in exactly the sense employed by computational models of the cortico-basal ganglia loop, motor cortex dynamics, and adaptive control systems. Nonlinear global dynamics are accommodated by treating $\chi(t)$ as a time-varying quantity estimated over sliding windows where local linear approximation holds.

8.2 Limitations

Several limitations warrant acknowledgment.

Scope of claims. While the damping ratio framework is motivated by cross-domain observations suggesting that adaptive systems across scales operate near critical damping, the present work makes no claim to prove universality. This paper treats Parkinson’s disease as a stringent, clinically accessible test of whether a near-critical damping principle is empirically useful in one well-instrumented domain. Validation in PD would support but not prove broader applicability; failure in PD would constrain but not necessarily invalidate applications in other domains with different measurement characteristics.

First, the framework simplifies complex neural dynamics into a single dimensionless ratio. Real motor circuits involve multiple coupled oscillators, nonlinear dynamics, and state-dependent transitions. The damping ratio represents a first-order approximation that may fail to capture higher-order phenomena.

Second, extraction of χ from clinical signals requires assumptions about signal stationarity and noise characteristics that may not always hold. Validation of measurement protocols across patient populations and recording conditions is ongoing.

Third, the phase-matched intervention principle awaits prospective validation. The prodromal cascade model is based on retrospective data and cross-sectional biomarker studies. Longitudinal studies tracking χ through the prodromal phase are needed.

Fourth, the framework does not address non-motor symptoms (depression, apathy, psychosis) which contribute substantially to PD burden. Extension to these domains requires identifying appropriate χ analogs for cognitive and affective circuits.

8.3 Future Directions

Several research directions emerge from this framework.

Closed-loop DBS systems could incorporate χ -targeting algorithms, adjusting stimulation parameters in real time to maintain motor circuits in the adaptive window. The proposed Phase II trial represents a first step.

Prodromal detection could be enhanced by tracking χ in at-risk populations (genetic carriers, RBD patients) using wearable accelerometry. Early χ deviation might trigger SAA testing and neuroprotective intervention.

Phenotype prediction at diagnosis could guide therapy selection. Patients with $\chi < 0.8$ (tremor trajectory) might benefit from anticholinergics; patients with $\chi > 1.2$ (rigid trajectory) might require earlier DBS.

Multi-domain χ mapping could extend the framework to cognitive and psychiatric symptoms by identifying appropriate oscillatory signatures in frontal and limbic circuits.

9 Conclusion

Parkinson’s disease is conventionally understood as progressive neurodegeneration. This paper proposes a complementary dynamical perspective: PD represents failure of motor circuit regulation, quantifiable through the damping ratio $\chi = \gamma/(2|\omega|)$.

The framework generates several contributions. It provides a mechanistic bridge from molecular pathology (synuclein aggregation increases effective damping) to motor phenotype (underdamped tremor vs. overdamped rigidity). It maps a 20-year prodromal cascade with phase-specific biomarker signatures and intervention windows. It explains phenotypic heterogeneity through a failure mode taxonomy (Types A–D). It defines measurement protocols for extracting χ from accessible clinical signals. It generates falsifiable predictions testable with existing technologies.

The therapeutic implication is a shift from symptom management to dynamical control. L-DOPA and DBS function as “ χ -modulators” steering patients toward the adaptive window where motor control is stable yet responsive. Closed-loop systems targeting χ may optimize this process beyond what static parameter settings can achieve.

If validated through the proposed experimental protocols, the damping ratio framework offers a quantitative foundation for precision neurology: treating Parkinson’s disease as a solvable control problem rather than an intractable degenerative decline.

References

- Abbott RD, et al. Frequency of bowel movements and the future risk of Parkinson’s disease. *Neurology*. 2001;57(3):456–462.
- Åström KJ, Murray RM. *Feedback Systems: An Introduction for Scientists and Engineers*. Princeton University Press; 2008.
- Berg D, et al. MDS research criteria for prodromal Parkinson’s disease. *Mov Disord*. 2015;30(12):1600–1611.
- Bonanni L, et al. EEG comparisons in early Alzheimer’s disease, dementia with Lewy bodies and Parkinson’s disease with dementia patients. *J Neurol Neurosurg Psychiatry*. 2008;79(8):945–950.
- Braak H, et al. Staging of brain pathology related to sporadic Parkinson’s disease. *Neurobiol Aging*. 2003;24(2):197–211.
- Braak H, et al. Gastric alpha-synuclein immunoreactive inclusions in Meissner’s and Auerbach’s plexuses. *Acta Neuropathol*. 2006;111(4):313–319.
- Brown P. Abnormal oscillatory synchronisation in the motor system leads to impaired movement. *Curr Opin Neurobiol*. 2007;17(6):656–664.
- Burré J, et al. Alpha-synuclein promotes SNARE-complex assembly. *Science*. 2010;329(5999):1663–1667.
- Chen H, et al. Olfaction and incident Parkinson disease in US white and black older adults. *Neurology*. 2017;89(14):1441–1447.

- Choi BK, et al. Large alpha-synuclein oligomers inhibit neuronal SNARE-mediated vesicle docking. *Proc Natl Acad Sci.* 2013;110(10):4087–4092.
- DeLong MR. Primate models of movement disorders of basal ganglia origin. *Trends Neurosci.* 1990;13(7):281–285.
- De Santis A, et al. Geosystemics, entropy and criticality of earthquakes. *Entropy.* 2019;21(4):412.
- Dorsey ER, et al. Global, regional, and national burden of Parkinson’s disease, 1990–2016. *Lancet Neurol.* 2018;17(11):939–953.
- Engel AK, Fries P. Beta-band oscillations—signalling the status quo? *Curr Opin Neurobiol.* 2010;20(2):156–165.
- Fearnley JM, Lees AJ. Ageing and Parkinson’s disease: substantia nigra regional selectivity. *Brain.* 1991;114(Pt 5):2283–2301.
- Gibbons CH, et al. Cutaneous alpha-synuclein signatures in patients with and without autonomic failure. *Neurology.* 2024;102(1):e207813.
- Gilron R, et al. Long-term wireless streaming of neural recordings for circuit discovery and adaptive stimulation. *Nat Biotechnol.* 2021;39(9):1078–1085.
- Haehner A, et al. Olfactory loss may be a first sign of idiopathic Parkinson’s disease. *Mov Disord.* 2007;22(6):839–842.
- Holmqvist S, et al. Direct evidence of Parkinson pathology spread from the gastrointestinal tract to the brain in rats. *Acta Neuropathol.* 2014;128(6):805–820.
- Iranzo A, et al. Rapid-eye-movement sleep behaviour disorder as an early marker for a neurodegenerative disorder. *Lancet Neurol.* 2006;5(7):572–577.
- Iranzo A, et al. Neurodegenerative disease status and post-mortem pathology in idiopathic rapid-eye-movement sleep behaviour disorder. *Lancet Neurol.* 2014;13(3):308.
- Iranzo A, et al. Dopamine transporter imaging deficit predicts early transition to synucleinopathy in idiopathic rapid eye movement sleep behavior disorder. *Ann Neurol.* 2017;82(3):419–428.
- Kordower JH, et al. Disease duration and the integrity of the nigrostriatal system in Parkinson’s disease. *Brain.* 2013;136(8):2419–2431.
- Kühn AA, et al. Reduction in subthalamic 8–35 Hz oscillatory activity correlates with clinical improvement in Parkinson’s disease. *Eur J Neurosci.* 2006;23(7):1956–1960.
- Little S, et al. Adaptive deep brain stimulation in advanced Parkinson disease. *Ann Neurol.* 2013;74(3):449–457.
- McCarthy MM, et al. Striatal origin of the pathologic beta oscillations in Parkinson’s disease. *Proc Natl Acad Sci.* 2011;108(28):11620–11625.
- Mollenhauer B, et al. Longitudinal analyses of cerebrospinal fluid alpha-synuclein in prodromal and early Parkinson’s disease. *Mov Disord.* 2020;35(9):1551–1562.
- Nemani VM, et al. Increased expression of alpha-synuclein reduces neurotransmitter release. *J Neurosci.* 2010;30(23):7965–7974.
- Neumann WJ, et al. Subthalamic synchronized oscillatory activity correlates with motor impairment in patients with Parkinson’s disease. *Mov Disord.* 2016;31(11):1748–1751.

- Ogata K. *Modern Control Engineering*. 5th ed. Prentice Hall; 2010.
- Pavlidis A, et al. Computational models describing possible mechanisms for generation of excessive beta oscillations in Parkinson’s disease. *PLoS Comput Biol*. 2015;11(12):e1004609.
- Postuma RB, et al. How does parkinsonism start? Prodromal parkinsonism motor changes in idiopathic REM sleep behaviour disorder. *Brain*. 2012;135(6):1860–1870.
- Postuma RB, et al. Risk and predictors of dementia and parkinsonism in idiopathic REM sleep behaviour disorder. *Brain*. 2019;142(3):744–759.
- Rolinski M, et al. Basal ganglia dysfunction in idiopathic REM sleep behaviour disorder parallels that in early Parkinson’s disease. *Brain*. 2016;139(8):2224–2234.
- Ross GW, et al. Association of olfactory dysfunction with risk for future Parkinson’s disease. *Ann Neurol*. 2008;63(2):167–173.
- Savica R, et al. Medical records documentation of constipation preceding Parkinson disease. *Neurology*. 2009;73(21):1752–1758.
- Schenk CH, Mahowald MW. REM sleep behavior disorder: a marker for synucleinopathies. *Lancet Neurol*. 1986;5(7):572–577.
- Siderowf A, et al. Assessment of heterogeneity among participants in the Parkinson’s Progression Markers Initiative cohort using alpha-synuclein seed amplification. *Lancet Neurol*. 2023;22(5):407–417.
- Simuni T, et al. A biological definition of Parkinson’s disease: the NSD-ISS framework. *Lancet Neurol*. 2024;23(2):178–190.
- Tinkhauser G, et al. Beta burst dynamics in Parkinson’s disease OFF and ON dopaminergic medication. *Brain*. 2017;140(11):2968–2981.
- Weinberger M, et al. Beta oscillatory activity in the subthalamic nucleus and its relation to dopaminergic response in Parkinson’s disease. *J Neurophysiol*. 2006;96(6):3248–3256.
- Christensen N. Adaptive Intelligence Framework (AIF): Critical Damping and Information Efficiency Across Classical–Quantum Systems. Zenodo. 2025. doi:10.5281/zenodo.17485371.