
Welcome! What medication are you looking for?

I'm a _Any_ year old Female v, and I'm insured by the ___Any__ insurance company.

The medication I want to search is _____

Search




Check side effects before and after you take your medications

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aenean euismod bibendum laoreet. Proin gravida dolor sit amet lacus accumsan et viverra justo commodo.

Check if your medication is covered in your insurance plan


Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aenean euismod bibendum laoreet. Proin gravida dolor sit amet lacus accumsan et viverra justo commodo.





See how other people experienced your medications

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aenean euismod bibendum laoreet. Proin gravida dolor sit amet lacus accumsan et viverra justo commodo.



Take it with a meal or a snack to prevent upset stomach.

Can make you bruise or bleed more easily and it can also cause stomach ulcers.

Don't take it with other blood thinners, like aspirin, unless your doctor tells you it's safe.

Can hurt your kidneys, so don't use it if you have a history of kidney problems.

Suggested Usage

Directions use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. use in a well ventilated area maximal treatment area for one applicator is approximately 13.2 in. × 13.2 in. (1126 cm 2). remove applicator from package; do not touch sponge hold the applicator with the sponge down. Pinch wing only once to activate the ampules and release the antiseptic. wet the sponge by pressing and releasing the sponge against the treatment ... read more

Interactivity with other medications or food

Do not take this medication with: Celexa Cymbalta Effexor Lexapro Prozac

