


[SIGN UP](#)[LOG IN](#)

Provide professional information and help of your medications





Check side effects before and after you take your medications

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aenean euismod bibendum laoreet. Proin gravida dolor sit amet lacus accumsan et viverra justo commodo.

See how other people experienced your medications

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aenean euismod bibendum laoreet. Proin gravida dolor sit amet lacus accumsan et viverra justo commodo.



Manage your own medication experience

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aenean euismod bibendum laoreet. Proin gravida dolor sit amet lacus accumsan et viverra justo commodo.

Sign up today to manage your medications today!



Create your account



SIGN UP

LOG IN

Advil, Motrin IB (Ibuprofen)

+ Add to my medicine


Prescription or OTC

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aenean euismod bibendum laoreet. Proin gravida dolor sit amet lacus accumsan et viverra justo commodo. Proin sodales pulvinar tempor. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Nam fermentum, nulla luctus pharetra vulputate, felis tellus mollis orci, sed rhoncus sapien nunc eget odio.

Used for: Pain Fever Headache Migraine

Overview Reviews 100 Alternatives

Effectiveness 100 reviews



Side effects

Upset stomach	<div><div></div><div></div><div></div><div></div><div></div></div>	severity: severe
High blood pressure	<div><div></div><div></div><div></div><div></div><div></div></div>	severity: strong
Difficult urinating	<div><div></div><div></div><div></div><div></div><div></div></div>	severity: strong
Unexplained weight gain	<div><div></div><div></div><div></div><div></div><div></div></div>	severity: mild
Headache	<div><div></div><div></div><div></div><div></div><div></div></div>	severity: light

Suggested Usage

Directions use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. use in a well ventilated area maximal treatment area for one applicator is approximately 13.2 in. × 13.2 in. (1126 cm 2). remove applicator from package; do not touch sponge hold the applicator with the sponge down. Pinch wing only once to activate the ampules and release the antiseptic. wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin completely wet the treatment area with antiseptic dry surgical sites (e.g., abdomen or arm): use gentle back-and-forth strokes for 30 seconds moist surgical sites (e.g., inguinal fold): use gentle back-and-forth strokes for 2 minutes do not allow solution to pool; tuck prep towels to absorb solution, and then remove allow the solution to completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away. discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Interactivity with other medications or food

Celexa	<div><div></div><div></div><div></div><div></div><div></div></div>	severity: severe
Cymbalta	<div><div></div><div></div><div></div><div></div><div></div></div>	severity: strong
Effexor	<div><div></div><div></div><div></div><div></div><div></div></div>	severity: strong
Lexapro	<div><div></div><div></div><div></div><div></div><div></div></div>	severity: mild
Prozac	<div><div></div><div></div><div></div><div></div><div></div></div>	severity: light



search

12

Q

My Medications

My Daily schedules

Add a new medication

Print

Share



Active from Apr 11, 15

Cymbalta (Duloxetine)

Use for headache

Reminder: every day 8:00 am

Expand to see dosage and warnings

Prescription on date Apr 11, 15

Write a review

End taking

