

Day B

Thursday, September 24, 2026 — light recovery session (left shoulder and left hip/thigh)

Stop rule: stop any exercise that brings on or worsens the left shoulder or left buttock/thigh irritation. Mild achiness that settles is fine. Sharp, burning, tingling, or pain that spreads is a stop. Not working through it today.

Lighter session, five exercises instead of nine. Skipped this time: Sumo Deadlift, Reverse Lunge, Suitcase Carry (heavy hip loading, and a one-sided carry), and Overhead Press (already out for the shoulder). Everything left is at or below 9/17's load, nothing progresses. Since sitting bothers the hip and walking eases it, stand or walk around between exercises rather than resting seated.

Last session (9/17)	Exercise	Sets × Reps (today)	Weight (today)	Actual
— (replaces Overhead Press)	Dumbbell Goblet Walk	3 × 30 ft (<i>walking is the one thing that eases the hip, so this one should feel fine</i>)	12.5 lb	
green band × 13/13/13	Band Bent-Over Row	3 × 10–13 (<i>stay more upright than usual, only a slight lean</i>)	green band	
5 lb × 20/20/20	Stability Ball Hip Bridge	2 × 10 (<i>gentle, bodyweight, skip it if it stirs up the buttock</i>)	bodyweight	
10 lb × 55/55 sec/side	Side Plank	2 × 30 sec/side (<i>no weight; stop if the left hip complains</i>)	bodyweight	
green band × 15/15/15, 2-sec pause	Band Face Pull	3 × 15, same 2-sec pause	green band	
10 lb × 8/8/8	Dumbbell Lateral Raise	2 × 10 (<i>stop if it pinches the front of the shoulder</i>)	8 lb	

Notes: This is a training adjustment, not a diagnosis. The hip pattern you described (irritation, stays above the knee, worse with sitting, better with walking) fits muscle-type irritation more than a nerve problem, but it is still worth having a physical therapist or your doctor look at the shoulder and hip together, and sooner if you get numbness, tingling, weakness, pain below the knee, night pain, or any bladder or bowel change. Walks and the easy hikes are fine as long as they stay flat and easy; skip the hill warm-up until the hip settles. Once both areas feel normal, the next worksheet can bring the deadlift and lunge back at a lower weight before building up again.