

Initiating Search

November 10, 2025, 2:40 PM

• Search:

CAS Lexicon Search:

Jones oxidation - Preferred Concept

Search Tasks

Task	Result Type	View
Returned Reference Results from CAS Lexicon (148)	 References	View Results
Exported: Retrieved Related Reaction Results + Filters (111)	 Reactions	View Results
Filtered By:		
Yield:	90-100%, 80-89%, 70-79%	
Document Type:	Journal	

Copyright © 2025 American Chemical Society (ACS). All Rights Reserved.


Internal use only. Redistribution is subject to the terms of your CAS SciFinder License Agreement and CAS information Use Policies.

Reactions (50)

[View in CAS SciFinder](#)

Scheme 1 (1 Reaction)

Steps: 1 Yield: 100%



31-614-CAS-38455009

Steps: 1 Yield: 100%

Discovery of a novel tetrahydroimidazo[1,2-a]pyridine-5-carboxylic acid derivative as a potent and selective heparanase-1 inhibitor utilizing an improved synthetic approach

By: Imai, Yudai; et al

Bioorganic & Medicinal Chemistry Letters (2023), 97, 129543.


1.1 **Reagents:** N-methylmorpholine N-oxide
Solvents: Chloroform; 20 min, rt

1.2 **Catalysts:** TPAP; 30 min, 0 °C; 2 h, rt

Experimental Protocols

Scheme 2 (1 Reaction)

Steps: 1 Yield: 100%


 Suppliers (52)

 Suppliers (11)

31-614-CAS-34334478

Steps: 1 Yield: 100%

Selectivity of Oxidizing Agents toward Axial and Equatorial Hydroxyl Groups

By: Kaspar, Miroslav; et al

Journal of Organic Chemistry (2022), 87(14), 9157-9170.


1.1 **Reagents:** Chromium trioxide
Solvents: Acetone, Water; 0 °C; 30 min, 0 °C

1.2 **Reagents:** Isopropanol

Experimental Protocols

Scheme 3 (1 Reaction)

Steps: 1 Yield: 100%



 Suppliers (15)

 Suppliers (45)

31-480-CAS-15048430	Steps: 1 Yield: 100%	Silica gel supported Jones reagent (SJR): a simple, versatile, and efficient reagent for oxidation of alcohols in non-aqueous media By: Ali, Mohammed Hashmat; et al Synthetic Communications (2001), 31(21), 3383-3393.
1.1 Solvents: Dichloromethane		
Experimental Protocols		

Scheme 4 (1 Reaction)

Steps: 1 Yield: 100%




Suppliers (78)

Suppliers (75)

31-480-CAS-4394770	Steps: 1 Yield: 100%	Silica gel supported Jones reagent (SJR): a simple, versatile, and efficient reagent for oxidation of alcohols in non-aqueous media By: Ali, Mohammed Hashmat; et al Synthetic Communications (2001), 31(21), 3383-3393.
1.1 Solvents: Dichloromethane		
Experimental Protocols		

Scheme 5 (2 Reactions)

Steps: 1 Yield: 98-100%



Absolute stereochemistry shown

Suppliers (103)

Absolute stereochemistry shown,
Rotation (+)

Suppliers (68)

31-614-CAS-32794144	Steps: 1 Yield: 100%	Synthesis, In Vitro and In Silico Analysis of New Oleanolic Acid and Lupeol Derivatives against Leukemia Cell Lines: Involvement of the NF-κB Pathway By: Fontana, Gianfranco; et al International Journal of Molecular Sciences (2022), 23(12), 6594.
1.1 Reagents: Jones reagent Solvents: Acetone; 0 °C		
Experimental Protocols		

31-480-CAS-15392472	Steps: 1 Yield: 98%	Convenient and chromatography-free partial syntheses of maslinic acid and augustic acid By: Sommerwerk, Sven; et al Tetrahedron Letters (2014), 55(37), 5156-5158.
1.1 Reagents: Silica Solvents: Acetone; 30 min, reflux; reflux → 0 °C		
1.2 Reagents: Chromium trioxide, Sulfuric acid Solvents: Water; 0 °C; 30 min, 0 °C		
1.3 Solvents: Methanol; 30 min, 0 °C		
Experimental Protocols		

Scheme 6 (1 Reaction)

Steps: 1 Yield: 99%



Suppliers (71)

Suppliers (85)

31-480-CAS-8986407

Steps: 1 Yield: 99%

Silica gel supported Jones reagent (SJR): a simple, versatile, and efficient reagent for oxidation of alcohols in non-aqueous media

1.1 Solvents: Dichloromethane


Experimental Protocols

By: Ali, Mohammed Hashmat; et al

Synthetic Communications (2001), 31(21), 3383-3393.

Scheme 7 (2 Reactions)

Steps: 1 Yield: 97-99%



Suppliers (97)

Suppliers (136)

31-480-CAS-3765084

Steps: 1 Yield: 99%

Silica gel supported Jones reagent (SJR): a simple, versatile, and efficient reagent for oxidation of alcohols in non-aqueous media

1.1 Solvents: Dichloromethane

Experimental Protocols

By: Ali, Mohammed Hashmat; et al

Synthetic Communications (2001), 31(21), 3383-3393.

31-480-CAS-7236181

Steps: 1 Yield: 97%

A new efficient oxidation of alcohols with Jones reagent supported on Kieselguhr

1.1 Solvents: Dichloromethane; 10 min, rt


Experimental Protocols

By: Lou, Ji-Dong; et al

Synthesis and Reactivity in Inorganic, Metal-Organic, and Nano-Metal Chemistry (2006), 36(5), 381-383.

Scheme 8 (1 Reaction)

Steps: 1 Yield: 98%




Absolute stereochemistry shown

Absolute stereochemistry shown

31-480-CAS-216417	Steps: 1 Yield: 98%	Straightforward asymmetric synthesis of Ala-Ψ[CF:CH]-Pro, a proline-containing pseudodipeptide bearing a fluoroolefin as a peptide bond mimic By: Dutheuil, Guillaume; et al New Journal of Chemistry (2013), 37(5), 1320-1325.
1.1 Reagents: Dipyridinium dichromate Solvents: Dichloromethane; 2 h, rt		
Experimental Protocols		

Scheme 9 (1 Reaction)


Steps: 1 Yield: 98%



31-480-CAS-16285018	Steps: 1 Yield: 98%	A one-pot tandem strategy in catalytic asymmetric vinylogous aldol reaction of homoallylic alcohols By: Hou, Xufeng; et al Molecules (2016), 21(7), 842/1-842/12.
1.1 Solvents: <i>tert</i> -Butylbenzene; 3.5 - 7 h, 0 °C 1.2 10 min, -10 °C 1.3 Reagents: Trisodium phosphate Catalysts: <i>N</i> -[3,5-Bis(trifluoromethyl)phenyl]- <i>N'</i> -[(1 <i>R</i> ,2 <i>R</i>)-2-(1-pyrrolidinyl)cyclohexyl]thiourea Solvents: <i>tert</i> -Butylbenzene; -10 °C; 60 h, -10 °C		
Experimental Protocols		

Scheme 10 (1 Reaction)


Steps: 1 Yield: 98%



31-480-CAS-8380006	Steps: 1 Yield: 98%	Silica gel supported Jones reagent (SJR): a simple, versatile, and efficient reagent for oxidation of alcohols in non-aqueous media By: Ali, Mohammed Hashmat; et al Synthetic Communications (2001), 31(21), 3383-3393.
1.1 Solvents: Dichloromethane		
Experimental Protocols		

Scheme 11 (1 Reaction)


Steps: 1 Yield: 97%



31-480-CAS-9449306	Steps: 1 Yield: 97%	Conversion of 1,4-Diketones into para-Disubstituted Benzenes By: Ziffle, Vincent E.; et al Journal of Organic Chemistry (2010), 75(23), 8024-8038.
1.1 Solvents: Acetone; rt → 0 °C		
1.2 Solvents: Acetone; 0 °C; 10 min, 0 °C		
1.3 Solvents: Methanol; 30 min, 0 °C → rt Experimental Protocols		

Scheme 12 (1 Reaction)

Steps: 1 Yield: 97%




Suppliers (3)

31-480-CAS-11256909	Steps: 1 Yield: 97%	Incorporation of a Michael acceptor enhances the antitumor activity of triterpenoic acids By: Heller, Lucie; et al European Journal of Medicinal Chemistry (2015), 101, 391-399.
1.1 Reagents: Chromium trioxide, Sulfuric acid Solvents: Acetone; 0 °C; 1 h, rt		
Experimental Protocols		

Scheme 13 (1 Reaction)

Steps: 1 Yield: 97%




Suppliers (51)

Suppliers (77)

31-480-CAS-15370470	Steps: 1 Yield: 97%	Silica gel supported Jones reagent (SJR): a simple, versatile, and efficient reagent for oxidation of alcohols in non-aqueous media By: Ali, Mohammed Hashmat; et al Synthetic Communications (2001), 31(21), 3383-3393.
1.1 Solvents: Dichloromethane		
Experimental Protocols		

Scheme 14 (1 Reaction)

Steps: 1 Yield: 97%




Suppliers (149)

Suppliers (69)

31-480-CAS-9989180	Steps: 1 Yield: 97%	A new efficient oxidation of alcohols with Jones reagent supported on Kieselguhr
1.1 Solvents: Dichloromethane; 10 min, rt		By: Lou, Ji-Dong; et al
Experimental Protocols		Synthesis and Reactivity in Inorganic, Metal-Organic, and Nano-Metal Chemistry (2006), 36(5), 381-383.

Scheme 15 (1 Reaction)


Steps: 1 Yield: 96%



31-480-CAS-6952935	Steps: 1 Yield: 96%	Conversion of 1,4-Diketones into para-Disubstituted Benzenes
1.1 Reagents: Oxalyl chloride		By: Ziffle, Vincent E.; et al
Solvents: Dichloromethane; rt → -78 °C		Journal of Organic Chemistry (2010), 75(23), 8024-8038.
1.2 Reagents: Dimethyl sulfoxide		
Solvents: Dichloromethane; 15 min, -78 °C		
1.3 35 min, -78 °C		
1.4 Reagents: Triethylamine; 5 min, -78 °C; 25 min, -78 °C → rt		
1.5 Solvents: Water; rt		
Experimental Protocols		

Scheme 16 (1 Reaction)


Steps: 1 Yield: 96%



31-480-CAS-2363977	Steps: 1 Yield: 96%	Silica gel supported Jones reagent (SJR): a simple, versatile, and efficient reagent for oxidation of alcohols in non-aqueous media
1.1 Solvents: Dichloromethane		By: Ali, Mohammed Hashmat; et al
Experimental Protocols		Synthetic Communications (2001), 31(21), 3383-3393.

Scheme 17 (2 Reactions)


Steps: 1 Yield: 93-96%



31-480-CAS-8702378 Steps: 1 Yield: 96% 1.1 Solvents: Dichloromethane Experimental Protocols	Silica gel supported Jones reagent (SJR): a simple, versatile, and efficient reagent for oxidation of alcohols in non-aqueous media By: Ali, Mohammed Hashmat; et al Synthetic Communications (2001), 31(21), 3383-3393.
31-480-CAS-5242103 Steps: 1 Yield: 93% 1.1 Solvents: Dichloromethane; 10 min, rt Experimental Protocols	A new efficient oxidation of alcohols with Jones reagent supported on Kieselguhr By: Lou, Ji-Dong; et al Synthesis and Reactivity in Inorganic, Metal-Organic, and Nano-Metal Chemistry (2006), 36(5), 381-383.

Scheme 18 (1 Reaction)


Steps: 1 Yield: 95%



31-480-CAS-19316871 Steps: 1 Yield: 95% 1.1 Reagents: Sulfuric acid Solvents: Acetone, Water; 2 h, rt Experimental Protocols	A Method for Synthesis of 3- Hydroxy-1-indanones via Cu-Catalyzed Intramolecular Annulation Reactions By: He, Guoxue; et al Journal of Organic Chemistry (2018), 83(21), 13356-13362.
--	--

Scheme 19 (1 Reaction)


Steps: 1 Yield: 95%



31-480-CAS-16254164 Steps: 1 Yield: 95% 1.1 Solvents: <i>tert</i>-Butylbenzene; 3.5 - 7 h, 0 °C 1.2 10 min, -10 °C 1.3 Reagents: Trisodium phosphate Catalysts: <i>N</i>-[3,5-Bis(trifluoromethyl)phenyl]-<i>N'</i>-[(1<i>R</i>,2<i>R</i>)-2-(1-pyrrolidinyl)cyclohexyl]thiourea Solvents: <i>tert</i>-Butylbenzene; -10 °C; 38 h, -10 °C Experimental Protocols	A one-pot tandem strategy in catalytic asymmetric vinylogous aldol reaction of homoallylic alcohols By: Hou, Xufeng; et al Molecules (2016), 21(7), 842/1-842/12.
--	--

Scheme 20 (1 Reaction)

Steps: 1 Yield: 95%



Supplier (1)

Supplier (1)

31-480-CAS-3385223

Steps: 1 Yield: 95%

Conversion of 1,4-Diketones into para-Disubstituted Benzenes

By: Ziffle, Vincent E.; et al

Journal of Organic Chemistry (2010), 75(23), 8024-8038.

1.1 Solvents: Acetone; rt → 0 °C


1.2 Solvents: Acetone; 0 °C; 20 min, 0 °C

1.3 Solvents: Methanol; 30 min, 0 °C → rt

Experimental Protocols

Scheme 21 (1 Reaction)

Steps: 1 Yield: 95%



Suppliers (105)

Suppliers (128)

31-480-CAS-14255248

Steps: 1 Yield: 95%

A new efficient oxidation of alcohols with Jones reagent supported on Kieselguhr

By: Lou, Ji-Dong; et al


Synthesis and Reactivity in Inorganic, Metal-Organic, and Nano-Metal Chemistry (2006), 36(5), 381-383.

1.1 Solvents: Dichloromethane; 10 min, rt

Experimental Protocols

Scheme 22 (1 Reaction)

Steps: 1 Yield: 94%



Relative stereochemistry shown

Relative stereochemistry shown

Suppliers (9)

Supplier (1)

31-480-CAS-17518248

Steps: 1 Yield: 94%

A diastereoselective synthesis of boceprevir's gem-dimethyl bicyclic [3.1.0] proline intermediate from an insecticide ingredient cis-cypermethric acid

By: Kallam, Srinivasa Reddy; et al

Tetrahedron (2017), 73(30), 4285-4294.

1.1 Reagents: Oxalyl chloride

Catalysts: Dimethyl sulfoxide


Solvents: Dichloromethane; 30 min, -78 °C

1.2 Reagents: Triethylamine

Solvents: Dichloromethane; -78 °C → 35 °C; 45 min, 30 - 35 °C

Scheme 23 (1 Reaction)

Steps: 1 Yield: 94%



31-480-CAS-16284214

Steps: 1 Yield: 94%

A one-pot tandem strategy in catalytic asymmetric vinylogous aldol reaction of homoallylic alcohols

By: Hou, Xufeng; et al

Molecules (2016), 21(7), 842/1-842/12.

1.1 Solvents: *tert*-Butylbenzene; 3.5 - 7 h, 0 °C

1.2 10 min, -10 °C


1.3 Reagents: Trisodium phosphate

Catalysts: *N*-[3,5-Bis(trifluoromethyl)phenyl]-*N'*-[(1*R*,2*R*)-2-(1-pyrrolidinyl)cyclohexyl]thioureaSolvents: *tert*-Butylbenzene; -10 °C; 61 h, -10 °C

Experimental Protocols

Scheme 24 (1 Reaction)

Steps: 1 Yield: 94%



Suppliers (132)

Suppliers (81)

31-480-CAS-4075671

Steps: 1 Yield: 94%

Silica gel supported Jones reagent (SJR): a simple, versatile, and efficient reagent for oxidation of alcohols in non-aqueous media

By: Ali, Mohammed Hashmat; et al


Synthetic Communications (2001), 31(21), 3383-3393.

1.1 Solvents: Dichloromethane

Experimental Protocols

Scheme 25 (1 Reaction)

Steps: 1 Yield: 94%



Suppliers (59)

Suppliers (78)

31-480-CAS-9311643

Steps: 1 Yield: 94%

Silica gel supported Jones reagent (SJR): a simple, versatile, and efficient reagent for oxidation of alcohols in non-aqueous media

By: Ali, Mohammed Hashmat; et al


Synthetic Communications (2001), 31(21), 3383-3393.

1.1 Solvents: Dichloromethane

Experimental Protocols

Scheme 26 (1 Reaction)

Steps: 1 Yield: 94%



Suppliers (71)

Suppliers (105)

31-480-CAS-8050440

Steps: 1 Yield: 94%

1.1 Solvents: Dichloromethane

Experimental Protocols


Silica gel supported Jones reagent (SJR): a simple, versatile, and efficient reagent for oxidation of alcohols in non-aqueous media

By: Ali, Mohammed Hashmat; et al

Synthetic Communications (2001), 31(21), 3383-3393.

Scheme 27 (1 Reaction)

Steps: 1 Yield: 93%



Suppliers (90)

Suppliers (87)

31-480-CAS-9039554

Steps: 1 Yield: 93%

1.1 Solvents: Dichloromethane

Experimental Protocols


Silica gel supported Jones reagent (SJR): a simple, versatile, and efficient reagent for oxidation of alcohols in non-aqueous media

By: Ali, Mohammed Hashmat; et al

Synthetic Communications (2001), 31(21), 3383-3393.

Scheme 28 (1 Reaction)

Steps: 1 Yield: 93%



Suppliers (60)

Suppliers (74)

31-480-CAS-2999003

Steps: 1 Yield: 93%

1.1 Solvents: Dichloromethane

Experimental Protocols


Silica gel supported Jones reagent (SJR): a simple, versatile, and efficient reagent for oxidation of alcohols in non-aqueous media

By: Ali, Mohammed Hashmat; et al

Synthetic Communications (2001), 31(21), 3383-3393.

Scheme 29 (1 Reaction)

Steps: 1 Yield: 93%



Suppliers (86)

Suppliers (52)

31-480-CAS-12125394

Steps: 1 Yield: 93%

A new efficient oxidation of alcohols with Jones reagent supported on Kieselguhr

By: Lou, Ji-Dong; et al


Synthesis and Reactivity in Inorganic, Metal-Organic, and Nano-Metal Chemistry (2006), 36(5), 381-383.

1.1 Solvents: Dichloromethane; 10 min, rt

Experimental Protocols

Scheme 30 (1 Reaction)

Steps: 1 Yield: 92%



Suppliers (19)

Suppliers (85)

Absolute stereochemistry shown,
Rotation (-)
Double bond geometry shown

31-480-CAS-16254171

Steps: 1 Yield: 92%

A one-pot tandem strategy in catalytic asymmetric vinylogous aldol reaction of homoallylic alcohols

By: Hou, Xufeng; et al

Molecules (2016), 21(7), 842/1-842/12.

1.1 Solvents: *tert*-Butylbenzene; 3.5 - 7 h, 0 °C

1.2 10 min, -10 °C

1.3 Reagents: Trisodium phosphate


Catalysts: *N*-[3,5-Bis(trifluoromethyl)phenyl]-*N'*-[(1*R*,2*R*)-2-(1-pyrrolidinyl)cyclohexyl]thiourea

Solvents: *tert*-Butylbenzene; -10 °C; 59 h, -10 °C

Experimental Protocols

Scheme 31 (1 Reaction)

Steps: 1 Yield: 92%



Suppliers (23)

Suppliers (85)

Absolute stereochemistry shown,
Rotation (-)
Double bond geometry shown

31-480-CAS-16254174

Steps: 1 Yield: 92%

A one-pot tandem strategy in catalytic asymmetric vinylogous aldol reaction of homoallylic alcohols

By: Hou, Xufeng; et al

Molecules (2016), 21(7), 842/1-842/12.

1.1 Solvents: *tert*-Butylbenzene; 3.5 - 7 h, 0 °C

1.2 10 min, -10 °C

1.3 Reagents: Trisodium phosphate


Catalysts: *N*-[3,5-Bis(trifluoromethyl)phenyl]-*N'*-[(1*R*,2*R*)-2-(1-pyrrolidinyl)cyclohexyl]thiourea

Solvents: *tert*-Butylbenzene; -10 °C; 64 h, -10 °C

Experimental Protocols

Scheme 32 (1 Reaction)

Steps: 1 Yield: 92%



Suppliers (35)

Suppliers (67)

31-480-CAS-7274479

Steps: 1 Yield: 92%

Silver-catalysed intramolecular cyclisation of 2-alkynylacetophenones and 3-acetyl-2-alkynylpyridines in the presence of ammonia


By: Dell'Acqua, Monica; et al

Organic & Biomolecular Chemistry (2011), 9(22), 7836-7848.

1.1 Reagents: Chromium trioxide
Solvents: Acetone; -70 - 25 °C

Scheme 33 (1 Reaction)

Steps: 1 Yield: 92%



Suppliers (63)

Suppliers (74)

31-480-CAS-10921634

Steps: 1 Yield: 92%

Oxidation of benzoin to benzil with the Jones' reagent supported on graphite


By: Lou, Ji-Dong; et al

Oxidation Communications (2010), 33(3), 550-554.

1.1 Reagents: Jones reagent
Solvents: Dichloromethane; 1 h, rt; 1 h, rt

Scheme 34 (1 Reaction)

Steps: 1 Yield: 92%



Suppliers (118)

Suppliers (101)

31-480-CAS-39192

Steps: 1 Yield: 92%

Oxidation of benzoin to benzil with the Jones' reagent supported on graphite


By: Lou, Ji-Dong; et al

Oxidation Communications (2010), 33(3), 550-554.

1.1 Reagents: Jones reagent
Solvents: Dichloromethane; 1 h, rt; 1 h, rt

Scheme 35 (1 Reaction)

Steps: 1 Yield: 91%



31-480-CAS-16254175

Steps: 1 Yield: 91%

A one-pot tandem strategy in catalytic asymmetric vinylogous aldol reaction of homoallylic alcohols

1.1 Solvents: *tert*-Butylbenzene; 3.5 - 7 h, 0 °C

1.2 10 min, -10 °C

1.3 Reagents: Trisodium phosphate

Catalysts: *N*-[3,5-Bis(trifluoromethyl)phenyl]-*N'*-[(1*R*,2*R*)-2-(1-pyrrolidinyl)cyclohexyl]thioureaSolvents: *tert*-Butylbenzene; -10 °C; 61 h, -10 °C


By: Hou, Xufeng; et al

Molecules (2016), 21(7), 842/1-842/12.

Experimental Protocols

Scheme 36 (1 Reaction)

Steps: 1 Yield: 91%



31-480-CAS-1243849

Steps: 1 Yield: 91%

Silver-catalysed intramolecular cyclisation of 2-alkynylacetophenones and 3-acetyl-2-alkynylpyridines in the presence of ammonia

1.1 Reagents: Chromium trioxide


Solvents: Acetone; -70 - 25 °C

By: Dell'Acqua, Monica; et al

Organic & Biomolecular Chemistry (2011), 9(22), 7836-7848.

Scheme 37 (1 Reaction)

Steps: 1 Yield: 90%



31-480-CAS-396628

Steps: 1 Yield: 90%

Conversion of 1,4-Diketones into para-Disubstituted Benzenes

1.1 Solvents: Acetone; rt → 0 °C

1.2 Solvents: Acetone; 0 °C; 30 min, 0 °C

1.3 Solvents: Methanol; 30 min, 0 °C → rt


By: Ziffle, Vincent E.; et al

Journal of Organic Chemistry (2010), 75(23), 8024-8038.

Experimental Protocols

Scheme 38 (1 Reaction)

Steps: 1 Yield: 90%



31-301-CAS-21165257

Steps: 1 Yield: 90%

Catalytic Environmentally Friendly Protocol for Achmatowicz Rearrangement


By: Li, Zhilong; et al

Journal of Organic Chemistry (2016), 81(11), 4847-4855.

1.1 **Reagents:** Sodium bicarbonate, Potassium peroxymonosulfate sulfate (2KHSO₅.KHSO₄.K₂SO₄)**Catalysts:** Potassium bromide**Solvents:** Tetrahydrofuran, Water; 0 °C; 0.5 h, 0 °C1.2 **Reagents:** Sodium bicarbonate**Solvents:** Water

Scheme 39 (1 Reaction)

Steps: 1 Yield: 90%



Supplier (1)

Supplier (1)

31-480-CAS-12516816

Steps: 1 Yield: 90%

Conversion of 1,4-Diketones into para-Disubstituted Benzenes

By: Ziffle, Vincent E.; et al


Journal of Organic Chemistry (2010), 75(23), 8024-8038.

1.1 **Solvents:** Acetone; rt → 0 °C1.2 **Solvents:** Acetone; 0 °C; 10 min, 0 °C1.3 **Solvents:** Methanol; 30 min, 0 °C → rt

Experimental Protocols

Scheme 40 (1 Reaction)

Steps: 1 Yield: 90%



Suppliers (30)

Suppliers (66)

31-480-CAS-8796682

Steps: 1 Yield: 90%

Oxidation of benzoin to benzil with the Jones' reagent supported on graphite


By: Lou, Ji-Dong; et al

Oxidation Communications (2010), 33(3), 550-554.

1.1 **Reagents:** Jones reagent**Solvents:** Dichloromethane; 1 h, rt; 1 h, rt

Scheme 41 (1 Reaction)

Steps: 1 Yield: 90%



Suppliers (103)

Suppliers (12)

31-614-CAS-37906935

Steps: 1 Yield: 90%

Anti-Diabetic Activity of Glycyrrhetic Acid Derivatives FC-114 and FC-122: Scale-Up, In Silico, In Vitro, and In Vivo Studies

By: Alvarez-Almazan, Samuel; et al

International Journal of Molecular Sciences (2023), 24(16), 12812.


1.1 Reagents: Sulfuric acid

Catalysts: Chromium trioxide

Solvents: Tetrahydrofuran; 1.5 h, rt

Scheme 42 (1 Reaction)

Steps: 1 Yield: 90%



Suppliers (63)

Suppliers (68)

31-480-CAS-21099077

Steps: 1 Yield: 90%

Preparation of β^2 -homologous amino acids bearing polar side chains via a collective synthesis strategy

By: Liu, Shi; et al

Journal of Organic Chemistry (2020), 85(3), 1718-1724.


1.1 Reagents: 1-Methylimidazole, Tempo

Catalysts: 2,2'-Bipyridine, Copper bromide (Cu Br)

Solvents: Acetonitrile; overnight, rt

Scheme 43 (1 Reaction)

Steps: 1 Yield: 89%



31-480-CAS-13660654

Steps: 1 Yield: 89%

One-pot preparation of hydroxylated potassium organotrifluoroborates and subsequent Jones oxidation to potassium organocarbonyltrifluoroborates

By: Kim, Dong-Su; et al

Tetrahedron (2011), 67(6), 1062-1070.


1.1 Solvents: Acetone; 10 min, 0 °C

1.2 Reagents: Potassium carbonate; 5 min, rt

Experimental Protocols

Scheme 44 (1 Reaction)

Steps: 1 Yield: 89%



Suppliers (65)

Suppliers (25)

31-480-CAS-867083

Steps: 1 Yield: 89%

A new efficient oxidation of alcohols with Jones reagent supported on Kieselguhr

1.1 Solvents: Dichloromethane; 10 min, rt


By: Lou, Ji-Dong; et al

Experimental Protocols

Synthesis and Reactivity in Inorganic, Metal-Organic, and Nano-Metal Chemistry (2006), 36(5), 381-383.

Scheme 45 (1 Reaction)

Steps: 1 Yield: 89%



Suppliers (34)

Suppliers (127)

31-480-CAS-1655878

Steps: 1 Yield: 89%

Silica gel supported Jones reagent (SJR): a simple, versatile, and efficient reagent for oxidation of alcohols in non-aqueous media

1.1 Solvents: Dichloromethane


By: Ali, Mohammed Hashmat; et al

Experimental Protocols

Synthetic Communications (2001), 31(21), 3383-3393.

Scheme 46 (1 Reaction)

Steps: 1 Yield: 88%



Absolute stereochemistry shown

Absolute stereochemistry shown,
Rotation (+)

Suppliers (21)

Suppliers (11)

31-480-CAS-12558386

Steps: 1 Yield: 88%

Incorporation of a Michael acceptor enhances the antitumor activity of triterpenoic acids

1.1 Reagents: Chromium trioxide, Sulfuric acid

By: Heller, Lucie; et al


Solvents: Acetone; 0 °C; 1 h, rt

Experimental Protocols

European Journal of Medicinal Chemistry (2015), 101, 391-399.


Scheme 47 (1 Reaction)

Steps: 1 Yield: 88%



Absolute stereochemistry shown,
Rotation (+)

Suppliers (2)



Absolute stereochemistry shown

Supplier (1)

31-480-CAS-9461591

Steps: 1 Yield: 88%

Incorporation of a Michael acceptor enhances the antitumor activity of triterpenoic acids

By: Heller, Lucie; et al

European Journal of Medicinal Chemistry (2015), 101, 391-399.

1.1 **Reagents:** Chromium trioxide, Sulfuric acid

Solvents: Acetone; 0 °C; 1 h, rt

Experimental Protocols