

Initiating Search

November 11, 2025, 2:59 PM

• Search:

CAS Lexicon Search:

Aryl azides - Preferred Concept

Search Tasks

Task	Result Type	View
Returned Reference Results from CAS Lexicon (1,914)	 References	View Results
Exported: Retrieved Related Reaction Results (389)	 Reactions	View Results

Copyright © 2025 American Chemical Society (ACS). All Rights Reserved.

Internal use only. Redistribution is subject to the terms of your CAS SciFinder License Agreement and CAS information Use Policies.

Reactions (50)

[View in CAS SciFinder](#)

Scheme 1 (1 Reaction)

Steps: 1 Yield: 99%



31-043-CAS-971240

Steps: 1 Yield: 99%

Designing a Cu(II)-ArCu(II)-ArCu(III)-Cu(I) Catalytic Cycle: Cu(II)-Catalyzed Oxidative Arene C-H Bond Azidation with Air as an Oxidant under Ambient Conditions

By: Yao, Bo; et al

Journal of Organic Chemistry (2014), 79(22), 11139-11145.

1.1 Reagents: *p*-Toluenesulfonic acid, Oxygen, Sodium azide
Catalysts: Cupric perchlorate
Solvents: Acetonitrile, Dichloromethane, Water; 24 h, rt

Experimental Protocols

Scheme 2 (1 Reaction)

Steps: 1 Yield: 98%



31-043-CAS-10067947

Steps: 1 Yield: 98%

Designing a Cu(II)-ArCu(II)-ArCu(III)-Cu(I) Catalytic Cycle: Cu(II)-Catalyzed Oxidative Arene C-H Bond Azidation with Air as an Oxidant under Ambient Conditions

By: Yao, Bo; et al


Journal of Organic Chemistry (2014), 79(22), 11139-11145.

1.1 Reagents: *p*-Toluenesulfonic acid, Oxygen, Sodium azide
Catalysts: Cupric perchlorate
Solvents: Acetonitrile, Dichloromethane, Water; 24 h, rt

Experimental Protocols

Scheme 3 (1 Reaction)

Steps: 1 Yield: 96%



31-043-CAS-11628535

Steps: 1 Yield: 96%

Designing a Cu(II)-ArCu(II)-ArCu(III)-Cu(I) Catalytic Cycle: Cu(II)-Catalyzed Oxidative Arene C-H Bond Azidation with Air as an Oxidant under Ambient Conditions

By: Yao, Bo; et al


Journal of Organic Chemistry (2014), 79(22), 11139-11145.

1.1 Reagents: *p*-Toluenesulfonic acid, Oxygen, Sodium azide
Catalysts: Cupric perchlorate
Solvents: Acetonitrile, Dichloromethane, Water; 24 h, rt

Experimental Protocols

Scheme 4 (1 Reaction)

Steps: 1 Yield: 96%



Suppliers (71)

Suppliers (13)

31-043-CAS-4285568

Steps: 1 Yield: 96%

Synthesis of aryl azides via post-cleavage modification of polymer-bound triazenes

By: Avemaria, Frank; et al


Synlett (2004), (7), 1163-1166.

- 1.1 **Reagents:** Boron trifluoride etherate
Solvents: Dichloromethane; rt; rt → -15 °C
- 1.2 **Reagents:** Isoamyl nitrite; 0.25 h, -15 °C; 4 h, -15 °C; -15 °C → -70 °C
- 1.3 **Solvents:** Acetonitrile; -40 °C
- 1.4 **Reagents:** Pyridine
Solvents: Tetrahydrofuran; -20 °C; 6 h, -20 °C; -20 °C → rt
- 1.5 **Reagents:** Trifluoroacetic acid
Solvents: Dichloromethane; 2 - 5 min, rt
- 1.6 **Reagents:** Azidotrimethylsilane; 5 - 10 min, rt

Experimental Protocols

Scheme 5 (1 Reaction)

Steps: 1 Yield: 96%



Suppliers (54)

Suppliers (3)

31-043-CAS-6440204

Steps: 1 Yield: 96%

Synthesis of aryl azides via post-cleavage modification of polymer-bound triazenes

By: Avemaria, Frank; et al


Synlett (2004), (7), 1163-1166.

- 1.1 **Reagents:** Boron trifluoride etherate
Solvents: Dichloromethane; rt; rt → -15 °C
- 1.2 **Reagents:** Isoamyl nitrite; 0.25 h, -15 °C; 4 h, -15 °C; -15 °C → -70 °C
- 1.3 **Solvents:** Acetonitrile; -40 °C
- 1.4 **Reagents:** Pyridine
Solvents: Tetrahydrofuran; -20 °C; 6 h, -20 °C; -20 °C → rt
- 1.5 **Reagents:** Trifluoroacetic acid
Solvents: Dichloromethane; 2 - 5 min, rt
- 1.6 **Reagents:** Azidotrimethylsilane; 5 - 10 min, rt

Experimental Protocols

Scheme 6 (1 Reaction)

Steps: 1 Yield: 96%



Suppliers (83)

Suppliers (4)

31-043-CAS-11295932

Steps: 1 Yield: 96%

Formal C-H-Azidation Based Shortcut to Diazido Building Blocks for the Versatile Preparation of Photoaffinity Labeling Probes

By: Yoshida, Suguru; et al


European Journal of Organic Chemistry (2014), 2014(19), 3991-3995.

- 1.1 **Reagents:** Bis(pinacolato)diborane
Catalysts: Bis[(1,2,5,6-η)-1,5-cyclooctadiene]di-μ-methoxydi-
 iridium, 4,4'-Bis(1,1-dimethylethyl)-2,2'-bipyridine
Solvents: Hexane; 24 - 144 h, rt
- 1.2 **Reagents:** Azidotrimethylsilane, Potassium fluoride
Catalysts: Cuprous chloride
Solvents: Methanol; 4 - 24 h, reflux

Experimental Protocols

Scheme 7 (1 Reaction)

Steps: 1 Yield: 95%



Suppliers (69)

Supplier (1)

31-043-CAS-13448283

Steps: 1 Yield: 95%

Formal C-H-Azidation Based Shortcut to Diazido Building Blocks for the Versatile Preparation of Photoaffinity Labeling Probes

By: Yoshida, Suguru; et al

European Journal of Organic Chemistry (2014), 2014(19), 3991-3995.

- 1.1 **Reagents:** Bis(pinacolato)diborane
Catalysts: Bis[(1,2,5,6-η)-1,5-cyclooctadiene]di-μ-methoxydi-
 iridium, 4,4'-Bis(1,1-dimethylethyl)-2,2'-bipyridine
Solvents: Hexane; 24 - 144 h, rt
- 1.2 **Reagents:** Azidotrimethylsilane, Potassium fluoride
Catalysts: Cuprous chloride
Solvents: Methanol; 4 - 24 h, reflux

Experimental Protocols

Scheme 8 (1 Reaction)

Steps: 1 Yield: 94%



31-043-CAS-3083903

Steps: 1 Yield: 94%

Designing a Cu(II)-ArCu(II)-ArCu(III)-Cu(I) Catalytic Cycle: Cu(II)-Catalyzed Oxidative Arene C-H Bond Azidation with Air as an Oxidant under Ambient Conditions

By: Yao, Bo; et al

Journal of Organic Chemistry (2014), 79(22), 11139-11145.

- 1.1 **Reagents:** *p*-Toluenesulfonic acid, Oxygen, Sodium azide
Catalysts: Cupric perchlorate
Solvents: Acetonitrile, Dichloromethane, Water; 24 h, rt

Experimental Protocols

Scheme 9 (1 Reaction)

Steps: 1 Yield: 93%



31-043-CAS-7343421

Steps: 1 Yield: 93%

1.1 **Reagents:** *p*-Toluenesulfonic acid, Oxygen, Sodium azide
Catalysts: Cupric perchlorate
Solvents: Acetonitrile, Dichloromethane, Water; 48 h, rt

Experimental Protocols


Designing a Cu(II)-ArCu(II)-ArCu(III)-Cu(I) Catalytic Cycle: Cu(II)-Catalyzed Oxidative Arene C-H Bond Azidation with Air as an Oxidant under Ambient Conditions

By: Yao, Bo; et al

Journal of Organic Chemistry (2014), 79(22), 11139-11145.

Scheme 10 (1 Reaction)

Steps: 1 Yield: 91%



Suppliers (75)

31-043-CAS-9185226

Steps: 1 Yield: 91%

1.1 **Reagents:** Bis(pinacolato)diborane
Catalysts: Bis[(1,2,5,6-η)-1,5-cyclooctadiene]di-μ-methoxydi-*iridium*, 4,4'-Bis(1,1-dimethylethyl)-2,2'-bipyridine
Solvents: Hexane; 39 h, rt

1.2 **Reagents:** Azidotrimethylsilane, Potassium fluoride
Catalysts: Cuprous chloride
Solvents: Methanol; 7 h, reflux

Experimental Protocols

Formal C-H-Azidation Based Shortcut to Diazido Building Blocks for the Versatile Preparation of Photoaffinity Labeling Probes

By: Yoshida, Suguru; et al

European Journal of Organic Chemistry (2014), 2014(19), 3991-3995.

Scheme 11 (1 Reaction)

Steps: 1 Yield: 90%



31-043-CAS-5214796

Steps: 1 Yield: 90%

1.1 **Reagents:** *p*-Toluenesulfonic acid, Oxygen, Sodium azide
Catalysts: Cupric perchlorate
Solvents: Acetonitrile, Dichloromethane, Water; 24 h, rt

Experimental Protocols


Designing a Cu(II)-ArCu(II)-ArCu(III)-Cu(I) Catalytic Cycle: Cu(II)-Catalyzed Oxidative Arene C-H Bond Azidation with Air as an Oxidant under Ambient Conditions

By: Yao, Bo; et al

Journal of Organic Chemistry (2014), 79(22), 11139-11145.

Scheme 12 (1 Reaction)

Steps: 1 Yield: 90%



Suppliers (54)

Suppliers (3)

31-043-CAS-13820538

Steps: 1 Yield: 90%

The development of carbene-stabilized N-O radical coupling strategy in metal-free regioselective C-H azidation of quinoline N-oxides

By: Li, Pan; et al

Organic Chemistry Frontiers (2015), 2(10), 1313-1317.


1.1 **Reagents:** Bis(trifluoroacetoxy)iodobenzene, Azidotrimethylsilane

Solvents: Ethyl acetate; 10 min, 25 °C

Experimental Protocols

Scheme 13 (1 Reaction)

Steps: 1 Yield: 90%



Suppliers (84)

Suppliers (30)

31-043-CAS-13260206

Steps: 1 Yield: 90%

Simple and efficient method for the preparation of aryl azides using sonication

By: Telvekar, Vikas N.; et al

Synthetic Communications (2012), 42(7), 1085-1089.


1.1 **Reagents:** Sodium azide, Iodate(1-), dichloro-, sodium

Solvents: Water; 10 min, 25 °C

Experimental Protocols

Scheme 14 (2 Reactions)

Steps: 1 Yield: 88-90%



Suppliers (77)

Suppliers (2)

31-043-CAS-20594551

Steps: 1 Yield: 90%

Direct Azidation of Phenols

By: Kitamura, Mitsuru; et al

European Journal of Organic Chemistry (2019), 2019(33), 5824-5827.

1.1 **Reagents:** Diisopropylamine, 1*H*-imidazolium, 1,3-bis[2,6-bis(1-methylethyl)phenyl]-2-chloro-, chloride (1:1)


Solvents: *N,N,N',N'*-Tetramethylethylenediamine; 1 h, 23 °C

1.2 **Reagents:** Sodium azide; 8 h, 23 °C

1.3 **Solvents:** Water; 23 °C

31-043-CAS-21803960 Steps: 1 Yield: 88% 1.1 Reagents: Diisopropylamine, 1 <i>H</i>-imidazolium, 2-azido-1,3-bis [2,6-bis(1-methylethyl)phenyl]-, hexafluorophosphate(1-) (1:1) Solvents: 2-Methoxyethanol; 30 min, rt 1.2 Reagents: Sodium azide; 4 h, rt	Regent for electrophilic azidation and diazotization Assignee: Kyushu Institute of Technology Japan, JP2019094331 A 2019-06-20 PatentPak available
--	---

Scheme 15 (1 Reaction)


Steps: **1** Yield: **90%**

Suppliers (81)

Suppliers (14)

31-043-CAS-17684559 Steps: 1 Yield: 90% 1.1 Reagents: Zincate(1-), dichloro(2,2,6,6-tetramethyl-1-piperi dyl)-, lithium (1:1) Solvents: Tetrahydrofuran, Hexane; 1 h, rt; rt → 0 °C 1.2 Reagents: 1-Azido-1,2-benziodoxol-3(1 <i>H</i>)-one Catalysts: Copper(II) 2-ethylhexanoate Solvents: Tetrahydrofuran; 0 °C; 0 °C → rt; 3 h, rt 1.3 Reagents: Isopropanol	Site-Selective Copper-Catalyzed Amination and Azidation of Arenes and Heteroarenes via Deprotonative Zincation By: Hendrick, Charles E.; et al Journal of the American Chemical Society (2017), 139(33), 11622-11628.
---	--


Scheme 16 (1 Reaction)

Steps: **1** Yield: **89%**

Suppliers (64)

31-043-CAS-8126092 Steps: 1 Yield: 89% 1.1 Reagents: Iodobenzene diacetate, p-Toluenesulfonic acid monohydrate Catalysts: Bis[dichloro[η⁵-(pentamethylcyclopentadienyl)] rhodium] Solvents: Acetone; 15 min, rt 1.2 Reagents: Sodium azide; 16 h, 50 °C	Rhodium(III)-Catalyzed Azidation and Nitration of Arenes by C-H Activation By: Xie, Fang; et al Angewandte Chemie, International Edition (2013), 52(45), 11862-11866.
--	--

Scheme 17 (1 Reaction)

Steps: **1** Yield: **88%**


Suppliers (76)

Suppliers (2)

31-043-CAS-8800272	Steps: 1 Yield: 88%	Room-temperature Cu(II)-catalyzed aromatic C-H azidation for the synthesis of ortho-azido anilines with excellent regioselectivity
1.1 Reagents: Oxygen, 1-Azido-1,2-benziodoxol-3(1 <i>H</i>)-one Catalysts: Cupric acetate Solvents: Tetrahydrofuran; 12 h, 20 °C		By: Fan, Yunpeng; et al
Experimental Protocols		Chemical Communications (Cambridge, United Kingdom) (2014), 50(43), 5733-5736.

Scheme 18 (1 Reaction)


Steps: 1 Yield: 88%



31-043-CAS-15397581	Steps: 1 Yield: 88%	Simple and efficient method for the preparation of aryl azides using sonication
1.1 Reagents: Sodium azide, Iodate(1-), dichloro-, sodium Solvents: Water; 10 min, 25 °C		By: Telvekar, Vikas N.; et al
Experimental Protocols		Synthetic Communications (2012), 42(7), 1085-1089.

Scheme 19 (1 Reaction)


Steps: 1 Yield: 88%



31-043-CAS-7647356	Steps: 1 Yield: 88%	Designing a Cu(II)-ArCu(II)-ArCu(III)-Cu(I) Catalytic Cycle: Cu(II)-Catalyzed Oxidative Arene C-H Bond Azidation with Air as an Oxidant under Ambient Conditions
1.1 Reagents: <i>p</i> -Toluenesulfonic acid, Oxygen, Sodium azide Catalysts: Cupric perchlorate Solvents: Acetonitrile, Dichloromethane, Water; 20 h, rt		By: Yao, Bo; et al
Experimental Protocols		Journal of Organic Chemistry (2014), 79(22), 11139-11145.

Scheme 20 (2 Reactions)


Steps: 1 Yield: 87-88%



31-043-CAS-5505056 Steps: 1 Yield: 88% 1.1 Reagents: Iodobenzene diacetate, p-Toluenesulfonic acid monohydrate Catalysts: Bis[dichloro[η⁵-(pentamethylcyclopentadienyl)]rhodium] Solvents: Acetone; 15 min, rt 1.2 Reagents: Sodium azide; 16 h, 50 °C	Rhodium(III)-Catalyzed Azidation and Nitration of Arenes by C-H Activation By: Xie, Fang; et al Angewandte Chemie, International Edition (2013), 52(45), 11862-11866.
31-043-CAS-14905106 Steps: 1 Yield: 87% 1.1 Reagents: Iodobenzene diacetate, p-Toluenesulfonic acid monohydrate Catalysts: Chloro[(1,2,3,4,5-η)-1,2,3,4,5-pentamethyl-2,4-cyclopentadien-1-yl][2-(2-pyridinyl-κM)phenyl-κC]rhodium Solvents: Acetone; 15 min, rt 1.2 Reagents: Sodium azide; 16 h, 50 °C	Rhodium(III)-Catalyzed Azidation and Nitration of Arenes by C-H Activation By: Xie, Fang; et al Angewandte Chemie, International Edition (2013), 52(45), 11862-11866.

Scheme 21 (1 Reaction)


Steps: 1 Yield: 87%



31-043-CAS-9498219 Steps: 1 Yield: 87% 1.1 Reagents: Perchloric acid, Oxygen, Sodium azide Catalysts: Cupric perchlorate Solvents: Acetonitrile, Dichloromethane, Water; 20 h, rt Experimental Protocols	Designing a Cu(II)-ArCu(II)-ArCu(III)-Cu(I) Catalytic Cycle: Cu(II)-Catalyzed Oxidative Arene C-H Bond Azidation with Air as an Oxidant under Ambient Conditions By: Yao, Bo; et al Journal of Organic Chemistry (2014), 79(22), 11139-11145.
---	---

Scheme 22 (2 Reactions)

Steps: 1 Yield: 87%



Suppliers (99)


Supplier (1)

31-043-CAS-23585428 Steps: 1 Yield: 87% 1.1 Catalysts: Copper sulfate Solvents: Acetonitrile, Water; 15 min, 40 °C 1.2 Reagents: Sodium hydroxide, Sodium persulfate, Sodium azide Solvents: Water; 1.5 h, pH 4.5, 40 °C; 5 h, pH 4.5, 40 °C; 40 °C → 25 °C 1.3 Reagents: Sodium thiosulfate; 25 °C 1.4 Reagents: Hydrochloric acid Solvents: Water; pH 1 Experimental Protocols	Regioselective C-H Azidation of Anilines and Application to Synthesis of Key Intermediate for Pharmaceutical By: Seki, Masahiko; et al Journal of Organic Chemistry (2021), 86(11), 7842-7848.
---	--

31-043-CAS-21184659 Steps: 1 Yield: 87% 1.1 Reagents: Copper sulfate Solvents: Acetonitrile, Water; 15 min, 40 °C 1.2 Reagents: Sodium hydroxide Solvents: Water; pH 4.5, 40 °C 1.3 Reagents: Sodium persulfate, Sodium azide; 5 h, pH 4 - 4.5, 40 °C; 4 h, pH 4 - 4.5, 40 °C 1.4 Reagents: Sodium persulfate; 3 h, acidified, 40 °C 1.5 Reagents: Sodium hydroxide Solvents: Water; pH 4 - 4.5	Method for producing α-azidoaniline derivative or α, α'-diazidoaniline derivative as intermediate for α-aminoaniline and benzimidazole derivative by regioselective azidation of aniline derivative Assignee: Tokuyama Corporation World Intellectual Property Organization, WO2020004043 A1 2020-01-02 PatentPak available
--	--

Scheme 23 (1 Reaction)


Steps: 1 Yield: 86%



31-043-CAS-4744672 Steps: 1 Yield: 86% 1.1 Reagents: Oxygen, 1-Azido-1,2-benziodoxol-3(1<i>H</i>)-one Catalysts: Cupric acetate Solvents: Tetrahydrofuran; 12 h, 30 °C Experimental Protocols	Room-temperature Cu(II)-catalyzed aromatic C-H azidation for the synthesis of ortho-azido anilines with excellent regioselectivity By: Fan, Yunpeng; et al Chemical Communications (Cambridge, United Kingdom) (2014), 50(43), 5733-5736.
---	---

Scheme 24 (2 Reactions)


Steps: 1 Yield: 84-86%



31-043-CAS-12465304 Steps: 1 Yield: 86% 1.1 Reagents: Oxygen, 1-Azido-1,2-benziodoxol-3(1<i>H</i>)-one Catalysts: Cupric acetate Solvents: Tetrahydrofuran; 12 h, 20 °C Experimental Protocols	Room-temperature Cu(II)-catalyzed aromatic C-H azidation for the synthesis of ortho-azido anilines with excellent regioselectivity By: Fan, Yunpeng; et al Chemical Communications (Cambridge, United Kingdom) (2014), 50(43), 5733-5736.
31-043-CAS-17816764 Steps: 1 Yield: 84% 1.1 Reagents: Hydrogen peroxide, Oxygen, Sodium azide Catalysts: Cupric acetate Solvents: Water; 2 h, rt	Regioselective and Direct Azidation of Anilines via Cu(II)-Catalyzed C-H Functionalization in Water By: Fang, Hongli; et al Journal of Organic Chemistry (2017), 82(20), 11212-11217.

Scheme 25 (1 Reaction)

Steps: 1 Yield: 85%



Suppliers (17)

31-043-CAS-2006648

Steps: 1 Yield: 85%

Rhodium(III)-Catalyzed Azidation and Nitration of Arenes by C-H Activation


By: Xie, Fang; et al

Angewandte Chemie, International Edition (2013), 52(45), 11862-11866.

1.1 **Reagents:** Iodobenzene diacetate, p-Toluenesulfonic acid monohydrate**Catalysts:** Bis[dichloro[η⁵-(pentamethylcyclopentadienyl)]rhodium]**Solvents:** Acetone; 15 min, rt1.2 **Reagents:** Sodium azide; 16 h, 50 °C

Scheme 26 (1 Reaction)

Steps: 1 Yield: 85%



Suppliers (127)

Suppliers (32)

31-043-CAS-11131839

Steps: 1 Yield: 85%

Simple and efficient method for the preparation of aryl azides using sonication

By: Telvekar, Vikas N.; et al


Synthetic Communications (2012), 42(7), 1085-1089.

1.1 **Reagents:** Sodium azide, Iodate(1-), dichloro-, sodium**Solvents:** Water; 10 min, 25 °C

Experimental Protocols

Scheme 27 (2 Reactions)

Steps: 1 Yield: 84%




Suppliers (61)

Supplier (1)

31-043-CAS-23586532 Steps: 1 Yield: 84% 1.1 Catalysts: Copper sulfate Solvents: Acetonitrile, Water; 15 min, 40 °C 1.2 Reagents: Sodium hydroxide, Sodium persulfate, Sodium azide Solvents: Water; 1.5 h, pH 4.5, 40 °C; 3 h, pH 4.5, 40 °C; 40 °C → 25 °C 1.3 Reagents: Sodium thiosulfate; 25 °C 1.4 Reagents: Hydrochloric acid Solvents: Water; pH 1 Experimental Protocols	Regioselective C-H Azidation of Anilines and Application to Synthesis of Key Intermediate for Pharmaceutical By: Seki, Masahiko; et al Journal of Organic Chemistry (2021), 86(11), 7842-7848.
31-043-CAS-21184656 Steps: 1 Yield: 84% 1.1 Reagents: Copper sulfate Solvents: Acetonitrile, Water; 15 min, 40 °C 1.2 Reagents: Sodium hydroxide Solvents: Water; pH 4.5, 40 °C 1.3 Reagents: Sodium persulfate, Sodium azide; 1.5 h, pH 4 - 4.5, 40 °C; 4 h, pH 4 - 4.5, 40 °C 1.4 Reagents: Sodium persulfate; 3 h, acidified, 40 °C 1.5 Reagents: Sodium hydroxide Solvents: Water; pH 4 - 4.5	Method for producing α-azidoaniline derivative or α,α'-diazidoaniline derivative as intermediate for α-aminoaniline and benzimidazole derivative by regioselective azidation of aniline derivative Assignee: Tokuyama Corporation World Intellectual Property Organization, WO2020004043 A1 2020-01-02 PatentPak available

Scheme 28 (1 Reaction)

Steps: 1 Yield: 83%




Suppliers (85)

Suppliers (2)

31-043-CAS-13547433 Steps: 1 Yield: 83% 1.1 Reagents: Sodium azide, Iodate(1-), dichloro-, sodium Solvents: Water; 15 min, 25 °C Experimental Protocols	Simple and efficient method for the preparation of aryl azides using sonication By: Telvekar, Vikas N.; et al Synthetic Communications (2012), 42(7), 1085-1089.
--	--

Scheme 29 (1 Reaction)

Steps: 1 Yield: 83%




Suppliers (78)

Suppliers (16)

31-043-CAS-9003017	Steps: 1 Yield: 83%	Simple and efficient method for the preparation of aryl azides using sonication
1.1 Reagents: Sodium azide, Iodate(1-), dichloro-, sodium Solvents: Water; 10 min, 25 °C		By: Telvekar, Vikas N.; et al
Experimental Protocols		Synthetic Communications (2012), 42(7), 1085-1089.

Scheme 30 (1 Reaction)

Steps: 1 Yield: 83%




Suppliers (43)

31-043-CAS-5610742	Steps: 1 Yield: 83%	Rhodium(III)-Catalyzed Azidation and Nitration of Arenes by C-H Activation
1.1 Reagents: Iodobenzene diacetate, p-Toluenesulfonic acid monohydrate Catalysts: Bis[dichloro[η ⁵ -(pentamethylcyclopentadienyl)]rhodium] Solvents: Acetone; 15 min, rt		By: Xie, Fang; et al
1.2 Reagents: Sodium azide; 16 h, 50 °C		Angewandte Chemie, International Edition (2013), 52(45), 11862-11866.

Scheme 31 (1 Reaction)

Steps: 1 Yield: 82%




Suppliers (27)

31-043-CAS-19137067	Steps: 1 Yield: 82%	Mild Dirhodium(II)-Catalyzed Chemo- and Regioselective Azidation of Arenes
1.1 Reagents: 1-Azido-1,2-benziodoxol-3(1H)-one Catalysts: Rhodium, bis[μ-[α,α,α',α'-tetramethyl-1,3-benzenedipropionato(2-)-κO ¹ ,κO ³ :κO ³ ,κO ¹]]di-, (Rh-Rh) Solvents: Methanol; 10 h, 50 °C		By: Wang, Yi; et al
Experimental Protocols		Organic Letters (2018), 20(18), 5732-5736.

Scheme 32 (1 Reaction)

Steps: 1 Yield: 82%




Suppliers (69)

31-043-CAS-4994595 Steps: 1 Yield: 82% 1.1 Reagents: Iodobenzene diacetate, p-Toluenesulfonic acid monohydrate Catalysts: Bis[dichloro[η⁵-(pentamethylcyclopentadienyl)]rhodium] Solvents: Acetone; 15 min, rt 1.2 Reagents: Sodium azide; 16 h, 50 °C	Rhodium(III)-Catalyzed Azidation and Nitration of Arenes by C-H Activation By: Xie, Fang; et al Angewandte Chemie, International Edition (2013), 52(45), 11862-11866.
---	---

Scheme 33 (1 Reaction)


Steps: 1 Yield: 82%



31-043-CAS-15076086 Steps: 1 Yield: 82% 1.1 Reagents: Sodium azide, Iodate(1-), dichloro-, sodium Solvents: Water; 10 min, 25 °C Experimental Protocols	Simple and efficient method for the preparation of aryl azides using sonication By: Telvekar, Vikas N.; et al Synthetic Communications (2012), 42(7), 1085-1089.
--	--

Scheme 34 (1 Reaction)


Steps: 1 Yield: 82%



31-043-CAS-181533 Steps: 1 Yield: 82% 1.1 Reagents: Sodium azide, Iodate(1-), dichloro-, sodium Solvents: Water; 15 min, 25 °C Experimental Protocols	Simple and efficient method for the preparation of aryl azides using sonication By: Telvekar, Vikas N.; et al Synthetic Communications (2012), 42(7), 1085-1089.
--	--


Scheme 35 (1 Reaction)

Steps: 1 Yield: 82%



31-043-CAS-7897434	Steps: 1 Yield: 82%	Rhodium(III)-Catalyzed Azidation and Nitration of Arenes by C-H Activation
1.1 Reagents: Iodobenzene diacetate, p-Toluenesulfonic acid monohydrate Catalysts: Bis[dichloro[η ⁵ -(pentamethylcyclopentadienyl)]rhodium] Solvents: Acetone; 15 min, rt		By: Xie, Fang; et al Angewandte Chemie, International Edition (2013), 52(45), 11862-11866.
1.2 Reagents: Sodium azide; 16 h, 50 °C		


Scheme 36 (1 Reaction)

Steps: **1** Yield: **81%**

Suppliers (55)

31-043-CAS-10034634	Steps: 1 Yield: 81%	Rhodium(III)-Catalyzed Azidation and Nitration of Arenes by C-H Activation
1.1 Reagents: Iodobenzene diacetate, p-Toluenesulfonic acid monohydrate Catalysts: Bis[dichloro[η ⁵ -(pentamethylcyclopentadienyl)]rhodium] Solvents: Acetone; 15 min, rt		By: Xie, Fang; et al Angewandte Chemie, International Edition (2013), 52(45), 11862-11866.
1.2 Reagents: Sodium azide; 16 h, 50 °C		


Scheme 37 (1 Reaction)

Steps: **1** Yield: **80%**

Suppliers (6)

31-043-CAS-2822299	Steps: 1 Yield: 80%	Rhodium(III)-Catalyzed Azidation and Nitration of Arenes by C-H Activation
1.1 Reagents: Iodobenzene diacetate, p-Toluenesulfonic acid monohydrate Catalysts: Bis[dichloro[η ⁵ -(pentamethylcyclopentadienyl)]rhodium] Solvents: Acetone; 15 min, rt		By: Xie, Fang; et al Angewandte Chemie, International Edition (2013), 52(45), 11862-11866.
1.2 Reagents: Sodium azide; 16 h, 50 °C		

Scheme 38 (1 Reaction)

Steps: **1** Yield: **80%**

Suppliers (58)


31-043-CAS-11620599 Steps: 1 Yield: 80% 1.1 Reagents: Iodobenzene diacetate, <i>p</i>-Toluenesulfonic acid monohydrate Catalysts: Bis[dichloro[η⁵-(pentamethylcyclopentadienyl)]rhodium] Solvents: Acetone; 15 min, rt 1.2 Reagents: Sodium azide; 16 h, 50 °C	Rhodium(III)-Catalyzed Azidation and Nitration of Arenes by C-H Activation By: Xie, Fang; et al Angewandte Chemie, International Edition (2013), 52(45), 11862-11866.
---	---

Scheme 39 (1 Reaction)

Steps: **1** Yield: **80%**


31-043-CAS-13758740 Steps: 1 Yield: 80% 1.1 Reagents: <i>p</i>-Toluenesulfonic acid, Oxygen, Sodium azide Catalysts: Cupric perchlorate Solvents: Acetonitrile, Dichloromethane, Water; 20 h, rt Experimental Protocols	Designing a Cu(II)-ArCu(II)-ArCu(III)-Cu(I) Catalytic Cycle: Cu(II)-Catalyzed Oxidative Arene C-H Bond Azidation with Air as an Oxidant under Ambient Conditions By: Yao, Bo; et al Journal of Organic Chemistry (2014), 79(22), 11139-11145.
--	---

Scheme 40 (1 Reaction)

Steps: **1** Yield: **80%**

31-043-CAS-9121825 Steps: 1 Yield: 80% 1.1 Reagents: Oxygen, 1-Azido-1,2-benziodoxol-3(1<i>H</i>)-one Catalysts: Cupric acetate Solvents: Tetrahydrofuran; 12 h, 20 °C Experimental Protocols	Room-temperature Cu(II)-catalyzed aromatic C-H azidation for the synthesis of ortho-azido anilines with excellent regioselectivity By: Fan, Yunpeng; et al Chemical Communications (Cambridge, United Kingdom) (2014), 50(43), 5733-5736.
--	---


Scheme 41 (1 Reaction)

Steps: **1** Yield: **80%**

31-043-CAS-6002384 Steps: 1 Yield: 80% 1.1 Reagents: Iodobenzene diacetate, p-Toluenesulfonic acid monohydrate Catalysts: Bis[dichloro[η⁵-(pentamethylcyclopentadienyl)]rhodium] Solvents: Acetone; 15 min, rt 1.2 Reagents: Sodium azide; 16 h, 50 °C	Rhodium(III)-Catalyzed Azidation and Nitration of Arenes by C-H Activation By: Xie, Fang; et al Angewandte Chemie, International Edition (2013), 52(45), 11862-11866.
---	---

Scheme 42 (1 Reaction)


Steps: 1 Yield: 80%



31-043-CAS-17816768 Steps: 1 Yield: 80% 1.1 Reagents: Hydrogen peroxide, Oxygen, Sodium azide Catalysts: Cupric acetate Solvents: Water; 2 h, rt	Regioselective and Direct Azidation of Anilines via Cu(II)-Catalyzed C-H Functionalization in Water By: Fang, Hongli; et al Journal of Organic Chemistry (2017), 82(20), 11212-11217.
---	---

Scheme 43 (2 Reactions)

Steps: 1 Yield: 79%




31-043-CAS-23587647 Steps: 1 Yield: 79% 1.1 Catalysts: Copper sulfate Solvents: Acetonitrile, Water; 15 min, 40 °C 1.2 Reagents: Sodium hydroxide, Sodium persulfate, Sodium azide Solvents: Water; 1.5 h, pH 4.5, 40 °C; 2.5 h, pH 4.5, 40 °C; 40 °C → 25 °C 1.3 Reagents: Sodium thiosulfate; 25 °C 1.4 Reagents: Hydrochloric acid Solvents: Water; pH 1 Experimental Protocols	Regioselective C-H Azidation of Anilines and Application to Synthesis of Key Intermediate for Pharmaceutical By: Seki, Masahiko; et al Journal of Organic Chemistry (2021), 86(11), 7842-7848.
---	--

31-043-CAS-21184660 Steps: 1 Yield: 79% 1.1 Reagents: Copper sulfate Solvents: Acetonitrile, Water; 15 min, 40 °C 1.2 Reagents: Sodium hydroxide Solvents: Water; pH 4.5, 40 °C 1.3 Reagents: Sodium persulfate, Sodium azide; 2.5 h, pH 4 - 4.5, 40 °C; 4 h, pH 4 - 4.5, 40 °C 1.4 Reagents: Sodium persulfate; 3 h, acidified, 40 °C 1.5 Reagents: Sodium hydroxide Solvents: Water; pH 4 - 4.5	Method for producing α-azidoaniline derivative or α,α'-diazidoaniline derivative as intermediate for α-aminoaniline and benzimidazole derivative by regioselective azidation of aniline derivative Assignee: Tokuyama Corporation World Intellectual Property Organization, WO2020004043 A1 2020-01-02 PatentPak available
---	---

Scheme 44 (1 Reaction)

Steps: 1 Yield: 79%



Suppliers (60)

Supplier (1)

31-043-CAS-12659292 Steps: 1 Yield: 79% 1.1 Reagents: Iodobenzene diacetate, p-Toluenesulfonic acid monohydrate Catalysts: Bis[dichloro[η^5-(pentamethylcyclopentadienyl)]rhodium] Solvents: Acetone; 15 min, rt 1.2 Reagents: Sodium azide; 16 h, 50 °C	Rhodium(III)-Catalyzed Azidation and Nitration of Arenes by C-H Activation By: Xie, Fang; et al Angewandte Chemie, International Edition (2013), 52(45), 11862-11866.
---	--