

Initiating Search

November 10, 2025, 8:55 PM

• Search:

CAS Lexicon Search:

Thiol-ene reaction - Preferred Concept

Search Tasks

Task	Result Type	View
Returned Reference Results from CAS Lexicon (2,485)	 References	View Results
Exported: Retrieved Related Reaction Results + Filters (566)	 Reactions	View Results
Filtered By:		
Yield:	90-100%, 80-89%, 70-79%	
Document Type:	Journal	
Language:	English	

Copyright © 2025 American Chemical Society (ACS). All Rights Reserved.


Internal use only. Redistribution is subject to the terms of your CAS SciFinder License Agreement and CAS information Use Policies.

Reactions (50)

[View in CAS SciFinder](#)

Scheme 1 (1 Reaction)

Steps: 1 Yield: 100%



31-238-CAS-17059372

Steps: 1 Yield: 100%

Thioether-functionalized picolinium ionic liquids: synthesis, physical properties and computational studies

By: Thigpen, A. Shay; et al


New Journal of Chemistry (2017), 41(4), 1625-1630.

 1.1 Catalysts: 2,2-Dimethoxy-2-phenylacetophenone
 Solvents: Methanol, Dichloromethane; 8 h, rt

Experimental Protocols

Scheme 2 (1 Reaction)

Steps: 1 Yield: 100%



31-238-CAS-13675340

Steps: 1 Yield: 100%

Orthogonal reactivity of thiols toward chlorovinylsilanes: selective thiol-ene chemistry

By: Jennings, Abby R.; et al


Tetrahedron Letters (2014), 55(50), 6773-6775.

 1.1 Catalysts: Benzophenone
 Solvents: Diethyl ether; 4 h

Experimental Protocols

Scheme 3 (1 Reaction)

Steps: 1 Yield: 100%



31-238-CAS-17059373

Steps: 1 Yield: 100%

Thioether-functionalized picolinium ionic liquids: synthesis, physical properties and computational studies

By: Thigpen, A. Shay; et al




New Journal of Chemistry (2017), 41(4), 1625-1630.

 1.1 Catalysts: 2,2-Dimethoxy-2-phenylacetophenone
 Solvents: Methanol, Dichloromethane; 8 h, rt

Experimental Protocols

Scheme 4 (1 Reaction)

Steps: 1 Yield: 100%



Suppliers (56)

Suppliers (60)

31-238-CAS-15671847

Steps: 1 Yield: 100%

Highly thermally resistant polyhedral oligomeric silsesquioxanes lubricating oil prepared via a thiol-ene click reaction

By: Wang, Xiaobai; et al


Science of Advanced Materials (2014), 6(7), 1553-1561.

1.1 Catalysts: Azobisisobutyronitrile; rt → 80 °C; 5 h, 80 °C

Scheme 5 (1 Reaction)

Steps: 1 Yield: 100%

Multi-component
structure image
available in CAS
SciFinder



Multi-component
structure image
available in CAS
SciFinder

Suppliers (63)

31-238-CAS-17059359

Steps: 1 Yield: 100%

Thioether-functionalized picolinium ionic liquids: synthesis, physical properties and computational studies

By: Thigpen, A. Shay; et al

New Journal of Chemistry (2017), 41(4), 1625-1630.


1.1 Catalysts: 2,2-Dimethoxy-2-phenylacetophenone
Solvents: Methanol, Dichloromethane; 8 h, rt

Experimental Protocols

Scheme 6 (1 Reaction)

Steps: 1 Yield: 100%

Multi-component
structure image
available in CAS
SciFinder



Multi-component
structure image
available in CAS
SciFinder

Suppliers (56)

31-238-CAS-17059365

Steps: 1 Yield: 100%

Thioether-functionalized picolinium ionic liquids: synthesis, physical properties and computational studies

By: Thigpen, A. Shay; et al


New Journal of Chemistry (2017), 41(4), 1625-1630.

1.1 Catalysts: 2,2-Dimethoxy-2-phenylacetophenone
Solvents: Methanol, Dichloromethane; 8 h, rt

Experimental Protocols

Scheme 7 (1 Reaction)

Steps: 1 Yield: 100%



Suppliers (57)

31-238-CAS-19408298

Steps: 1 Yield: 100%

Post-polymerization modification of the side chain in optically active polymers by thiol-ene reaction

1.1 **Catalysts:** 2,2-Dimethoxy-2-phenylacetophenone
Solvents: Toluene; 16 h, rt


By: Kanbayashi, Naoya; et al

Polymer Chemistry (2017), 8(6), 985-994.

Scheme 8 (1 Reaction)

Steps: 1 Yield: 100%

Multi-component
structure image
available in CAS
SciFinder



Multi-component
structure image
available in CAS
SciFinder

Suppliers (51)

31-238-CAS-17059358

Steps: 1 Yield: 100%

Thioether-functionalized picolinium ionic liquids: synthesis, physical properties and computational studies

1.1 **Catalysts:** 2,2-Dimethoxy-2-phenylacetophenone
Solvents: Methanol, Dichloromethane; 8 h, rt

By: Thigpen, A. Shay; et al


New Journal of Chemistry (2017), 41(4), 1625-1630.

Experimental Protocols

Scheme 9 (1 Reaction)

Steps: 1 Yield: 100%

Multi-component
structure image
available in CAS
SciFinder



Multi-component
structure image
available in CAS
SciFinder

Suppliers (48)

31-238-CAS-17059362

Steps: 1 Yield: 100%

Thioether-functionalized picolinium ionic liquids: synthesis, physical properties and computational studies

1.1 **Catalysts:** 2,2-Dimethoxy-2-phenylacetophenone
Solvents: Methanol, Dichloromethane; 8 h, rt

By: Thigpen, A. Shay; et al


New Journal of Chemistry (2017), 41(4), 1625-1630.

Experimental Protocols

Scheme 10 (1 Reaction)

Steps: 1 Yield: 100%

Multi-component
structure image
available in CAS
SciFinder



Suppliers (60)

Multi-component
structure image
available in CAS
SciFinder

31-238-CAS-17059366

Steps: 1 Yield: 100%

Thioether-functionalized picolinium ionic liquids: synthesis, physical properties and computational studies

By: Thigpen, A. Shay; et al

New Journal of Chemistry (2017), 41(4), 1625-1630.


1.1 **Catalysts:** 2,2-Dimethoxy-2-phenylacetophenone
Solvents: Methanol, Dichloromethane; 8 h, rt

Experimental Protocols

Scheme 11 (1 Reaction)

Steps: 1 Yield: 100%

Multi-component
structure image
available in CAS
SciFinder



Suppliers (57)

Multi-component
structure image
available in CAS
SciFinder

31-238-CAS-10441578

Steps: 1 Yield: 100%

Synthesis of New Lipid-Inspired Ionic Liquids by Thiol-ene Chemistry: Profound Solvent Effect on Reaction Pathway

By: Mirjafari, Arsalan; et al


Chemistry - A European Journal (2014), 20(25), 7576-7580.

1.1 **Catalysts:** 2,2-Dimethoxy-2-phenylacetophenone
Solvents: Methanol; 8 h, rt


Experimental Protocols

Scheme 12 (1 Reaction)


Steps: 1 Yield: 100%



Suppliers (56)



Suppliers (60)



31-238-CAS-18889801

Steps: 1 Yield: 100%

Highly efficient preparation of hyperbranched epoxy resins by UV-initiated thiol-ene click reaction

By: Zhang, Daohong; et al

Progress in Organic Coatings (2016), 101, 178-185.

1.1 **Catalysts:** 1-Hydroxycyclohexyl phenyl ketone

Experimental Protocols

Steps: **1** Yield: **100%**



New Journal of Chemistry (2017), 41(4), 1625-1630.

Steps: 1 Yield: 100%



 Suppliers (60)

Progress in Organic Coatings (2016), 101, 178-185.




Steps: 1 Yield: 100%



New Journal of Chemistry (2017), 41(4), 1625-1630.

Scheme 16 (1 Reaction)

Steps: 1 Yield: 100%



Suppliers (56)

Suppliers (57)

31-238-CAS-15632588

Steps: 1 Yield: 100%

Highly thermally resistant polyhedral oligomeric silsesquioxanes lubricating oil prepared via a thiol-ene click reaction

By: Wang, Xiaobai; et al


Science of Advanced Materials (2014), 6(7), 1553-1561.

1.1 Catalysts: Azobisisobutyronitrile; rt → 80 °C; 5 h, 80 °C

Scheme 17 (1 Reaction)

Steps: 1 Yield: 100%

Multi-component
structure image
available in CAS
SciFinder



Multi-component
structure image
available in CAS
SciFinder

Suppliers (48)

31-238-CAS-17059363

Steps: 1 Yield: 100%

Thioether-functionalized picolinium ionic liquids: synthesis, physical properties and computational studies

By: Thigpen, A. Shay; et al

New Journal of Chemistry (2017), 41(4), 1625-1630.


1.1 Catalysts: 2,2-Dimethoxy-2-phenylacetophenone
Solvents: Methanol, Dichloromethane; 8 h, rt

Experimental Protocols

Scheme 18 (1 Reaction)

Steps: 1 Yield: 100%

Multi-component
structure image
available in CAS
SciFinder



Multi-component
structure image
available in CAS
SciFinder

Suppliers (44)

31-238-CAS-17059368

Steps: 1 Yield: 100%

Thioether-functionalized picolinium ionic liquids: synthesis, physical properties and computational studies

By: Thigpen, A. Shay; et al


New Journal of Chemistry (2017), 41(4), 1625-1630.

1.1 Catalysts: 2,2-Dimethoxy-2-phenylacetophenone
Solvents: Methanol, Dichloromethane; 8 h, rt

Experimental Protocols

Scheme 19 (1 Reaction)

Steps: 1 Yield: 100%



Absolute stereochemistry shown,
Rotation (+)

Suppliers (108)

Suppliers (103)

31-614-CAS-33027508

Steps: 1 Yield: 100%

1.1 **Catalysts:** Bis(2,4,6-trimethylbenzoyl)phenylphosphine oxide
Solvents: Ethanol; 8 h

Experimental Protocols


Synthesis of a Curing Agent Derived from Limonene and the Study of Its Performance to Polymerize a Biobased Epoxy Resin Using the Epoxy/Thiol-Ene Photopolymerization Technique

By: Acosta Ortiz, Ricardo; et al

Polymers (Basel, Switzerland) (2022), 14(11), 2192.

Scheme 20 (1 Reaction)

Steps: 1 Yield: 100%



Suppliers (38)

Suppliers (49)

Suppliers (2)

31-238-CAS-7312641

Steps: 1 Yield: 100%

1.1 **Catalysts:** Benzophenone; 4 h

Experimental Protocols


Orthogonal reactivity of thiols toward chlorovinylsilanes: selective thiol-ene chemistry

By: Jennings, Abby R.; et al

Tetrahedron Letters (2014), 55(50), 6773-6775.

Scheme 21 (1 Reaction)

Steps: 1 Yield: 100%



Suppliers (119)

31-614-CAS-37959237

Steps: 1 Yield: 100%

1.1 **Catalysts:** Azobisisobutyronitrile
Solvents: Dichloromethane; 24 - 46 h, reflux

Experimental Protocols


Systematic Order-Made Synthesis of Sequence-Defined Polyurethanes with Length, Types, and Topologies

By: Yamauchi, Haruka; et al

ACS Macro Letters (2023), 12(9), 1264-1271.

Scheme 22 (1 Reaction)

Steps: 1 Yield: 100%



Suppliers (180)

31-238-CAS-18623494

Steps: 1 Yield: 100%

A double-click approach to the protecting group free synthesis of glycoconjugates

By: Alexander, S. R.; et al

Organic & Biomolecular Chemistry (2018), 16(8), 1258-1262.


1.1 Reagents: 2,2-Dimethoxy-2-phenylacetophenone

Solvents: *tert*-Butanol, Water; 4 h, pH 4

Experimental Protocols

Scheme 23 (1 Reaction)

Steps: 1 Yield: 100%



Suppliers (66)

Suppliers (60)

31-238-CAS-18889795

Steps: 1 Yield: 100%

Highly efficient preparation of hyperbranched epoxy resins by UV-initiated thiol-ene click reaction

By: Zhang, Daohong; et al


Progress in Organic Coatings (2016), 101, 178-185.

1.1 Catalysts: 1-Hydroxycyclohexyl phenyl ketone; 5 min, rt

Experimental Protocols

Scheme 24 (1 Reaction)

Steps: 1 Yield: 100%



Suppliers (63)

Suppliers (79)

31-238-CAS-23565505

Steps: 1 Yield: 100%

UV-Resistant Trifluoropropyl-Substituted Open-Cage Silsesquioxane-Pendant Polysiloxanes

By: Li, Lina; et al

ACS Applied Polymer Materials (2021), 3(3), 1368-1375.


1.1 Catalysts: Azobisisobutyronitrile

Solvents: Tetrahydrofuran; 48 h, 50 °C

Experimental Protocols

Scheme 25 (1 Reaction)

Steps: 1 Yield: 100%



Suppliers (60)

31-238-CAS-18889796

Steps: 1 Yield: 100%

Highly efficient preparation of hyperbranched epoxy resins by UV-initiated thiol-ene click reaction

1.1 Catalysts: 1-Hydroxycyclohexyl phenyl ketone; 5 min, rt


Experimental Protocols

By: Zhang, Daohong; et al

Progress in Organic Coatings (2016), 101, 178-185.

Scheme 26 (1 Reaction)

Steps: 1 Yield: 100%

 Multi-component
structure image
available in CAS
SciFinder

 Multi-component
structure image
available in CAS
SciFinder

Suppliers (57)

31-238-CAS-17059369

Steps: 1 Yield: 100%

Thioether-functionalized picolinium ionic liquids: synthesis, physical properties and computational studies
1.1 Catalysts: 2,2-Dimethoxy-2-phenylacetophenone
Solvents: Methanol, Dichloromethane; 8 h, rt


Experimental Protocols

By: Thigpen, A. Shay; et al


New Journal of Chemistry (2017), 41(4), 1625-1630.

Scheme 27 (1 Reaction)


Steps: 1 Yield: 100%



Suppliers (56)



Suppliers (67)



31-238-CAS-15571183

Steps: 1 Yield: 100%

Highly thermally resistant polyhedral oligomeric silsesquioxanes lubricating oil prepared via a thiol-ene click reaction


1.1 Catalysts: Azobisisobutyronitrile; rt → 80 °C; 5 h, 80 °C

By: Wang, Xiaobai; et al

Science of Advanced Materials (2014), 6(7), 1553-1561.

Scheme 28 (1 Reaction)

Steps: 1 Yield: 100%



Suppliers (8)

Suppliers (60)

31-238-CAS-18889791

Steps: 1 Yield: 100%

Highly efficient preparation of hyperbranched epoxy resins by UV-initiated thiol-ene click reaction

1.1 Catalysts: 1-Hydroxycyclohexyl phenyl ketone; 5 min, rt


By: Zhang, Daohong; et al

Experimental Protocols

Progress in Organic Coatings (2016), 101, 178-185.

Scheme 29 (1 Reaction)

Steps: 1 Yield: 100%



Suppliers (40)

31-238-CAS-4253724

Steps: 1 Yield: 100%

Orthogonal reactivity of thiols toward chlorovinylsilanes: selective thiol-ene chemistry

1.1 Catalysts: Benzophenone
Solvents: Diethyl ether; 4 h


By: Jennings, Abby R.; et al

Experimental Protocols

Tetrahedron Letters (2014), 55(50), 6773-6775.

Scheme 30 (1 Reaction)

Steps: 1 Yield: 100%



Suppliers (43)

31-238-CAS-17059360

Steps: 1 Yield: 100%

Thioether-functionalized picolinium ionic liquids: synthesis, physical properties and computational studies

1.1 Catalysts: 2,2-Dimethoxy-2-phenylacetophenone
Solvents: Methanol, Dichloromethane; 8 h, rt


By: Thigpen, A. Shay; et al

Experimental Protocols

New Journal of Chemistry (2017), 41(4), 1625-1630.

Scheme 31 (1 Reaction)

Steps: 1 Yield: 100%



Supplier (1)

Suppliers (57)

31-238-CAS-15584888

Steps: 1 Yield: 100%

Highly thermally resistant polyhedral oligomeric silsesquioxanes lubricating oil prepared via a thiol-ene click reaction


By: Wang, Xiaobai; et al

Science of Advanced Materials (2014), 6(7), 1553-1561.

1.1 Catalysts: Azobisisobutyronitrile; rt → 80 °C; 5 h, 80 °C

Scheme 32 (1 Reaction)

Steps: 1 Yield: 100%



Suppliers (119)

31-614-CAS-37959251

Steps: 1 Yield: 100%

Systematic Order-Made Synthesis of Sequence-Defined Polyurethanes with Length, Types, and Topologies

By: Yamauchi, Haruka; et al


ACS Macro Letters (2023), 12(9), 1264-1271.

1.1 Catalysts: Azobisisobutyronitrile
Solvents: Dichloromethane; 2 d, 45 °C

Experimental Protocols

Scheme 33 (1 Reaction)

Steps: 1 Yield: 100%



Suppliers (51)

31-238-CAS-11870006

Steps: 1 Yield: 100%

Facile and Efficient Synthesis of Carbosiloxane Dendrimers via Orthogonal Click Chemistry Between Thiol and Ene

By: Zhang, Zhida; et al


Macromolecular Rapid Communications (2016), 37(4), 318-322.

1.1 Catalysts: 2,2-Dimethoxy-2-phenylacetophenone
Solvents: Tetrahydrofuran; 10 min, rt

Experimental Protocols

Scheme 34 (1 Reaction)

Steps: 1 Yield: 100%



31-238-CAS-17059371

Steps: 1 Yield: 100%

Thioether-functionalized picolinium ionic liquids: synthesis, physical properties and computational studies

1.1 Catalysts: 2,2-Dimethoxy-2-phenylacetophenone
Solvents: Methanol, Dichloromethane; 8 h, rt


By: Thigpen, A. Shay; et al

Experimental Protocols

New Journal of Chemistry (2017), 41(4), 1625-1630.

Scheme 35 (1 Reaction)

Steps: 1 Yield: 100%



31-614-CAS-33648654

Steps: 1 Yield: 100%

Synthesis of Thiomorpholine via a Telescoped Photochemical Thiol-Ene/Cyclization Sequence in Continuous Flow

1.1 Catalysts: Fluorenone
Solvents: Methanol; 30 min, 20 °C


By: Steiner, Alexander; et al

Experimental Protocols

Organic Process Research & Development (2022), 26(8), 2532-2539.

Scheme 36 (1 Reaction)

Steps: 1 Yield: 100%



31-614-CAS-37959228

Steps: 1 Yield: 100%

Systematic Order-Made Synthesis of Sequence-Defined Polyurethanes with Length, Types, and Topologies

1.1 Catalysts: Azobisisobutyronitrile
Solvents: Dichloromethane; 1 - 2 d, 45 °C

By: Yamauchi, Haruka; et al


Experimental Protocols

ACS Macro Letters (2023), 12(9), 1264-1271.

Scheme 37 (1 Reaction)

Steps: 1 Yield: 100%

Multi-component
structure image
available in CAS
SciFinder



Multi-component
structure image
available in CAS
SciFinder

Suppliers (56)

31-238-CAS-17059364

Steps: 1 Yield: 100%

Thioether-functionalized picolinium ionic liquids: synthesis, physical properties and computational studies

By: Thigpen, A. Shay; et al

New Journal of Chemistry (2017), 41(4), 1625-1630.


1.1 **Catalysts:** 2,2-Dimethoxy-2-phenylacetophenone
Solvents: Methanol, Dichloromethane; 8 h, rt

Experimental Protocols

Scheme 38 (1 Reaction)

Steps: 1 Yield: 100%

Multi-component
structure image
available in CAS
SciFinder



Multi-component
structure image
available in CAS
SciFinder

Suppliers (60)

31-238-CAS-17059367

Steps: 1 Yield: 100%

Thioether-functionalized picolinium ionic liquids: synthesis, physical properties and computational studies

By: Thigpen, A. Shay; et al


New Journal of Chemistry (2017), 41(4), 1625-1630.

1.1 **Catalysts:** 2,2-Dimethoxy-2-phenylacetophenone
Solvents: Methanol, Dichloromethane; 8 h, rt

Experimental Protocols


Scheme 39 (1 Reaction)

Steps: 1 Yield: 100%




Double bond geometry shown

+



→



Suppliers (62)

Suppliers (5)

31-614-CAS-31181065

Steps: 1 Yield: 100%

Salen-decorated Periodic Mesoporous Organosilica: From Metal-assisted Epoxidation to Metal-free CO₂ Insertion

By: Haghighat, Mahdiah; et al


Chemistry - An Asian Journal (2021), 16(15), 2126-2135.

1.1 **Reagents:** 2,2-Dimethoxy-2-phenylacetophenone

Experimental Protocols

Scheme 40 (1 Reaction)

Steps: 1 Yield: 100%



Suppliers (58)

Suppliers (46)

Supplier (1)

31-238-CAS-20219323

Steps: 1 Yield: 100%

Visible-Light-Mediated Organocatalyzed Thiol-Ene Reaction Initiated by a Proton-Coupled Electron Transfer


1.1 Catalysts: 7*H*-Benzimidazo[2,1-*a*]benz[*de*]isoquinolin-7-one
Solvents: Cyclohexane; 1 h, rt

By: Levin, Vitalij V.; et al

Journal of Organic Chemistry (2019), 84(12), 8337-8343.

Scheme 41 (1 Reaction)

Steps: 1 Yield: 100%



Suppliers (56)

Suppliers (60)

31-238-CAS-15551743

Steps: 1 Yield: 100%

Highly thermally resistant polyhedral oligomeric silsesquioxanes lubricating oil prepared via a thiol-ene click reaction


1.1 Catalysts: Azobisisobutyronitrile; rt → 80 °C; 5 h, 80 °C

By: Wang, Xiaobai; et al

Science of Advanced Materials (2014), 6(7), 1553-1561.

Scheme 42 (1 Reaction)

Steps: 1 Yield: 100%



Suppliers (80)

Suppliers (119)

31-614-CAS-42491586

Steps: 1 Yield: 100%

Synthesis of Novel Eugenol-Containing Polysilsesquioxanes with a Flexible Spacer and Their Use for Functional Anticorrosive Coatings

1.1 Catalysts: 2-Hydroxy-2-methyl-1-phenylpropan-1-one; 24 h, 35 °C


By: Ageenkov, Alexander D.; et al

ChemistrySelect (2024), 9(37), e202401623.

Experimental Protocols

Scheme 43 (1 Reaction)

Steps: 1 Yield: 100%



Suppliers (56)

Suppliers (56)

31-238-CAS-15591296

Steps: 1 Yield: 100%

Highly thermally resistant polyhedral oligomeric silsesquioxanes lubricating oil prepared via a thiol-ene click reaction


By: Wang, Xiaobai; et al

Science of Advanced Materials (2014), 6(7), 1553-1561.

1.1 Catalysts: Azobisisobutyronitrile; rt → 80 °C; 5 h, 80 °C

Scheme 44 (1 Reaction)

Steps: 1 Yield: 100%



Suppliers (46)

Suppliers (62)

Suppliers (2)

31-238-CAS-20219324

Steps: 1 Yield: 100%

Visible-Light-Mediated Organocatalyzed Thiol-Ene Reaction Initiated by a Proton-Coupled Electron Transfer


By: Levin, Vitalij V.; et al

Journal of Organic Chemistry (2019), 84(12), 8337-8343.

1.1 Catalysts: 7*H*-Benzimidazo[2,1-*a*]benz[*de*]isoquinolin-7-one
Solvents: Cyclohexane; 1 h, rt

Scheme 45 (1 Reaction)

Steps: 1 Yield: 100%



Suppliers (58)

31-238-CAS-13398185

Steps: 1 Yield: 100%

Orthogonal reactivity of thiols toward chlorovinylsilanes: selective thiol-ene chemistry

By: Jennings, Abby R.; et al

Tetrahedron Letters (2014), 55(50), 6773-6775.


1.1 Catalysts: Benzophenone
Solvents: Diethyl ether; 4 h

Experimental Protocols

Scheme 46 (1 Reaction)

Steps: 1 Yield: 100%

Multi-component
structure image
available in CAS
SciFinder



Multi-component
structure image
available in CAS
SciFinder

Suppliers (57)

31-238-CAS-17059370

Steps: 1 Yield: 100%

Thioether-functionalized picolinium ionic liquids: synthesis, physical properties and computational studies

By: Thigpen, A. Shay; et al




New Journal of Chemistry (2017), 41(4), 1625-1630.

1.1 Catalysts: 2,2-Dimethoxy-2-phenylacetophenone
Solvents: Methanol, Dichloromethane; 8 h, rt

Experimental Protocols

Scheme 47 (1 Reaction)

Steps: 1 Yield: 100%



Supplier (1)

Suppliers (60)

31-238-CAS-15649362

Steps: 1 Yield: 100%

Highly thermally resistant polyhedral oligomeric silsesquioxanes lubricating oil prepared via a thiol-ene click reaction


By: Wang, Xiaobai; et al

Science of Advanced Materials (2014), 6(7), 1553-1561.

1.1 Catalysts: Azobisisobutyronitrile; rt → 80 °C; 5 h, 80 °C

Scheme 48 (1 Reaction)

Steps: 1 Yield: 99%



Suppliers (78)

Suppliers (53)

Suppliers (3)

31-238-CAS-20219331

Steps: 1 Yield: 99%

Visible-Light-Mediated Organocatalyzed Thiol-Ene Reaction Initiated by a Proton-Coupled Electron Transfer


By: Levin, Vitalij V.; et al

Journal of Organic Chemistry (2019), 84(12), 8337-8343.

1.1 Catalysts: Acridine Orange base
Solvents: Cyclohexane; 1 h, rt

Scheme 49 (1 Reaction)

Steps: 1 Yield: 99%



Suppliers (46)

Suppliers (53)

31-238-CAS-20219322

Steps: 1 Yield: 99%

Visible-Light-Mediated Organocatalyzed Thiol-Ene Reaction Initiated by a Proton-Coupled Electron Transfer


1.1 **Catalysts:** Acridine Orange base
Solvents: Cyclohexane; 1 h, rt

By: Levin, Vitalij V.; et al

Journal of Organic Chemistry (2019), 84(12), 8337-8343.

Scheme 50 (1 Reaction)

Steps: 1 Yield: 99%



Suppliers (56)

Suppliers (115)

Suppliers (3)

31-238-CAS-20219341

Steps: 1 Yield: 99%

Visible-Light-Mediated Organocatalyzed Thiol-Ene Reaction Initiated by a Proton-Coupled Electron Transfer

1.1 **Catalysts:** 7-*H*-Benzimidazo[2,1-*a*]benz[*de*]isoquinolin-7-one
Solvents: Cyclohexane; 1 h, rt

By: Levin, Vitalij V.; et al

Journal of Organic Chemistry (2019), 84(12), 8337-8343.