

Initiating Search

November 4, 2025, 2:38 PM

• Search:

CAS Lexicon Search:

Wittig reaction - Preferred Concept

Search Tasks

Task	Result Type	View
Returned Reference Results from CAS Lexicon (7,049)	 References	View Results
Exported: Retrieved Related Reaction Results + Filters (1,688)	 Reactions	View Results
Filtered By:		
Yield:	90-100%, 80-89%, 70-79%	
Language:	English	
Publication Year:	2015 to 2025	

Copyright © 2025 American Chemical Society (ACS). All Rights Reserved.


Internal use only. Redistribution is subject to the terms of your CAS SciFinder License Agreement and CAS information Use Policies.

Reactions (50)

[View in CAS SciFinder](#)

Scheme 1 (1 Reaction)

Steps: 1 Yield: 100%



31-339-CAS-18902281

Steps: 1 Yield: 100%

Application of Cp₂TiCl-Promoted Radical Cyclization: A Unified Strategy for the Syntheses of Iridoid Monoterpenes


By: Khan, Hina P. A.; et al

Journal of Organic Chemistry (2018), 83(11), 6086-6092.

1.1 Solvents: Dichloromethane; 0 °C; 12 h, rt

Scheme 2 (1 Reaction)

Steps: 1 Yield: 100%



31-614-CAS-32317094

Steps: 1 Yield: 100%


Synthesis of phthalazinone PARP inhibitors treating cell proliferation disorders

Assignee: Rhizen Pharmaceuticals AG

World Intellectual Property Organization, WO2022090938 A1
2022-05-05**PatentPak** available1.1 Reagents: Triethylamine
Solvents: Dichloromethane; 1 h, rt

Scheme 3 (1 Reaction)


Steps: 1 Yield: 100%



31-339-CAS-21501973	Steps: 1 Yield: 100%	An Efficient Synthesis of Angelmarin and its Analogs
1.1 Reagents: Potassium carbonate Solvents: Dichloromethane; 6 h, rt		By: Liu, Su-You; et al
Experimental Protocols		Natural Product Communications (2017), 12(12), 1934578X1701201215.

Scheme 4 (1 Reaction)


Steps: 1 Yield: 100%



31-614-CAS-32317116	Steps: 1 Yield: 100%	Synthesis of phthalazinone PARP inhibitors treating cell proliferation disorders
1.1 Reagents: Triethylamine Solvents: Dichloromethane; 1 h, rt		Assignee: Rhizen Pharmaceuticals AG
		World Intellectual Property Organization, WO2022090938 A1 2022-05-05
		PatentPak available

Scheme 5 (1 Reaction)


Steps: 1 Yield: 100%



31-614-CAS-38514605	Steps: 1 Yield: 100%	Novel esters of 1,2,3,4,4a,5,6,7,8,9,10,10a-dodecahydrobenzo [g]quinolin-6-ol compounds, preparation method and application for treating Parkinson's disease
1.1 Reagents: Lithium hydroxide Solvents: Dimethylformamide; 0 °C; 30 min, 0 °C		Assignee: Integrative Research Laboratories Sweden AB
1.2 Solvents: Dimethylformamide; 0 °C; 4 h, cooled; 18 h, rt		World Intellectual Property Organization, WO2023208869 A1 2023-11-02
1.3 Reagents: Sodium carbonate; pH 10		PatentPak available

Scheme 6 (1 Reaction)


Steps: 1 Yield: 100%



31-339-CAS-18902280 Steps: 1 Yield: 100% 1.1 Solvents: Dichloromethane; 0 °C; 12 h, rt	Application of Cp₂TiCl-Promoted Radical Cyclization: A Unified Strategy for the Syntheses of Iridoid Monoterpenes By: Khan, Hina P. A.; et al Journal of Organic Chemistry (2018), 83(11), 6086-6092.
---	---

Scheme 7 (1 Reaction)

Steps: 1 Yield: 100%




Suppliers (3)

31-614-CAS-32317135 Steps: 1 Yield: 100% 1.1 Reagents: Triethylamine Solvents: Dichloromethane; 1 h, rt	Synthesis of phthalazinone PARP inhibitors treating cell proliferation disorders Assignee: Rhizen Pharmaceuticals AG World Intellectual Property Organization, WO2022090938 A1 2022-05-05 PatentPak available
--	---

Scheme 8 (1 Reaction)

Steps: 1 Yield: 100%




Suppliers (3)

31-614-CAS-32317104 Steps: 1 Yield: 100% 1.1 Reagents: Triethylamine Solvents: Dichloromethane; 1 h, rt	Synthesis of phthalazinone PARP inhibitors treating cell proliferation disorders Assignee: Rhizen Pharmaceuticals AG World Intellectual Property Organization, WO2022090938 A1 2022-05-05 PatentPak available
--	---

Scheme 9 (1 Reaction)

Steps: 1 Yield: 100%




Suppliers (80)

31-339-CAS-23500286 Steps: 1 Yield: 100% 1.1 Reagents: Diisobutylaluminum hydride Solvents: Dichloromethane, Hexane; 1 h, -78 °C; 10 min, -78 °C 1.2 Reagents: Potassium sodium tartrate Solvents: Water; 12 h, -78 °C 1.3 Reagents: Manganese oxide (MnO₂) Solvents: Dichloromethane; 22 h, rt 1.4 Reagents: Sodium bis(trimethylsilyl)amide Solvents: Tetrahydrofuran; 15 min, 0 °C; 15 min, 0 °C 1.5 Solvents: Tetrahydrofuran; 30 min, 0 °C 1.6 Reagents: Ammonium chloride Solvents: Water Experimental Protocols	Towards the total synthesis of aurodiox: Preparation of the key hemiacetal-lactone By: Ohara, Motoyoshi; et al Tetrahedron Letters (2021), 66, 152799.
---	--

Scheme 10 (1 Reaction)

Steps: 1 Yield: 100%




Suppliers (3)

31-614-CAS-32317131 Steps: 1 Yield: 100% 1.1 Reagents: Triethylamine Solvents: Dichloromethane; 1 h, rt	Synthesis of phthalazinone PARP inhibitors treating cell proliferation disorders Assignee: Rhizen Pharmaceuticals AG World Intellectual Property Organization, WO2022090938 A1 2022-05-05 PatentPak available
--	---

Scheme 11 (1 Reaction)

Steps: 1 Yield: 100%




Suppliers (3)

31-614-CAS-32317110 Steps: 1 Yield: 100% 1.1 Reagents: Triethylamine Solvents: Dichloromethane; 1 h, rt	Synthesis of phthalazinone PARP inhibitors treating cell proliferation disorders Assignee: Rhizen Pharmaceuticals AG World Intellectual Property Organization, WO2022090938 A1 2022-05-05 PatentPak available
--	---

Scheme 12 (1 Reaction)

Steps: 1 Yield: 100%



31-339-CAS-19051986

Steps: 1 Yield: 100%

Synthesis of 5,6-dehydrokawain and some fluorinated analogues

1.1 Solvents: Dichloromethane; 16 h, rt


By: Obi, Grace; et al

Experimental Protocols

Synthetic Communications (2018), 48(12), 1482-1486.

Scheme 13 (1 Reaction)

Steps: 1 Yield: 100%



31-339-CAS-17318523

Steps: 1 Yield: 100%

Bichromatic Photosynthesis of Coumarins by UV Filter-Enabled Olefin Metathesis

1.1 Reagents: Potassium *tert*-butoxide

By: Eivgi, Or; et al

Solvents: Tetrahydrofuran; 30 min, rt; rt → -78 °C

1.2 Solvents: Tetrahydrofuran; -78 °C; -78 °C → rt; overnight, rt

Advanced Synthesis & Catalysis (2017), 359(14), 2352-2357.


1.3 Reagents: Hydrochloric acid

Solvents: Water

Experimental Protocols

Scheme 14 (1 Reaction)

Steps: 1 Yield: 100%



31-339-CAS-18522802

Steps: 1 Yield: 100%

Highly-functionalized arene synthesis based on palladium on carbon-catalyzed aqueous dehydrogenation of cyclohexadienes and cyclohexenes

1.1 Reagents: Butyllithium

By: Yasukawa, Naoki; et al

Solvents: Tetrahydrofuran, Hexane; 15 min, 0 °C

1.2 0 °C → rt

Green Chemistry (2018), 20(6), 1213-1217.

1.3 Reagents: Ammonium chloride

Solvents: Water

Experimental Protocols

Scheme 15 (1 Reaction)

Steps: 1 Yield: 99%



31-339-CAS-22710699

Steps: 1 Yield: 99%

Process for the preparation of iloprost

1.1 **Reagents:** Potassium carbonate, 18-Crown-6
Solvents: Toluene; rt → 90 °C; 90 °C → rt


Assignee: Chinoin Gyogyszer es Vegyeszeti Termek Gyara Zrt.

World Intellectual Property Organization, WO2019202345 A2 2019-10-24

PatentPak available

Scheme 16 (1 Reaction)

Steps: 1 Yield: 99%



Suppliers (83)

31-614-CAS-36091408

Steps: 1 Yield: 99%

Synthesis of some 2-alkyl-5-(4-vinylphenyl)-2H-tetrazole compounds via the Wittig reaction and investigation by X-ray crystallography

1.1 **Reagents:** Potassium carbonate
Solvents: 1,4-Dioxane; 30 h, reflux


By: Doyduk, Dogukan; et al

Organic Communications (2022), 15(4), 333-345.

Experimental Protocols

Scheme 17 (1 Reaction)

Steps: 1 Yield: 99%



Suppliers (55)

Suppliers (85)

Double bond geometry shown

Suppliers (4)

31-339-CAS-2715751

Steps: 1 Yield: 99%

A new class of hybrid anticancer agents inspired by the synergistic effects of curcumin and genistein: Design, synthesis, and anti-proliferative evaluation

By: Chen, Qiao-Hong; et al

Bioorganic & Medicinal Chemistry Letters (2015), 25(20), 4553-4556.

1.1 **Solvents:** Toluene; 5 h, 70 °C


1.2 **Reagents:** Hydrochloric acid
Solvents: Water; pH 3

1.3 **Reagents:** Sodium bicarbonate
Solvents: Water; pH 9

Experimental Protocols

Scheme 18 (1 Reaction)

Steps: 1 Yield: 99%



Suppliers (56)

Suppliers (82)

Double bond geometry shown

31-614-CAS-40326729

Steps: 1 Yield: 99%

Catalytic 1,1-diazidation of alkenes


- 1.1 **Reagents:** Sodium hydride
Solvents: Tetrahydrofuran; 0 °C; 30 min, 0 °C
- 1.2 **Solvents:** Tetrahydrofuran; 0 °C; 0 °C → rt; overnight, rt
- 1.3 **Solvents:** Water; rt

By: Qiu, Wangzhen; et al
 Nature Communications (2024), 15(1), 3632.

Experimental Protocols

Scheme 19 (1 Reaction)

Steps: 1 Yield: 99%



Double bond geometry shown

Suppliers (86)

Double bond geometry shown

31-614-CAS-35648063

Steps: 1 Yield: 99%

Unexplored Vinylic-Substituted 5-Benzylidenethiazolidine-2,4-diones: Synthesis and DFT/NMR Stereochemical Assignment

- 1.1 **Solvents:** Tetrahydrofuran; 15 h, rt


By: Dupommier, Dorian; et al

Experimental Protocols

Journal of Organic Chemistry (2023), 88(6), 3724-3739.

Scheme 20 (1 Reaction)


Steps: 1 Yield: 99%



Absolute stereochemistry shown
 Double bond geometry shown

Suppliers (4)

Suppliers (188)




Absolute stereochemistry shown,
 Rotation (+)
 Double bond geometry shown

31-614-CAS-35036998	Steps: 1 Yield: 99%	Stereocontrolled α-Allylation of α-Branched N-tert-Butanesulfinyl Ketimines via a Michael-Wittig Cascade for the Construction of Acyclic Quaternary Stereocenters
1.1 Reagents: Potassium <i>tert</i> -butoxide Solvents: Tetrahydrofuran; -78 °C; 30 min, -78 °C		By: Zhu, Chong-Lin; et al
1.2 Solvents: Tetrahydrofuran; -78 °C; 30 min, -78 °C $\rightarrow$ 0 °C		Organic Letters (2022), 24(48), 8925-8929.
1.3 Reagents: Lithium bis(trimethylsilyl)amide Solvents: Tetrahydrofuran; 0 °C; 0 °C $\rightarrow$ rt; 12 h, rt		
1.4 Reagents: Ammonium chloride Solvents: Water		
Experimental Protocols		

Scheme 21 (1 Reaction)

Steps: 1 Yield: 99%




Suppliers (83)

31-339-CAS-23471040	Steps: 1 Yield: 99%	Donor-acceptor conjugated ladder polymer via aromatization-driven thermodynamic annulation
1.1 Reagents: Potassium <i>tert</i> -butoxide Solvents: Tetrahydrofuran; 10 min, rt		By: Lee, Jongbok; et al
1.2 6 h, reflux		Polymer Chemistry (2018), 9(13), 1603-1609.
1.3 Solvents: Water		
Experimental Protocols		

Scheme 22 (1 Reaction)

Steps: 1 Yield: 99%



Suppliers (93)

Suppliers (90)


Double bond geometry shown

Suppliers (9)

31-339-CAS-19051990	Steps: 1 Yield: 99%	Synthesis of 5,6-dehydrokawain and some fluorinated analogues
1.1 Solvents: Dichloromethane; 16 h, rt		By: Obi, Grace; et al
Experimental Protocols		Synthetic Communications (2018), 48(12), 1482-1486.

Scheme 23 (1 Reaction)

Steps: 1 Yield: 99%




Suppliers (83)

31-614-CAS-37514350 Steps: 1 Yield: 99% 1.1 Catalysts: Potassium <i>tert</i>-butoxide Solvents: Tetrahydrofuran; 60 h, 0 °C → 20 °C 1.2 Reagents: Ammonium chloride Solvents: Water Experimental Protocols	Cationic β-Scission of C-H and C-C Bonds for Selective Dimerization and Subsequent Sulfur-Free RAFT Polymerization of α-Methylstyrene and Isobutylene By: Tanimoto, Takayuki; et al Angewandte Chemie, International Edition (2023), 62(43), e202307791.
---	---

Scheme 24 (1 Reaction)


Steps: 1 Yield: 99%



31-614-CAS-40326725 Steps: 1 Yield: 99% 1.1 Reagents: Sodium hydride Solvents: Tetrahydrofuran; 0 °C; 30 min, 0 °C 1.2 Solvents: Tetrahydrofuran; 0 °C; 0 °C → rt; overnight, rt 1.3 Solvents: Water; rt Experimental Protocols	Catalytic 1,1-diazidation of alkenes By: Qiu, Wangzhen; et al Nature Communications (2024), 15(1), 3632.
---	--

Scheme 25 (1 Reaction)


Steps: 1 Yield: 99%




31-339-CAS-21501976 Steps: 1 Yield: 99% 1.1 Reagents: Potassium carbonate Solvents: Dichloromethane; 6 h, rt Experimental Protocols	An Efficient Synthesis of Angelmarin and its Analogs By: Liu, Su-You; et al Natural Product Communications (2017), 12(12), 1934578X1701201215.
---	--

Scheme 26 (1 Reaction)


Steps: 1 Yield: 99%




31-614-CAS-40326738	Steps: 1 Yield: 99%	Catalytic 1,1-diazidation of alkenes
1.1 Reagents: Sodium hydride Solvents: Tetrahydrofuran; 0 °C; 30 min, 0 °C	By: Qiu, Wangzhen; et al Nature Communications (2024), 15(1), 3632.	
1.2 Solvents: Tetrahydrofuran; 0 °C; 0 °C → rt; overnight, rt		
1.3 Solvents: Water; rt		
Experimental Protocols		

Scheme 27 (1 Reaction)		Steps: 1 Yield: 99%
 Absolute stereochemistry shown Absolute stereochemistry shown, Rotation (-)		
 Suppliers (83)		

31-339-CAS-19609316	Steps: 1 Yield: 99%	Studies towards the synthesis of ertugli flozin from L-Arabinose
1.1 Reagents: Butyllithium Catalysts: 12-Crown-4 Solvents: Tetrahydrofuran, Hexane; -78 °C; 1 h, -40 °C; -40 °C → -78 °C		By: Triantakonsanti, Virginia V.; et al Tetrahedron (2018), 74(39), 5700-5708.
1.2 Solvents: Tetrahydrofuran; -78 °C; 2 h, < -60 °C; 12 h, 20 °C		
Experimental Protocols		

Scheme 28 (1 Reaction)		Steps: 1 Yield: 99%
 Double bond geometry shown		
 Suppliers (5)		
 Suppliers (85)		


31-339-CAS-4523553	Steps: 1 Yield: 99%	A new class of hybrid anticancer agents inspired by the synergistic effects of curcumin and genistein: Design, synthesis, and anti-proliferative evaluation By: Chen, Qiao-Hong; et al Bioorganic & Medicinal Chemistry Letters (2015), 25(20), 4553-4556.
1.1 Solvents: Toluene; 5 h, 70 °C		
1.2 Reagents: Hydrochloric acid Solvents: Water; pH 3		
1.3 Reagents: Sodium bicarbonate Solvents: Water; pH 9		
Experimental Protocols		

Scheme 29 (1 Reaction)		Steps: 1 Yield: 99%
 Absolute stereochemistry shown		
 Suppliers (69)		

31-049-CAS-15661133 Steps: 1 Yield: 99% 1.1 Reagents: Potassium carbonate Catalysts: 18-Crown-6 Solvents: Dichloromethane; 21.5 h, reflux	Minimum structural requirements for cell membrane leakage-mediated anti-MRSA activity of macrocyclic bis (bibenzyl)s By: Fujii, Kana; et al Bioorganic & Medicinal Chemistry Letters (2016), 26(9), 2324-2327.
--	---

Scheme 30 (1 Reaction)

Steps: 1 Yield: 99%



Suppliers (5)


Suppliers (75)

Double bond geometry shown

31-339-CAS-19873345 Steps: 1 Yield: 99% 1.1 Solvents: Dichloromethane; rt; 24 h, rt Experimental Protocols	Applications of Thermal Activation, Ball-milling and Aqueous Medium in Stereoselective Michael Addition of Nitromethane to En-ynones Catalyzed by Chiral Squaramides By: Ignatiuk, Zaneta A.; et al Advanced Synthesis & Catalysis (2019), 361(5), 1108-1116.
--	--

Scheme 31 (1 Reaction)

Steps: 1 Yield: 99%



Suppliers (90)

Suppliers (82)


Double bond geometry shown

Suppliers (16)

31-339-CAS-19051988 Steps: 1 Yield: 99% 1.1 Solvents: Dichloromethane; 16 h, rt Experimental Protocols	Synthesis of 5,6-dehydrokawain and some fluorinated analogues By: Obi, Grace; et al Synthetic Communications (2018), 48(12), 1482-1486.
--	--

Scheme 32 (1 Reaction)

Steps: 1 Yield: 99%



Suppliers (56)

Suppliers (86)

Double bond geometry shown

31-614-CAS-40326736	Steps: 1 Yield: 99%	Catalytic 1,1-diazidation of alkenes
1.1 Reagents: Sodium hydride Solvents: Tetrahydrofuran; 0 °C; 30 min, 0 °C	By: Qiu, Wangzhen; et al Nature Communications (2024), 15(1), 3632.	
1.2 Solvents: Tetrahydrofuran; 0 °C; 0 °C → rt; overnight, rt		
1.3 Solvents: Water; rt		
Experimental Protocols		

Scheme 33 (1 Reaction)		Steps: 1 Yield: 99%
Suppliers (83)	Double bond geometry shown Suppliers (48)	Double bond geometry shown Suppliers (3)


31-339-CAS-18522804	Steps: 1 Yield: 99%	Highly-functionalized arene synthesis based on palladium on carbon-catalyzed aqueous dehydrogenation of cyclohexadienes and cyclohexenes By: Yasukawa, Naoki; et al Green Chemistry (2018), 20(6), 1213-1217.
1.1 Reagents: Butyllithium Solvents: Tetrahydrofuran, Hexane; 15 min, 0 °C		
1.2 0 °C → rt		
1.3 Reagents: Ammonium chloride Solvents: Water		
Experimental Protocols		

Scheme 34 (1 Reaction)		Steps: 1 Yield: 99%
Suppliers (83)	Suppliers (55)	Suppliers (65)

31-339-CAS-20730149	Steps: 1 Yield: 99%	A Simple and Robust Process for Large- Scale Synthesis of 4-Methylenepiperidine Hydrochloride
1.1 Reagents: Potassium <i>tert</i> -butoxide Solvents: <i>tert</i> -Butyl methyl ether; 30 min, 20 - 30 °C; 2 h, 20 - 30 °C	By: Chen, Ao; et al Letters in Organic Chemistry (2019), 16(10), 802-806.	
1.2 1 h, 20 - 30 °C; 3 h, 20 - 30 °C		
1.3 Reagents: Water; 0.5 h, 20 - 30 °C		
1.4 Reagents: Hydrochloric acid Solvents: Methanol; 2 h, 0 - 10 °C; 3 h, 0 - 5 °C		
Experimental Protocols		

Scheme 35 (1 Reaction)

Steps: 1 Yield: 98%



Suppliers (5)

Suppliers (83)

Suppliers (8)

31-614-CAS-45598149

Steps: 1 Yield: 98%

Development of a Practical and Scalable Manufacturing Process for Ciprofol

By: Dong, Boliang; et al


Organic Process Research & Development (2025), 29(5), 1291-1298.

- 1.1 **Reagents:** Potassium *tert*-butoxide
Solvents: Tetrahydrofuran; 30 min, 10 - 20 °C; 10 h, 20 °C → 65 °C; 65 °C → 0 °C
- 1.2 **Reagents:** Hydrochloric acid
Solvents: Water; pH 6 - 7, 0 - 10 °C

Experimental Protocols

Scheme 36 (1 Reaction)


Steps: 1 Yield: 98%



Suppliers (45)

Suppliers (75)

Suppliers (75)



Absolute stereochemistry shown
 Double bond geometry shown

31-339-CAS-18801553

Steps: 1 Yield: 98%

Two-Pot Synthesis of Chiral 1,3-syn-Diols through Asymmetric Organocatalytic Aldol and Wittig Reactions Followed by Domino Hemiacetal/Oxy-Michael Reactions

By: Hayashi, Yujiro; et al


Chemistry - A European Journal (2018), 24(19), 4909-4915.

- 1.1 **Catalysts:** (2*S*)- α,α -Bis[3,5-bis(trifluoromethyl)phenyl]-2-pyrrolidinemethanol
Solvents: Acetonitrile, Toluene, Water; 24 h, rt; 24 °C → 0 °C
- 1.2 **Reagents:** Sodium hydride
Solvents: Dimethylformamide, Tetrahydrofuran; 0 °C; 30 min, rt
- 1.3 **Reagents:** Ammonium chloride
Solvents: Water; rt

Experimental Protocols

Scheme 37 (1 Reaction)

Steps: 1 Yield: 98%



Suppliers (56)

Suppliers (89)

Double bond geometry shown

Suppliers (7)

31-614-CAS-40326723

Steps: 1 Yield: 98%

Catalytic 1,1-diazidation of alkenes


- 1.1 **Reagents:** Sodium hydride
Solvents: Tetrahydrofuran; 0 °C; 30 min, 0 °C
- 1.2 **Solvents:** Tetrahydrofuran; 0 °C; 0 °C → rt; overnight, rt
- 1.3 **Solvents:** Water; rt

By: Qiu, Wangzhen; et al
 Nature Communications (2024), 15(1), 3632.

Experimental Protocols

Scheme 38 (1 Reaction)

Steps: 1 Yield: 98%



Suppliers (37)

Suppliers (128)

Suppliers (7)

31-614-CAS-34094616

Steps: 1 Yield: 98%

Radical cation Diels-Alder reactions of arylidene cycloalkanes


- 1.1 **Reagents:** Potassium *tert*-butoxide
Solvents: Tetrahydrofuran; rt; 1 h, 70 °C
- 1.2 3 h, 70 °C; 70 °C → rt; 1 h, rt

By: Nakayama, Kaii; et al
 Beilstein Journal of Organic Chemistry (2022), 18, 1100-1106.

Experimental Protocols

Scheme 39 (1 Reaction)


Steps: 1 Yield: 98%



Suppliers (20)

Suppliers (90)

Suppliers (188)




Double bond geometry shown

31-339-CAS-20621896 Steps: 1 Yield: 98% 1.1 Catalysts: [1,1'-Bis(diphenylphosphino)ferrocene]dichloropalladium, Methanesulfonamide, 1,1,1-trifluoro-<i>N</i>-[(trifluoromethyl)sulfonyl]-, barium salt (2:1) Solvents: Dimethyl sulfoxide; 48 h, 100 °C 1.2 Solvents: Water; 12 h, rt	A Ba/Pd Catalytic System Enables Dehydrative Cross-Coupling and Excellent E-Selective Wittig Reactions By: Xie, Peizhong; et al Organic Letters (2019), 21(17), 7055-7059.
--	--

Scheme 40 (1 Reaction)


Steps: 1 Yield: 98%



31-339-CAS-9711159 Steps: 1 Yield: 98% 1.1 Solvents: Dichloromethane; 0 °C; 0 °C → rt; 12 h, rt Experimental Protocols	Asymmetric Total Synthesis of Bioactive Natural Lipid Mycalol By: Das, Subhendu; et al Journal of Organic Chemistry (2015), 80(12), 6467-6489.
---	--

Scheme 41 (1 Reaction)


Steps: 1 Yield: 98%



31-614-CAS-35473008 Steps: 1 Yield: 98% 1.1 Reagents: Sodium hydride Solvents: Tetrahydrofuran; 45 - 90 min, rt 1.2 29 h, rt 1.3 Reagents: Hydrochloric acid Solvents: Water; pH 4 - 5, 0 °C Experimental Protocols	Ultrasound-assisted wittig reaction for the synthesis of 3-substituted 4-chloroquinolines and quinolin-4(1H)-ones with extended π-conjugated systems By: Frites, Wahiba; et al Journal of Chemistry (2022), 4807767.
---	--

Scheme 42 (1 Reaction)


Steps: 1 Yield: 98%



31-339-CAS-19051991	Steps: 1 Yield: 98%	Synthesis of 5,6-dehydrokawain and some fluorinated analogues
1.1 Solvents: Dichloromethane; 16 h, rt		By: Obi, Grace; et al
Experimental Protocols		Synthetic Communications (2018), 48(12), 1482-1486.

Scheme 43 (1 Reaction)


Steps: 1 Yield: 98%



31-339-CAS-22828505	Steps: 1 Yield: 98%	Asymmetric total synthesis of dihydroiso coumarins: 6-methoxymellein, kigelin and fusarentin 6, 7 dimethyl ether by employing proline catalyzed asymmetric α -aminooxylation
1.1 Solvents: Dichloromethane; 12 h, rt		By: Markad, Sachin B.; et al
Experimental Protocols		Tetrahedron (2020), 76(46), 131524.

Scheme 44 (1 Reaction)


Steps: 1 Yield: 98%



31-614-CAS-36091398	Steps: 1 Yield: 98%	Synthesis of some 2-alkyl-5-(4-vinylphenyl)-2H-tetrazole compounds via the Wittig reaction and investigation by X-ray crystallography
1.1 Reagents: Potassium carbonate Solvents: 1,4-Dioxane; 30 h, reflux		By: Doyduk, Dogukan; et al
Experimental Protocols		Organic Communications (2022), 15(4), 333-345.

Scheme 45 (1 Reaction)


Steps: 1 Yield: 98%



31-339-CAS-22829074 Steps: 1 Yield: 98%	Asymmetric total synthesis of dihydroiso coumarins: 6-methoxymellein, kigelin and fusarentin 6, 7 dimethyl ether by employing proline catalyzed asymmetric α-aminoxylation
1.1 Solvents: Dichloromethane; 12 h, rt Experimental Protocols	By: Markad, Sachin B.; et al Tetrahedron (2020), 76(46), 131524.

Scheme 46 (1 Reaction)

Steps: 1 Yield: 98%




Suppliers (49)

Suppliers (12)

31-339-CAS-18213499 Steps: 1 Yield: 98%	Methods for preparation of lycopenes from C15-Wittig salts and methods for purification of high all-E containing and high 6Z containing C15-Wittig salts
1.1 Solvents: Methanol; rt $\rightarrow$ 30 $^{\circ}$ C; 30 min, 30 $^{\circ}$ C 1.2 Reagents: Potassium carbonate Solvents: Ethyl acetate, Heptane, Water; 30 $^{\circ}$ C $\rightarrow$ rt; rt $\rightarrow$ 60 $^{\circ}$ C; 3 h, 60 $^{\circ}$ C; 60 $^{\circ}$ C $\rightarrow$ 100 $^{\circ}$ C; 100 $^{\circ}$ C $\rightarrow$ 80 $^{\circ}$ C; 16 h, 80 $^{\circ}$ C Experimental Protocols	Assignee: Allied Biotech Corporation European Patent Organization, EP2957565 A1 2015-12-23 PatentPak available

Scheme 47 (1 Reaction)

Steps: 1 Yield: 98%



Suppliers (77)


Suppliers (90)

Suppliers (25)

31-339-CAS-16193708 Steps: 1 Yield: 98%	Facile, direct reaction of benzaldehydes to 3-arylprop-2-enoic acids and 3-arylprop-2-ynoic acids in aqueous medium
1.1 Reagents: Sodium hydroxide Solvents: Water; 23 h, 75 $^{\circ}$ C; cooled 1.2 Reagents: Hydrochloric acid Solvents: Water; acidified Experimental Protocols	By: Thiemann, Thies; et al International Journal of Organic Chemistry (2016), 6(2), 126-141.

Scheme 48 (1 Reaction)

Steps: 1 Yield: 98%



Suppliers (83)


Suppliers (79)

Suppliers (5)

31-614-CAS-42509075	Steps: 1 Yield: 98%	High-Energy Ball Milling Enables an Ultra- Fast Wittig Olefination Under Ambient and Solvent-free Conditions
1.1 Reagents: Potassium <i>tert</i> -butoxide; 30 s, 25 °C		By: Templ, Johanna; et al
1.2 Reagents: Ammonium chloride Solvents: Water		Angewandte Chemie, International Edition (2024), 63(49), e202411536.
Experimental Protocols		

Scheme 49 (1 Reaction)


Steps: 1 Yield: 98%



31-339-CAS-19051987	Steps: 1 Yield: 98%	Synthesis of 5,6-dehydrokawain and some fluorinated analogues
1.1 Solvents: Dichloromethane; 16 h, rt		By: Obi, Grace; et al
Experimental Protocols		Synthetic Communications (2018), 48(12), 1482-1486.

Scheme 50 (1 Reaction)

Steps: 1 Yield: 98%



31-339-CAS-17869069	Steps: 1 Yield: 98%	Synthesis of stilbene-quinone hybrids through Heck reactions in PEG-400
1.1 Reagents: Potassium <i>tert</i> -butoxide Solvents: Tetrahydrofuran; 10 min, rt		By: Demidoff, Felipe C.; et al
1.2 Solvents: Tetrahydrofuran; rt; 60 min, rt		Synthesis (2017), 49(23), 5217-5223.
1.3 Reagents: Ammonium chloride Solvents: Water		