

Initiating Search

September 22, 2025, 3:18 PM

• Search:

Filtered By:


Yield: 90-100%, 80-89%, 70-79%, <10%

Document Type: Journal

Language: English

Publication Year: 2015 to 2025

Publication Name: Journal of Medicinal Chemistry, European Journal of Medicinal Chemistry, Organic Letters, Journal of Organic Chemistry, Angewandte Chemie, International Edition, Bioorganic & Medicinal Chemistry Letters, Bioorganic & Medicinal Chemistry, Journal of the American Chemical Society, Chemistry - A European Journal, Chemical Communications (Cambridge, United Kingdom), European Journal of Organic Chemistry, ACS Medicinal Chemistry Letters, Nature (London, United Kingdom), Nature Communications, Chinese Chemical Letters, Green Chemistry, Nature Chemistry



Structure Match: Substructure

Search Tasks

Task	Result Type	View
Exported: Returned Reaction Results + Filters (38,955)	 Reactions	View Results

Copyright © 2025 American Chemical Society (ACS). All Rights Reserved.


Internal use only. Redistribution is subject to the terms of your CAS SciFinder License Agreement and CAS information Use Policies.

Reactions (20)

[View in CAS SciFinder](#)

Scheme 1 (1 Reaction)

Steps: 1 Yield: 100%


 Supplier (1)

31-367-CAS-16263743

Steps: 1 Yield: 100%

1.1 **Reagents:** 4-Methylmorpholine, Phosphorodichloridic acid, phenyl ester
Solvents: Ethyl acetate; 5 min, -20 °C; 1 h, -20 °C

Experimental Protocols


Novel broad-spectrum and long-acting parenteral cephalosporins having an acyl cyanamide moiety at the C-3 terminal: Synthesis and structure-activity relationships

By: Yokoo, Katsuki; et al

European Journal of Medicinal Chemistry (2016), 124, 698-712.

Scheme 2 (1 Reaction)

Steps: 1 Yield: 100%


 Suppliers (3)

 Suppliers (70)

31-614-CAS-42895510

Steps: 1 Yield: 100%

1.1 **Reagents:** 1-Hydroxybenzotriazole, Diisopropylethylamine, 1-Ethyl-3-(3'-dimethylaminopropyl)carbodiimide hydrochloride
Solvents: Dimethylformamide; 2 h, rt

1.2 **Reagents:** Water; rt

Experimental Protocols



Rational Design of Macrocyclic Noncovalent Inhibitors of SAR S-CoV-2 M^{Pro} from a DNA-Encoded Chemical Library Screening Hit That Demonstrate Potent Inhibition against Pan-Coronavirus Homologues and Nirmatrelvir-Resistant Variants

By: Wang, Xu; et al

Journal of Medicinal Chemistry (2024), 67(21), 19623-19667.

Scheme 3 (1 Reaction)

Steps: 1 Yield: 100%



Suppliers (111)

Supplier (1)

31-367-CAS-16466988

Steps: 1 Yield: 100%

[2.2.1]-Bicyclic sultams as potent androgen receptor antagonists

By: Shan, Weifang; et al

Bioorganic & Medicinal Chemistry Letters (2016), 26(23), 5707-5711.


1.1 Reagents: Oxalyl chloride

Catalysts: Pyridine; 50 °C

1.2 -

Scheme 4 (1 Reaction)



Steps: 1 Yield: 100%



Suppliers (69)

Suppliers (81)

Suppliers (67)



Suppliers (324)

Suppliers (88)

31-367-CAS-19877594

Steps: 1 Yield: 100%

Consecutive Five-Component Ugi-4CR-CAL B-Catalyzed Aminolysis Sequence and Concatenation with Transition Metal Catalysis in a One-Pot Fashion to Substituted Triamides

By: Gesse, Pascal; et al

European Journal of Organic Chemistry (2019), 2019(11), 2150-2157.

1.1 Solvents: Methanol; 90 min, 20 °C


1.2 24 h, 20 °C

1.3 Catalysts: Lipase CaLB, Candida antarctica; 24 h, 45 °C

Experimental Protocols

Scheme 5 (1 Reaction)

Steps: 1 Yield: 100%




Suppliers (87)

31-367-CAS-19469920 Steps: 1 Yield: 100% 1.1 Reagents: 1-Hydroxybenzotriazole, 1-Ethyl-3-(3'-dimethylamino)carbodiimide hydrochloride Solvents: Dimethylformamide; 1 h, rt	Cefiderocol (S-649266), A new siderophore cephalosporin exhibiting potent activities against <i>Pseudomonas aeruginosa</i> and other gram-negative pathogens including multi-drug resistant bacteria: Structure activity relationship By: Aoki, Toshiaki; et al European Journal of Medicinal Chemistry (2018), 155, 847-868.
Experimental Protocols	

Scheme 6 (1 Reaction)


Steps: 1 Yield: 100%



31-367-CAS-23203826 Steps: 1 Yield: 100% 1.1 Reagents: 1,1'-Carbonyldiimidazole Solvents: Dimethylformamide; 10 min, rt 1.2 0.5 - 2 h, rt 1.3 Reagents: Ammonium chloride Solvents: Water 1.4 Reagents: Hydrochloric acid Solvents: Water; neutralized	Novel amide and imidazole compounds as potent hematopoietic prostaglandin D₂ synthase inhibitors By: Olson, Kirk L.; et al Bioorganic & Medicinal Chemistry Letters (2021), 34, 127759.
Experimental Protocols	

Scheme 7 (1 Reaction)


Steps: 1 Yield: 100%



31-367-CAS-23663323 Steps: 1 Yield: 100% 1.1 Reagents: Diisopropylethylamine, HOAt, <i>O</i> -(7-Azabenzotriazol-1-yl)- <i>N,N,N,N</i> -tetramethyluronium hexafluoro phosphate Solvents: Dimethylformamide; 3 h, rt	Automated design of macrocycles for therapeutic applications: From small molecules to peptides and proteins By: Sindhikara, Dan; et al Journal of Medicinal Chemistry (2020), 63(20), 12100-12115.
Experimental Protocols	

Scheme 8 (1 Reaction)

Steps: 1 Yield: 100%



Suppliers (83)

Suppliers (81)

Suppliers (3)

31-367-CAS-23202236

Steps: 1 Yield: 100%

Novel amide and imidazole compounds as potent hematopoietic prostaglandin D₂ synthase inhibitors

By: Olson, Kirk L.; et al



Bioorganic & Medicinal Chemistry Letters (2021), 34, 127759.

- 1.1 **Reagents:** 1,1'-Carbonyldiimidazole
Solvents: Dimethylformamide; 10 min, rt
- 1.2 0.5 - 2 h, rt
- 1.3 **Reagents:** Ammonium chloride
Solvents: Water
- 1.4 **Reagents:** Hydrochloric acid
Solvents: Water; neutralized


Experimental Protocols

Scheme 9 (1 Reaction)

Steps: 1 Yield: 100%



Suppliers (81)



Supplier (1)

31-367-CAS-19933620

Steps: 1 Yield: 100%

Exploration of novel macrocyclic dipeptide N-benzyl amides as proteasome inhibitors

By: Yu, Jianjun; et al



European Journal of Medicinal Chemistry (2019), 164, 423-439.

- 1.1 **Reagents:** 1-Hydroxybenzotriazole, 1-Ethyl-3-(3'-dimethylaminopropyl)carbodiimide hydrochloride
Solvents: Dichloromethane; 30 min, 0 °C
- 1.2 **Reagents:** Diisopropylethylamine; 5 h, rt


Experimental Protocols

Scheme 10 (1 Reaction)

Steps: 1 Yield: 100%



Suppliers (69)




Suppliers (70)

31-614-CAS-34006645	Steps: 1 Yield: 100%	Discovery, Structure-Activity Relationships, and In Vivo Evaluation of Novel Aryl Amides as Brain Penetrant Adaptor Protein 2-Associated Kinase 1 (AAK1) Inhibitors for the Treatment of Neuropathic Pain
1.1 Reagents: Diisopropylethylamine, <i>O</i> -(7-Azabenzotriazol-1-yl)- <i>N,N,N,N</i> -tetramethyluronium hexafluorophosphate Solvents: Dichloromethane; 16 h, rt		By: Hartz, Richard A.; et al Journal of Medicinal Chemistry (2021), 64(15), 11090-11128.
Experimental Protocols		

Scheme 11 (1 Reaction)


Steps: 1 Yield: 100%



31-614-CAS-31186804	Steps: 1 Yield: 100%	Generation of Highly Selective, Potent, and Covalent G Protein-Coupled Receptor Kinase 5 Inhibitors
1.1 Reagents: Diisopropylethylamine, <i>O</i> -(7-Azabenzotriazol-1-yl)- <i>N,N,N,N</i> -tetramethyluronium hexafluorophosphate Solvents: Dimethylformamide; 12 h, rt		By: Rowlands, Rachel A.; et al Journal of Medicinal Chemistry (2021), 64(1), 566-585.
Experimental Protocols		

Scheme 12 (1 Reaction)


Steps: 1 Yield: 100%



31-367-CAS-16417467	Steps: 1 Yield: 100%	Synthesis of Multivalent Organotellurium Chain-Transfer Agents by Post-modification and Their Applications in Living Radical Polymerization
1.1 Reagents: Diisopropylethylamine, <i>N</i> -[1-(Cyano-2-ethoxy-2-oxoethylideneaminoxy)dimethylamino(morpholino)uronium hexafluorophosphate Solvents: Dimethylformamide; 0.5 h, rt		By: Fan, Weijia; et al Chemistry - A European Journal (2016), 22(47), 17006-17010.
1.2 3 h, rt		
Experimental Protocols		

Scheme 13 (1 Reaction)


Steps: 1 Yield: 100%



31-614-CAS-38842313	Steps: 1 Yield: 100%	Discovery of PFI-6, a small-molecule chemical probe for the YE ATS domain of MLLT1 and MLLT3
1.1 Reagents: Diisopropylethylamine, <i>O</i> -(7-Azabenzotriazol-1-yl)- <i>N,N,N,N</i> -tetramethyluronium hexafluorophosphate Solvents: Dimethylformamide, Tetrahydrofuran; 16 h, rt		By: Raux, Brigitt; et al
Experimental Protocols		Bioorganic & Medicinal Chemistry Letters (2024), 98, 129546.

Scheme 14 (1 Reaction)

Steps: 1 Yield: 100%




Suppliers (112)

31-367-CAS-17755169	Steps: 1 Yield: 100%	Synthesis and antibacterial evaluation of novel 11-O-carbamoyl clarithromycin ketolides
1.1 Reagents: Dicyclohexylcarbodiimide, 1-Hydroxybenzotriazole Solvents: Tetrahydrofuran; 0 °C; 6 h, 0 °C		By: Jia, Li; et al
1.2 4 h, rt		Bioorganic & Medicinal Chemistry Letters (2017), 27(16), 3693-3697.
Experimental Protocols		

Scheme 15 (1 Reaction)

Steps: 1 Yield: 100%



Double bond geometry shown

Suppliers (81)

Double bond geometry shown


Suppliers (76)

Suppliers (15)

31-367-CAS-5606384	Steps: 1 Yield: 100%	Direct amidation of carboxylic acids with amines under microwave irradiation using silica gel as a solid support
1.1 Catalysts: Silica; 80 min, 130 °C		By: Ojeda-Porras, Andrea; et al
Experimental Protocols		Green Chemistry (2015), 17(5), 3157-3163.

Scheme 16 (1 Reaction)

Steps: 1 Yield: 100%



Suppliers (74)


Suppliers (81)

Supplier (1)

31-367-CAS-23204560 Steps: 1 Yield: 100% 1.1 Reagents: 1,1'-Carbonyldiimidazole Solvents: Dimethylformamide; 10 min, rt 1.2 0.5 - 2 h, rt 1.3 Reagents: Ammonium chloride Solvents: Water 1.4 Reagents: Hydrochloric acid Solvents: Water; neutralized Experimental Protocols	Novel amide and imidazole compounds as potent hematopoietic prostaglandin D₂ synthase inhibitors By: Olson, Kirk L.; et al Bioorganic & Medicinal Chemistry Letters (2021), 34, 127759.
--	--

Scheme 17 (1 Reaction)


Steps: 1 Yield: 100%



31-614-CAS-36469225 Steps: 1 Yield: 100% 1.1 Reagents: Oxalyl chloride Catalysts: Dimethylformamide Solvents: Dichloromethane; 4 h, rt 1.2 Reagents: Triethylamine Solvents: Toluene; 5 h, rt Experimental Protocols	Functionalized sulfonyl anthranilic acid derivatives inhibit replication of all the four dengue serotypes By: Felicetti, Tommaso; et al European Journal of Medicinal Chemistry (2023), 252, 115283.
--	--

Scheme 18 (1 Reaction)


Steps: 1 Yield: 100%




31-367-CAS-21800748 Steps: 1 Yield: 100% 1.1 Reagents: Triethylamine, Isobutyl chloroformate Solvents: Dichloromethane; 0 °C; 10 min, 0 °C; 0 °C → rt; 20 min, rt 1.2 60 min, rt Experimental Protocols	Amide Synthesis by Nickel/Photoredox-Catalyzed Direct Carbamoylation of (Hetero)Aryl Bromides By: Alandini, Nurtalya; et al Angewandte Chemie, International Edition (2020), 59(13), 5248-5253.
--	---

Scheme 19 (1 Reaction)

Steps: 1 Yield: 100%



31-367-CAS-23200316 Steps: 1 Yield: 100% 1.1 Reagents: 1,1'-Carbonyldiimidazole Solvents: Dimethylformamide; 10 min, rt 1.2 0.5 - 2 h, rt 1.3 Reagents: Ammonium chloride Solvents: Water 1.4 Reagents: Hydrochloric acid Solvents: Water; neutralized Experimental Protocols	Novel amide and imidazole compounds as potent hematopoietic prostaglandin D₂ synthase inhibitors By: Olson, Kirk L.; et al Bioorganic & Medicinal Chemistry Letters (2021), 34, 127759.
---	---



31-614-CAS-40274070 Steps: 1 Yield: 100% 1.1 Catalysts: Boronic acid, <i>B</i>-[2,6-bis[4-(trifluoromethyl)phenoxy]phenyl]-; 2 h, 1 mbar, 250 °C; 250 °C → 25 °C 1.2 Reagents: 1,1,2,2-Tetrachloroethane Solvents: Dichloromethane; 15 min, 25 °C 1.3 4 h, 25 °C Experimental Protocols	A Strategy for Improving the Efficiency of Boronic Acid Catalysis in the Synthesis of Amides By: Almetwali, Fatima; et al European Journal of Organic Chemistry (2023), 26(45), e202300720.
---	--