

Initiating Search

November 4, 2025, 10:08 PM

•• Search: Hofmann rearrangement

Search Tasks

Task	Result Type	View
Returned Reference Results (1,295)	 References	View Results
Exported: Retrieved Related Reaction Results (631)	 Reactions	View Results

Copyright © 2025 American Chemical Society (ACS). All Rights Reserved.


Internal use only. Redistribution is subject to the terms of your CAS SciFinder License Agreement and CAS information Use Policies.

Reactions (50)

[View in CAS SciFinder](#)

Scheme 1 (1 Reaction)

Steps: 1 Yield: 100%


 Suppliers (72)

 Suppliers (444)

 Suppliers (37)

31-448-CAS-1807173

Steps: 1 Yield: 100%

Hofmann Rearrangement of Carboxamides Mediated by Hypervalent Iodine Species Generated in Situ from Iodobenzene and Oxone: Reaction Scope and Limitations

1.1 Reagents: Iodobenzene, Potassium peroxymonosulfate sulfate (2KHSO₅.KHSO₄.K₂SO₄); 7 - 12 h, rt


Experimental Protocols

By: Zagulyaeva, Aleksandra A.; et al

Organic Letters (2010), 12(20), 4644-4647.

Scheme 2 (2 Reactions)

Steps: 1 Yield: 98-100%


 Suppliers (55)

 Suppliers (444)

 Suppliers (6)

31-448-CAS-3937974

Steps: 1 Yield: 100%

Hofmann Rearrangement of Carboxamides Mediated by Hypervalent Iodine Species Generated in Situ from Iodobenzene and Oxone: Reaction Scope and Limitations

1.1 Reagents: Iodobenzene, Potassium peroxymonosulfate sulfate (2KHSO₅.KHSO₄.K₂SO₄); 7 - 12 h, rt

Experimental Protocols

By: Zagulyaeva, Aleksandra A.; et al

Organic Letters (2010), 12(20), 4644-4647.

31-448-CAS-12291373

Steps: 1 Yield: 98%

Electrochemically induced Hofmann rearrangement

1.1 Reagents: Tetraethylammonium bromide
Solvents: Acetonitrile

By: Matsumura, Yoshihiro; et al

Tetrahedron Letters (1997), 38(51), 8879-8882.

Scheme 3 (3 Reactions)

Steps: 1 Yield: 100%



Suppliers (62)


Suppliers (444)

Suppliers (5)

31-448-CAS-10417053 Steps: 1 Yield: 100%	New method for Hofmann rearrangement
1.1 Reagents: <i>N</i> -Bromosuccinimide, Mercuric acetate Solvents: Dimethylformamide	By: Jew, Sang Sup; et al Tetrahedron Letters (1990), 31(11), 1559-62.
31-448-CAS-1282632 Steps: 1 Yield: 100%	New method for Hofmann rearrangement
1.1 Reagents: 1,3-Dibromo-5,5-dimethylhydantoin, Silver acetate Solvents: Dimethylformamide	By: Jew, Sang Sup; et al Tetrahedron Letters (1990), 31(11), 1559-62.
31-448-CAS-12545923 Steps: 1 Yield: 100%	New method for Hofmann rearrangement
1.1 Reagents: 1,3-Dibromo-5,5-dimethylhydantoin, Mercuric acetate Solvents: Dimethylformamide	By: Jew, Sang Sup; et al Tetrahedron Letters (1990), 31(11), 1559-62.

Scheme 4 (1 Reaction)

Steps: 1 Yield: 100%




Suppliers (82)

Suppliers (82)

31-448-CAS-8921805 Steps: 1 Yield: 100%	A catalytic version of hypervalent aryl-λ^3-iodane-induced Hofmann rearrangement of primary carboxamides: iodobenzene as an organocatalyst and <i>m</i>-chloroperbenzoic acid as a terminal oxidant
1.1 Reagents: <i>m</i> -Chloroperbenzoic acid, Tetrafluoroboric acid Catalysts: Iodobenzene Solvents: Dichloromethane, Water; 39 h, 25 °C 1.2 Reagents: Hydrochloric acid Solvents: Water; 25 °C	By: Miyamoto, Kazunori; et al Chemical Communications (Cambridge, United Kingdom) (2012), 48(7), 982-984.
Experimental Protocols	

Scheme 5 (1 Reaction)

Steps: 1 Yield: 100%



Suppliers (41)


Suppliers (444)

Suppliers (4)

31-614-CAS-36414481	Steps: 1 Yield: 100%	Synthesis of Cyclopropylamines through an Electro-Induced Hofmann Rearrangement
1.1 Reagents: Sodium bromide Solvents: Methanol, Acetonitrile; 5 min, rt; 26 min, rt		By: Cantin, Thomas; et al
Experimental Protocols		Synthesis (2023), 55(18), 2943-2950.

Scheme 6 (1 Reaction)


Steps: 1 Yield: 100%



31-448-CAS-8741292	Steps: 1 Yield: 100%	A facile synthesis of 2-oxazolidinones via Hofmann rearrangement mediated by bis(trifluoroacetoxy)iodobenzene
1.1 Reagents: Bis(trifluoroacetoxy)iodobenzene Solvents: Acetonitrile		By: Yu, C.; et al
		Tetrahedron Letters (2001), 42(8), 1449-1452.

Scheme 7 (1 Reaction)

Steps: 1 Yield: 100%



Suppliers (15)

Suppliers (7)

31-448-CAS-13008097	Steps: 1 Yield: 100%	A facile synthesis of 2-oxazolidinones via Hofmann rearrangement mediated by bis(trifluoroacetoxy)iodobenzene
1.1 Reagents: Bis(trifluoroacetoxy)iodobenzene Solvents: Acetonitrile		By: Yu, C.; et al
		Tetrahedron Letters (2001), 42(8), 1449-1452.

Scheme 8 (7 Reactions)


Steps: 1 Yield: 96-100%



31-448-CAS-17712633 Steps: 1 Yield: 100% 1.1 Reagents: Isobutyraldehyde, Oxygen Catalysts: 1-Iodo-2,3,4,5,6-pentamethylbenzene Solvents: 1,2-Dichloroethane, 2,2-Dimethylpropanenitrile; 3 h, 40 °C 1.2 24 h, 40 °C 1.3 Solvents: Water	Iodoarene-catalyzed oxidative transformations using molecular oxygen By: Miyamoto, K.; et al Chemical Communications (Cambridge, United Kingdom) (2017), 53(70), 9781-9784.
31-448-CAS-14270820 Steps: 1 Yield: 100% 1.1 Reagents: Silver acetate, <i>N</i> -Bromophthalimide Solvents: Dimethylformamide Experimental Protocols	Hofmann rearrangement by using N-bromophthalimide-silver acetate in DMF By: Park, Min Soo; et al Archives of Pharmacal Research (1993), 16(2), 152-4.
31-448-CAS-8879523 Steps: 1 Yield: 100% 1.1 Reagents: 1,3-Dibromo-5,5-dimethylhydantoin, Mercuric oxide Solvents: Dimethylformamide Experimental Protocols	Practical Hofmann rearrangement By: Jew, Sang Sup; et al Archives of Pharmacal Research (1992), 15(4), 333-5.
31-448-CAS-4126571 Steps: 1 Yield: 98% 1.1 Reagents: Potassium peroxymonosulfate sulfate (2KHSO ₅ .KH SO ₄ .K ₂ SO ₄) Catalysts: Iodobenzene Solvents: 1,1,1,3,3,3-Hexafluoro-2-propanol, Water; 3 h, 40 °C Experimental Protocols	Hypervalent Iodine Catalyzed Hofmann Rearrangement of Carboxamides Using Oxone as Terminal Oxidant By: Yoshimura, Akira; et al Journal of Organic Chemistry (2012), 77(24), 11399-11404.
31-448-CAS-7899895 Steps: 1 Yield: 97% 1.1 Reagents: (Tosylimino)phenyliodinane Solvents: Dichloromethane; rt; 1.5 h, rt 1.2 rt; 4.5 h, rt Experimental Protocols	(Tosylimino)phenyl-λ³-iodane as a reagent for the synthesis of methyl carbamates via Hofmann rearrangement of aromatic and aliphatic carboxamides By: Yoshimura, Akira; et al Journal of Organic Chemistry (2012), 77(4), 2087-2091.
31-448-CAS-6906088 Steps: 1 Yield: 97% 1.1 Reagents: Iodobenzene, Potassium peroxymonosulfate sulfate (2KHSO ₅ .KHSO ₄ .K ₂ SO ₄); 7 - 12 h, rt Experimental Protocols	Hofmann Rearrangement of Carboxamides Mediated by Hypervalent Iodine Species Generated in Situ from Iodobenzene and Oxone: Reaction Scope and Limitations By: Zagulyaeva, Aleksandra A.; et al Organic Letters (2010), 12(20), 4644-4647.
31-614-CAS-39119958 Steps: 1 Yield: 96% 1.1 Catalysts: Sodium bromide Solvents: Acetonitrile; 5 min, rt; rt → 50 °C Experimental Protocols	Multigram Electrochemical Hofmann Rearrangement Using a Spinning Three-Dimensional Anode By: Malviya, Bhanwar K.; et al Organic Process Research & Development (2023), 27(11), 2183-2191.

Scheme 9 (1 Reaction)

Steps: 1 Yield: 100%



Suppliers (71)

Suppliers (444)

Suppliers (22)

31-448-CAS-6747554

Steps: 1 Yield: 100%

Practical Hofmann rearrangement

1.1 Reagents: 1,3-Dibromo-5,5-dimethylhydantoin, Mercuric oxide

By: Jew, Sang Sup; et al


Solvents: Dimethylformamide

Archives of Pharmacal Research (1992), 15(4), 333-5.

Experimental Protocols

Scheme 10 (1 Reaction)

Steps: 1 Yield: 100%



Suppliers (106)

Suppliers (113)

31-448-CAS-9761276

Steps: 1 Yield: 100%

Multijet Oscillating Disc Millireactor: A Novel Approach for Continuous Flow Organic Synthesis

1.1 Reagents: Sodium hydroxide, Sodium hypochlorite

By: Liguori, Lucia; et al


Solvents: 1,4-Dioxane, Water; 15 min, 80 °C

Organic Process Research & Development (2011), 15(5), 997-1009.

Experimental Protocols

Scheme 11 (2 Reactions)

Steps: 1 Yield: 99%



Suppliers (65)

Suppliers (89)

31-448-CAS-6162877

Steps: 1 Yield: 99%

Method of producing p-phenylenediamine

1.1 Reagents: Sodium hydroxide

Assignee: OAO "Kaustik"

Solvents: Water; 5 °C

Russian Federation, RU2449983 C1 2012-05-10

1.2 Reagents: Sodium hypochlorite


Solvents: Water; 5 - 7 °C; 3 h, 10 °C; < 37 °C; 0.25 h, 60 °C; 1 h, 80 °C

PatentPak available

31-448-CAS-5690709	Steps: 1 Yield: 99%	Method for production of p-phenylenediamine by Hofmann rearrangement of terephthalamide
1.1 Reagents: Sodium hydroxide, Sodium hypochlorite Solvents: Water; rt → -5 °C; 3 h, 8 °C; 0.25 h, 60 °C; 1 h, 80 °C; cooled		Assignee: OAO "Kaustik" Russian Federation, RU2378248 C1 2010-01-10 PatentPak available

Scheme 12 (3 Reactions)


Steps: 1 Yield: 98-99%



31-448-CAS-7684373	Steps: 1 Yield: 99%	Various oxidative reactions with novel ion-supported (diacetoxyiodo)benzenes
1.1 Reagents: Potassium hydroxide Solvents: Methanol; rt; 5 - 10 °C		By: Iinuma, Masataka; et al
1.2 Reagents: Iodine(1+), bis(acetato-κO)[4-[3-(1-methylpyrrolidinio)propoxy]phenyl]-, 4-methylbenzenesulfonate (1:1); 15 min, 0 °C; 1.5 h, 0 °C → rt		Tetrahedron (2013), 69(14), 2961-2970.
Experimental Protocols		
31-448-CAS-10424199	Steps: 1 Yield: 99%	Various oxidative reactions with novel ion-supported (diacetoxyiodo)benzenes
1.1 Reagents: Potassium hydroxide Solvents: Methanol; rt; 5 - 10 °C		By: Iinuma, Masataka; et al
1.2 Reagents: Iodine(1+), bis(acetato-κO)[4-[3-(trimethylammonio)propoxy]phenyl]-, 4-methylbenzenesulfonate (1:1); 15 min, 0 °C; 1.5 h, 0 °C → rt		Tetrahedron (2013), 69(14), 2961-2970.
Experimental Protocols		
31-448-CAS-14717669	Steps: 1 Yield: 98%	Hypervalent Iodine Catalyzed Hofmann Rearrangement of Carboxamides Using Oxone as Terminal Oxidant
1.1 Reagents: Potassium peroxymonosulfate sulfate (2KHSO ₅ ·KH SO ₄ ·K ₂ SO ₄) Catalysts: Iodobenzene Solvents: 1,1,1,3,3,3-Hexafluoro-2-propanol, Water; 5 - 9 h, 40 °C		By: Yoshimura, Akira; et al Journal of Organic Chemistry (2012), 77(24), 11399-11404.
Experimental Protocols		

Scheme 13 (1 Reaction)


Steps: 1 Yield: 99%



31-448-CAS-14654344	Steps: 1 Yield: 99%	The coumarin indole transformation - a method for preparing 4-halo-5-hydroxyindoles from coumarins
1.1 Reagents: Lead tetraacetate Solvents: <i>tert</i> -Butanol; 75 °C; 20 min, 75 °C		By: Stoffman, Elia J. L.; et al
Experimental Protocols		Organic & Biomolecular Chemistry (2009), 7(23), 4862-4870.

Scheme 14 (1 Reaction)

Steps: 1 Yield: 99%




Suppliers (63)

Suppliers (73)

31-448-CAS-2789868	Steps: 1 Yield: 99%	New strategies for the Hofmann reaction
No Data Available		By: Rane, Dhananjay S.; et al
		Journal of Chemical Technology and Biotechnology (1994), 59(3), 271-7.

Scheme 15 (1 Reaction)

Steps: 1 Yield: 99%




Suppliers (444)


31-448-CAS-2201497	Steps: 1 Yield: 99%	Preparation of Alkyl Carbamate of 1-Protected Indole-3-methylamine as a Precursor of Indole-3-methylamine
1.1 Reagents: Iodobenzene diacetate Solvents: Methanol; 4 h, rt		By: Song, Hao; et al
Experimental Protocols		Synthetic Communications (2005), 35(21), 2735-2748.

Scheme 16 (1 Reaction)

Steps: 1 Yield: 99%



Absolute stereochemistry shown




Suppliers (84)

Multi-component
structure image
available in CAS
SciFinder

31-350-CAS-21022629	Steps: 1 Yield: 99%	Synthesis and epimerization of phenylalanyl 4-aminocyclophosphamides
1.1 Solvents: Acetonitrile, Water; 4 h, rt		By: Jiang, Yongying; et al
Experimental Protocols		Tetrahedron (2007), 63(43), 10637-10645.

Scheme 17 (1 Reaction)

Steps: 1 Yield: 98%



Suppliers (17)

Suppliers (9)

31-448-CAS-15137569

Steps: 1 Yield: 98%

A facile synthesis of 2-oxazolidinones via Hofmann rearrangement mediated by bis(trifluoroacetoxy)iodobenzene


By: Yu, C.; et al

Tetrahedron Letters (2001), 42(8), 1449-1452.

1.1 **Reagents:** Bis(trifluoroacetoxy)iodobenzene
Solvents: Acetonitrile

Scheme 18 (1 Reaction)

Steps: 1 Yield: 98%



Suppliers (54)

Suppliers (79)

31-448-CAS-14152466

Steps: 1 Yield: 98%

Synthesis of 4-amino-4-methylpiperidine di-p-toluenesulfonate

By: Tan, Hongmei; et al


Guangdong Yaoxueyuan Xuebao (2011), 27(3), 247-249.

1.1 **Reagents:** Potassium hydroxide
Solvents: Acetonitrile, Water; 0 - 5 °C; 5 °C → 0 °C

1.2 **Reagents:** 1,3-Dibromo-5,5-dimethylhydantoin; 0 - 5 °C; 30 min, 0 - 5 °C; 1 h, rt

Scheme 19 (1 Reaction)

Steps: 1 Yield: 98%



31-448-CAS-16117245

Steps: 1 Yield: 98%

Preparation method of rivaroxaban


1.1 **Reagents:** Bis(trifluoroacetoxy)iodobenzene
Solvents: Dimethylformamide; rt; 2 h, rt

Assignee: Shandong Luoxin Pharmaceutical Group Co., Ltd.
 China, CN105801572 A 2016-07-27

PatentPak available

Scheme 20 (1 Reaction)

Steps: 1 Yield: 98%



Suppliers (71)

Suppliers (65)

31-614-CAS-40880329

Steps: 1 Yield: 98%

Micelle-Enabled Hofmann Rearrangement in Water

- 1.1 **Reagents:** Bis(trifluoroacetoxy)iodobenzene
Solvents: Water, Poly(oxy-1,2-ethanediyl), α -[[[(2,5)-1-(1-oxododecyl)-2-pyrrolidinyl]carbonyl]- ω -methoxy-; 12 h, rt
- 1.2 **Reagents:** Hydrochloric acid
Solvents: Water; 30 min

By: Xing, Xiaoqi; et al

Organic Process Research & Development (2024), 28(7), 2945-2950.

Experimental Protocols

Scheme 21 (1 Reaction)

Steps: 1 Yield: 98%



Suppliers (61)

Suppliers (92)

31-448-CAS-1409691

Steps: 1 Yield: 98%

Preliminary study on seawater instead of freshwater in organic chemical reaction


- 1.1 **Reagents:** Sodium hydroxide, Bromine
Solvents: Water; 0 - 5 °C; 20 min, 0 - 5 °C
- 1.2 1.5 h, 0 - 5 °C; 5 °C $\rightarrow$ rt; 2 h, rt $\rightarrow$ 65 °C; 2 h, 65 °C $\rightarrow$ 75 °C

By: Gang, Liu; et al

Huaxue Tongbao (2012), 75(9), 832-836.

Scheme 22 (2 Reactions)

Steps: 1 Yield: 97-98%



Double bond geometry shown

Suppliers (444)

Double bond geometry shown

Suppliers (65)

Suppliers (4)

31-448-CAS-1504869

Steps: 1 Yield: 98%

Preparation and Reactivities of Novel (Diacetoxyiodo)arenes Bearing Heteroaromatics

- 1.1 **Reagents:** Potassium hydroxide, Bis(acetato- κO)-3-thienyl iodine
Solvents: Methanol

By: Togo, Hideo; et al

Journal of Organic Chemistry (2000), 65(24), 8391-8394.

Experimental Protocols

31-448-CAS-12527360	Steps: 1 Yield: 97%	Various oxidative reactions with novel ion-supported (diacetoxyiodo)benzenes
1.1 Reagents: Potassium hydroxide Solvents: Methanol; rt; 5 - 10 °C		By: Iinuma, Masataka; et al
1.2 Reagents: Iodine(1+), bis(acetato-κO)[4-[3-(trimethylammonio)propoxy]phenyl]-, 4-methylbenzenesulfonate (1:1); 15 min, 0 °C; 1.5 h, 0 °C → rt		Tetrahedron (2013), 69(14), 2961-2970.
Experimental Protocols		

Scheme 23 (1 Reaction)	Steps: 1 Yield: 98%
Absolute stereochemistry shown, Rotation (+) Suppliers (1)	Absolute stereochemistry shown Suppliers (2)

31-448-CAS-123665	Steps: 1 Yield: 98%	Enantioselective biotransformation of α,α-disubstituted dinitriles to the corresponding 2-cyanoacetamides using <i>Rhodococcus</i> sp. CGMCC 0497
1.1 Reagents: <i>N</i> -Bromosuccinimide, Mercuric acetate Solvents: Dimethylformamide; 16 h, rt		By: Wu, Zhong-Liu; et al
1.2 Reagents: Water		Tetrahedron: Asymmetry (2003), 14(15), 2133-2142.

Scheme 24 (1 Reaction)	Steps: 1 Yield: 98%
Suppliers (7)	Suppliers (77)

31-448-CAS-14152626	Steps: 1 Yield: 98%	Method for the production of racemic tert-leucine and its homologs or their nitriles
1.1 Reagents: Sodium hydroxide, Sodium hypochlorite		Assignee: Huels A.-G.
		Germany, DE19724086 A1 1997-12-11
		PatentPak available

Scheme 25 (1 Reaction)	Steps: 1 Yield: 98%
Suppliers (4)	Suppliers (18)

31-448-CAS-4778963	Steps: 1 Yield: 98%	Preparation of amino acid esters substituted at the alpha position by a tertiary hydrocarbon chain
1.1 Reagents: Sodium hypochlorite Solvents: Water		Assignee: Degussa-Huls Aktiengesellschaft
Experimental Protocols		European Patent Organization, EP970944 A1 2000-01-12 PatentPak available

Scheme 26 (2 Reactions)

Steps: 1 Yield: 96-97%




31-448-CAS-182442	Steps: 1 Yield: 97%	(Tosylimino)phenyl- λ^3 -iodane as a reagent for the synthesis of methyl carbamates via Hofmann rearrangement of aromatic and aliphatic carboxamides
1.1 Reagents: (Tosylimino)phenyliodine Solvents: Dichloromethane; rt; 1.5 h, rt		By: Yoshimura, Akira; et al
1.2 rt; 4.5 h, rt		Journal of Organic Chemistry (2012), 77(4), 2087-2091.
Experimental Protocols		

31-448-CAS-12913265	Steps: 1 Yield: 96%	Hypervalent Iodine Catalyzed Hofmann Rearrangement of Carboxamides Using Oxone as Terminal Oxidant
1.1 Reagents: Potassium peroxymonosulfate sulfate (2KHSO ₅ .KH SO ₄ .K ₂ SO ₄) Catalysts: Iodobenzene Solvents: 1,1,1,3,3,3-Hexafluoro-2-propanol, Water; 5 - 9 h, 40 °C		By: Yoshimura, Akira; et al
Experimental Protocols		Journal of Organic Chemistry (2012), 77(24), 11399-11404.

Scheme 27 (1 Reaction)


Steps: 1 Yield: 97%



31-448-CAS-10870790	Steps: 1 Yield: 97%	A facile synthesis of 2-oxazolidinones via Hofmann rearrangement mediated by bis(trifluoroacetoxy)iodobenzene
1.1 Reagents: Bis(trifluoroacetoxy)iodobenzene Solvents: Acetonitrile		By: Yu, C.; et al
		Tetrahedron Letters (2001), 42(8), 1449-1452.

Scheme 28 (1 Reaction)

Steps: 1 Yield: 97%



Suppliers (158)

Suppliers (70)

31-448-CAS-21178451

Steps: 1 Yield: 97%

Synthesis of 2,3-dichloropyridine


1.1 **Reagents:** Sodium hydroxide, Sodium hypochlorite
Solvents: Water; 10 °C; < 10 °C; 0 °C → 90 °C

By: Liu, Wei; et al

Huaxue Shiji (2015), 37(3), 283-285, 288.

Scheme 29 (1 Reaction)

Steps: 1 Yield: 97%



Suppliers (54)

Suppliers (2)

31-614-CAS-40880325

Steps: 1 Yield: 97%

Micelle-Enabled Hofmann Rearrangement in Water

1.1 **Reagents:** Bis(trifluoroacetoxy)iodobenzene
Solvents: Water, Poly(oxy-1,2-ethanediyl), α-[[[(2S)-1-(1-oxododecyl)-2-pyrrolidinyl]carbonyl]-ω-methoxy-; 12 h, rt

By: Xing, Xiaoqi; et al


Organic Process Research & Development (2024), 28(7), 2945-2950.

1.2 **Reagents:** Hydrochloric acid
Solvents: Water; 30 min

Experimental Protocols

Scheme 30 (1 Reaction)

Steps: 1 Yield: 97%



Suppliers (75)

Suppliers (75)

31-448-CAS-2524801

Steps: 1 Yield: 97%

New strategies for the Hofmann reaction


No Data Available

By: Rane, Dhananjay S.; et al

Journal of Chemical Technology and Biotechnology (1994), 59(3), 271-7.

Scheme 31 (1 Reaction)

Steps: 1 Yield: 97%



Suppliers (40)

Suppliers (444)

31-448-CAS-59930

Steps: 1 Yield: 97%

Preparation of Alkyl Carbamate of 1-Protected Indole-3-methylamine as a Precursor of Indole-3-methylamine

1.1 Reagents: Iodobenzene diacetate
Solvents: Methanol; 4 h, rt


By: Song, Hao; et al

Experimental Protocols

Synthetic Communications (2005), 35(21), 2735-2748.

Scheme 32 (1 Reaction)

Steps: 1 Yield: 97%



Absolute stereochemistry shown

Absolute stereochemistry shown

Supplier (1)

Suppliers (7)

31-448-CAS-10756501

Steps: 1 Yield: 97%

Method for producing optically active 3-aminopiperidine or salt thereof

1.1 Reagents: Sodium hydroxide, Sodium hypochlorite
Solvents: Water; 6 h, 25 °C

Assignee: Kaneka Corporation


Experimental Protocols

World Intellectual Property Organization, WO2008102720 A1 2008-08-28

[PatentPak](#) available

Scheme 33 (1 Reaction)

Steps: 1 Yield: 97%



Suppliers (83)

Suppliers (6)

31-448-CAS-18293796

Steps: 1 Yield: 97%

Method for synthesizing alogliptin intermediate (R)-3-aminopiperidine dihydrochloride

1.1 Reagents: 1-Fluoronaphthalene, Hydrogen peroxide, Tetrafluoroboric acid
Solvents: Ethanol, Water; 0.5 h, rt

Assignees: Weihai Disu Pharmaceutical Co., Ltd.; Disha Pharmaceutical Group Co., Ltd.; Shandong Disha Pharmaceutical Co., Ltd., Disha Pharmaceutical Group

1.2 8 h, rt


China, CN103319399 A 2013-09-25

1.3 Reagents: Hydrochloric acid
Solvents: Ethyl acetate, Water; 0.5 h, rt

[PatentPak](#) available

Scheme 34 (1 Reaction)

Steps: 1 Yield: 97%



Absolute stereochemistry shown

Absolute stereochemistry shown

Supplier (1)

Suppliers (99)

Suppliers (17)

31-448-CAS-9023821

Steps: 1 Yield: 97%

Synthesis of (s)-(+)-3-(aminomethyl)-5-methylhexanoic acid

1.1 Reagents: Bromine

Solvents: Methanol; rt → 5 °C; -25 °C → -15 °C; -25 °C → -15 °C; 1 - 2 h, -25 °C → -15 °C; -15 °C → 0 °C; 0 °C → 65 °C; 1 - 2 h

Assignees: Teva Pharmaceutical Industries Ltd.; Teva Pharmaceuticals USA, Inc.


Experimental Protocols

World Intellectual Property Organization, WO2008118427 A2 2008-10-02

PatentPak available

Scheme 35 (1 Reaction)

Steps: 1 Yield: 96%



Suppliers (26)

Suppliers (444)

Suppliers (3)

31-448-CAS-6256068

Steps: 1 Yield: 96%

Hypervalent Iodine Catalyzed Hofmann Rearrangement of Carboxamides Using Oxone as Terminal Oxidant

1.1 Reagents: Potassium peroxydisulfate sulfate (2KHSO₅·KH SO₄·K₂SO₄)

Catalysts: Iodobenzene

Solvents: 1,1,1,3,3,3-Hexafluoro-2-propanol, Water; 5 - 9 h, 40 °C

By: Yoshimura, Akira; et al

Journal of Organic Chemistry (2012), 77(24), 11399-11404.

Experimental Protocols

Scheme 36 (1 Reaction)

Steps: 1 Yield: 96%



Suppliers (54)

Suppliers (444)

Suppliers (5)

31-448-CAS-17712629

Steps: 1 Yield: 96%

Iodoarene-catalyzed oxidative transformations using molecular oxygen

1.1 Reagents: Isobutyraldehyde, Oxygen

Catalysts: 1-Iodo-2,3,4,5,6-pentamethylbenzene

Solvents: 1,2-Dichloroethane, 2,2-Dimethylpropanenitrile; 3 h, 40 °C

By: Miyamoto, K.; et al

Chemical Communications (Cambridge, United Kingdom) (2017), 53(70), 9781-9784.

1.2 24 h, 40 °C

1.3 Solvents: Water

Copyright © 2025 American Chemical Society (ACS). All Rights Reserved.

Internal use only. Redistribution is subject to the terms of your CAS SciFinder License Agreement and CAS information Use Policies.