

Initiating Search

November 10, 2025, 2:43 PM

• Search:

CAS Lexicon Search:

3-Chloroperbenzoic acid - Preferred Concept

Search Tasks

Task	Result Type	View
Returned Reference Results from CAS Lexicon (8,912)	 References	View Results
Exported: Retrieved Related Reaction Results + Filters (1,235)	 Reactions	View Results
Filtered By:		
Yield:	90-100%, 80-89%, 70-79%	
Document Type:	Journal	

Copyright © 2025 American Chemical Society (ACS). All Rights Reserved.


Internal use only. Redistribution is subject to the terms of your CAS SciFinder License Agreement and CAS information Use Policies.

Reactions (50)

[View in CAS SciFinder](#)

Scheme 1 (1 Reaction)

Steps: 1 Yield: 100%



Relative stereochemistry shown

 Supplier (1)

31-279-CAS-7274453

Steps: 1 Yield: 100%

Dioxirane oxidations of compounds other than alkenes


1.1 Reagents: Dimethyldioxirane
Solvents: Acetone

By: Adam, Waldemar; et al

Organic Reactions (Hoboken, NJ, United States) (2007), 69, 1-346.

Scheme 2 (1 Reaction)

Steps: 1 Yield: 100%



Absolute stereochemistry shown

 Suppliers (3)

 Suppliers (2)

31-279-CAS-6980510

Steps: 1 Yield: 100%

Is hole transfer involved in metalloporphyrin-catalyzed epoxidation?


1.1 Reagents: Nitrogen, Phenylseleninyl benzeneseleninate
Catalysts: Antimonate(1-), hexachloro-, (OC-6-11)-, salt with 4-bromo-*N,N*-bis(4-bromophenyl)benzenamine (1:1)
Solvents: Dichloromethane

By: Kim, Taisun; et al

Journal of the American Chemical Society (1993), 115(17), 7653-64.

Scheme 3 (1 Reaction)

Steps: 1 Yield: 100%



31-279-CAS-13514403	Steps: 1 Yield: 100%	Effects of alkyl side chains and terminal hydroph ilicity on vitamin D receptor (VDR) agonistic activity based on the diphenylpentane skeleton
1.1 Reagents: <i>m</i> -Chloroperbenzoic acid Solvents: Dichloromethane; rt		By: Misawa, Takashi; et al Bioorganic & Medicinal Chemistry Letters (2015), 25(22), 5362-5366.

Scheme 4 (1 Reaction)


Steps: 1 Yield: 100%

 Supplier (1)

31-279-CAS-2788699	Steps: 1 Yield: 100%	Dioxirane oxidations of compounds other than alkenes
1.1 Reagents: Dimethyldioxirane Solvents: Acetone, Dichloromethane		By: Adam, Waldemar; et al Organic Reactions (Hoboken, NJ, United States) (2007), 69, 1-346.

Scheme 5 (1 Reaction)

Steps: 1 Yield: 100%




Relative stereochemistry shown

 Suppliers (6) Supplier (1)

31-279-CAS-9915473	Steps: 1 Yield: 100%	Diastereoselective Catalytic Epoxidation of Chiral Allylic Alcohols by the TS-1 and Ti-β Zeolites: Evidence for a Hydrogen-Bonded, Peroxy-Type Loaded Complex as Oxidizing Species
1.1 Reagents: <i>tert</i> -Butyl hydroperoxide Catalysts: Titanium isopropoxide Solvents: Chloroform- <i>d</i>		By: Adam, Waldemar; et al Journal of Organic Chemistry (1997), 62(11), 3631-3637.

Scheme 6 (1 Reaction)


Steps: 1 Yield: 100%

 Suppliers (15)

31-279-CAS-13879710	Steps: 1 Yield: 100% Dioxirane oxidations of compounds other than alkenes
1.1 Reagents: Dimethyldioxirane Solvents: Acetone, Dichloromethane	By: Adam, Waldemar; et al Organic Reactions (Hoboken, NJ, United States) (2007), 69, 1-346.

Scheme 7 (1 Reaction)

Steps: 1 Yield: 100%




Suppliers (3)

31-279-CAS-11088396	Steps: 1 Yield: 100% Dioxirane oxidations of compounds other than alkenes
1.1 Reagents: Dimethyldioxirane Solvents: Acetone	By: Adam, Waldemar; et al Organic Reactions (Hoboken, NJ, United States) (2007), 69, 1-346.

Scheme 8 (1 Reaction)

Steps: 1 Yield: 100%




Absolute stereochemistry shown

Absolute stereochemistry shown

31-279-CAS-15348267	Steps: 1 Yield: 100% Synthesis of C-disaccharides via a hetero-Diels-Alder reaction and further stereocontrolled transformations
1.1 Reagents: Dimethyldioxirane Solvents: Acetone, Dichloromethane; 5 h, 0 °C	By: Wan, Qian; et al
Experimental Protocols	Carbohydrate Research (2008), 343(10-11), 1754-1765.

Scheme 9 (1 Reaction)

Steps: 1 Yield: 100%




Suppliers (5)

Suppliers (3)

31-279-CAS-11425816	Steps: 1 Yield: 100% Dioxirane oxidations of compounds other than alkenes
1.1 Reagents: Dimethyldioxirane Solvents: Acetone	By: Adam, Waldemar; et al Organic Reactions (Hoboken, NJ, United States) (2007), 69, 1-346.

Scheme 10 (1 Reaction)

Steps: 1 Yield: 100%



Double bond geometry shown

31-279-CAS-22654875

Steps: 1 Yield: 100%

Access to Vinyl Ethers and Ketones with Hypervalent Iodine Reagents as Oxy-Allyl Cation Synthetic Equivalents

By: Declas, Nina; et al

Angewandte Chemie, International Edition (2020), 59(41), 18256-18260.

1.1 **Reagents:** Sodium bicarbonate, *m*-Chloroperbenzoic acid
Solvents: Dichloromethane; 5 h, rt

1.2 **Reagents:** Sodium bicarbonate
Solvents: Water; 10 min, rt

Experimental Protocols

Scheme 11 (2 Reactions)

Steps: 1 Yield: 100%

Multi-component
structure image
available in CAS
SciFinder



Multi-component
structure image
available in CAS
SciFinder

31-279-CAS-20672241

Steps: 1 Yield: 100%

Antibacterial surface based on new epoxy-amine networks from ionic liquid monomers

By: Livi, Sebastien; et al

European Polymer Journal (2019), 116, 56-64.

1.1 **Reagents:** Dimethyldioxirane
Solvents: Acetone; 12 h, rt

1.2 **Reagents:** Dimethyl sulfide

Experimental Protocols

31-279-CAS-17952133

Steps: 1 Yield: 100%

Dimethyldioxirane (DMDO) as a valuable oxidant for the synthesis of polyfunctional aromatic imidazolium monomers bearing epoxides

By: Chardin, C.; et al

Green Chemistry (2017), 19(21), 5054-5059.


1.1 **Reagents:** Dimethyldioxirane
Solvents: Acetone; 12 h, rt

1.2 **Reagents:** Dimethyl sulfide; rt

Experimental Protocols

Scheme 12 (1 Reaction)

Steps: 1 Yield: 100%



Suppliers (6)

Suppliers (2)

31-279-CAS-7580209

Steps: 1 Yield: 100%

Dioxirane oxidations of compounds other than alkenes


By: Adam, Waldemar; et al

Organic Reactions (Hoboken, NJ, United States) (2007), 69, 1-346.

1.1 **Reagents:** Dimethyldioxirane
Solvents: Acetone, Dichloromethane

Scheme 13 (1 Reaction)

Steps: 1 Yield: 100%



Suppliers (63)

31-279-CAS-2029753

Steps: 1 Yield: 100%

Dioxirane oxidations of compounds other than alkenes


1.1 Reagents: Dimethyldioxirane
Solvents: Acetone, Dichloromethane

By: Adam, Waldemar; et al

Organic Reactions (Hoboken, NJ, United States) (2007), 69, 1-346.

Scheme 14 (1 Reaction)

Steps: 1 Yield: 100%



Supplier (1)

31-279-CAS-16877599

Steps: 1 Yield: 100%

1-Heteroarylpropan-2-ones as inhibitors of fatty acid amide hydrolase: Studies on structure-activity relationships and metabolic stability

1.1 Reagents: *m*-Chloroperbenzoic acid
Solvents: Dichloromethane; 6 d, rt


By: Zahov, Stefan; et al

Bioorganic & Medicinal Chemistry (2017), 25(3), 825-837.

Experimental Protocols

Scheme 15 (1 Reaction)

Steps: 1 Yield: 100%



31-279-CAS-8079338

Steps: 1 Yield: 100%

Synthesis of the originally proposed structures of elatenyne and an enyne from *Laurencia majuscula*

1.1 Reagents: Sodium bicarbonate, Dimethyldioxirane
Solvents: Dichloromethane; 0 °C; 20 min, 0 °C


By: Sheldrake, Helen M.; et al

Organic & Biomolecular Chemistry (2009), 7(2), 238-252.

Experimental Protocols

Scheme 16 (1 Reaction)

Steps: 1 Yield: 100%



Double bond geometry shown

31-279-CAS-22654876

Steps: 1 Yield: 100%

Access to Vinyl Ethers and Ketones with Hypervalent Iodine Reagents as Oxy-Allyl Cation Synthetic Equivalents

By: Declas, Nina; et al

Angewandte Chemie, International Edition (2020), 59(41), 18256-18260.


1.1 **Reagents:** Sodium bicarbonate, *m*-Chloroperbenzoic acid
Solvents: Dichloromethane; 5 h, rt

1.2 **Reagents:** Sodium bicarbonate
Solvents: Water; 10 min, rt

Experimental Protocols

Scheme 17 (1 Reaction)

Steps: 1 Yield: 100%



Relative stereochemistry shown

Supplier (1)

31-279-CAS-9063511

Steps: 1 Yield: 100%

Dioxirane oxidations of compounds other than alkenes

By: Adam, Waldemar; et al

Organic Reactions (Hoboken, NJ, United States) (2007), 69, 1-346.

1.1 **Reagents:** Dimethyldioxirane
Solvents: Acetone

Scheme 18 (1 Reaction)

Steps: 1 Yield: 100%

Multi-component
structure image
available in CAS
SciFinder



Multi-component
structure image
available in CAS
SciFinder

31-279-CAS-20672240

Steps: 1 Yield: 100%

Antibacterial surface based on new epoxy-amine networks from ionic liquid monomers

By: Livi, Sebastien; et al

European Polymer Journal (2019), 116, 56-64.


1.1 **Reagents:** Dimethyldioxirane
Solvents: Acetone; 2 h, rt

1.2 **Reagents:** Dimethyl sulfide

Experimental Protocols

Scheme 19 (1 Reaction)

Steps: 1 Yield: 100%



Suppliers (6)

Supplier (1)

31-279-CAS-12430836

Steps: 1 Yield: 100%

Structure-Activity Relationship Studies on Derivatives of Eudesmanolides from *Inula helenium* as Toxicants against *Aedes aegypti* Larvae and Adults

1.1 **Reagents:** *m*-Chloroperbenzoic acid
Solvents: Dichloromethane; overnight, cooled


Experimental Protocols

By: Cantrell, Charles L.; et al

Chemistry & Biodiversity (2010), 7(7), 1681-1697.

Scheme 20 (1 Reaction)

Steps: 1 Yield: 100%



Suppliers (7)

Suppliers (3)

31-279-CAS-1107138

Steps: 1 Yield: 100%

Diastereoselective Catalytic Epoxidation of Chiral Allylic Alcohols by the TS-1 and Ti-β Zeolites: Evidence for a Hydrogen-Bonded, Peroxy-Type Loaded Complex as Oxidizing Species


1.1 **Reagents:** *tert*-Butyl hydroperoxide
Catalysts: Titanium isopropoxide
Solvents: Chloroform-*d*

By: Adam, Waldemar; et al

Journal of Organic Chemistry (1997), 62(11), 3631-3637.

Scheme 21 (1 Reaction)

Steps: 1 Yield: 100%




Suppliers (34)

Suppliers (4)

31-279-CAS-12368577 Steps: 1 Yield: 100%	Diastereoselective Catalytic Epoxidation of Chiral Allylic Alcohols by the TS-1 and Ti-β Zeolites: Evidence for a Hydrogen-Bonded, Peroxy-Type Loaded Complex as Oxidizing Species By: Adam, Waldemar; et al Journal of Organic Chemistry (1997), 62(11), 3631-3637.
1.1 Reagents: Hydrogen peroxide Catalysts: Titanium Solvents: Acetonitrile, Water	

Scheme 22 (1 Reaction)

Steps: 1 Yield: 100%




Supplier (1)

Suppliers (3)

31-279-CAS-12991764 Steps: 1 Yield: 100%	Dioxirane oxidations of compounds other than alkenes By: Adam, Waldemar; et al Organic Reactions (Hoboken, NJ, United States) (2007), 69, 1-346.
1.1 Reagents: Potassium carbonate, Dimethyldioxirane Solvents: Acetone	

Scheme 23 (1 Reaction)

Steps: 1 Yield: 100%




Suppliers (8)

Suppliers (3)

31-279-CAS-6064212 Steps: 1 Yield: 100%	Quasi-biomimetic ring contraction promoted by a cysteine-based nucleophile: Total synthesis of Sch-642305, some analogs and their putative anti-HIV activities By: Dermenci, Alpay; et al Chemical Science (2011), 2(8), 1568-1572.
1.1 Reagents: Sodium bicarbonate, <i>m</i> -Chloroperbenzoic acid Solvents: Dichloromethane; cooled; overnight	
Experimental Protocols	

Scheme 24 (1 Reaction)

Steps: 1 Yield: 100%



Relative stereochemistry shown


Suppliers (2)

Supplier (1)

31-279-CAS-132481	Steps: 1 Yield: 100%	Dioxirane oxidations of compounds other than alkenes
1.1 Reagents: Dimethyldioxirane Solvents: Acetone		By: Adam, Waldemar; et al Organic Reactions (Hoboken, NJ, United States) (2007), 69, 1-346.

Scheme 25 (1 Reaction)

Steps: 1 Yield: 100%



Suppliers (8)

Suppliers (4)

31-279-CAS-9826363	Steps: 1 Yield: 100%	Dioxirane oxidations of compounds other than alkenes
1.1 Reagents: Dimethyldioxirane Solvents: Acetone, Dichloromethane		By: Adam, Waldemar; et al Organic Reactions (Hoboken, NJ, United States) (2007), 69, 1-346.

Scheme 26 (1 Reaction)

Steps: 1 Yield: 100%

Multi-component
structure image
available in CAS
SciFinder




Multi-component
structure image
available in CAS
SciFinder

31-279-CAS-17952134	Steps: 1 Yield: 100%	Dimethyldioxirane (DMDO) as a valuable oxidant for the synthesis of polyfunctional aromatic imidazolium monomers bearing epoxides
1.1 Reagents: Dimethyldioxirane Solvents: Acetone; 12 h, rt		By: Chardin, C.; et al
1.2 Reagents: Dimethyl sulfide; rt		Green Chemistry (2017), 19(21), 5054-5059.
Experimental Protocols		

Scheme 27 (1 Reaction)

Steps: 1 Yield: 100%



Double bond geometry shown

Relative stereochemistry shown


Suppliers (3)

Supplier (1)

31-279-CAS-472473 Steps: 1 Yield: 100%	Diastereoselective Catalytic Epoxidation of Chiral Allylic Alcohols by the TS-1 and Ti-β Zeolites: Evidence for a Hydrogen-Bonded, Peroxy-Type Loaded Complex as Oxidizing Species By: Adam, Waldemar; et al Journal of Organic Chemistry (1997), 62(11), 3631-3637.
1.1 Reagents: <i>m</i> -Chloroperbenzoic acid Solvents: Dichloromethane	

Scheme 28 (1 Reaction)

Steps: 1 Yield: 100%




Supplier (1)

31-279-CAS-7585873 Steps: 1 Yield: 100%	Stereoselective functionalization of SuperQuat enamides: asymmetric synthesis of homochiral 1,2-diols and α-benzyloxy carbonyl compounds By: Aciro, Caroline; et al Tetrahedron (2008), 64(39), 9320-9344.
1.1 Reagents: Dimethyldioxirane Solvents: Acetone; 0 °C; 1 h, 0 °C → rt	
Experimental Protocols	

Scheme 29 (1 Reaction)

Steps: 1 Yield: 100%




Suppliers (70)

Suppliers (6)

31-279-CAS-21167692 Steps: 1 Yield: 100%	A Chemical Probe for the ATAD2 Bromodomain By: Bamborough, Paul; et al Angewandte Chemie, International Edition (2016), 55(38), 11382-11386.
1.1 Reagents: <i>m</i> -Chloroperbenzoic acid Solvents: Dichloromethane, Water; 0 °C; 15 min, 0 °C → rt; 16 h, rt 1.2 Reagents: Sodium thiosulfate Solvents: Water; 15 min, rt	
Experimental Protocols	

Scheme 30 (1 Reaction)

Steps: 1 Yield: 100%



Suppliers (34)

Suppliers (5)

31-279-CAS-3407690

Steps: 1 Yield: 100%

Diastereoselective Catalytic Epoxidation of Chiral Allylic Alcohols by the TS-1 and Ti-β Zeolites: Evidence for a Hydrogen-Bonded, Peroxy-Type Loaded Complex as Oxidizing Species

By: Adam, Waldemar; et al

Journal of Organic Chemistry (1997), 62(11), 3631-3637.

1.1 **Reagents:** *tert*-Butyl hydroperoxide
Catalysts: Vanadyl acetylacetonate
Solvents: Benzene

Scheme 31 (1 Reaction)

Steps: 1 Yield: 100%

Multi-component
structure image
available in CAS
SciFinder



Multi-component
structure image
available in CAS
SciFinder

31-279-CAS-17952121

Steps: 1 Yield: 100%

Dimethyldioxirane (DMDO) as a valuable oxidant for the synthesis of polyfunctional aromatic imidazolium monomers bearing epoxides

By: Chardin, C.; et al


Green Chemistry (2017), 19(21), 5054-5059.

1.1 **Reagents:** Dimethyldioxirane
Solvents: Acetone; 12 h, rt
 1.2 **Reagents:** Dimethyl sulfide; rt

Experimental Protocols

Scheme 32 (1 Reaction)

Steps: 1 Yield: 100%



Suppliers (8)

Suppliers (63)

31-014-CAS-9442975

Steps: 1 Yield: 100%

Ammonium-directed dihydroxylation of 3-aminocyclohex-1-enes: development of a metal-free dihydroxylation protocol

By: Aciro, Caroline; et al


Organic & Biomolecular Chemistry (2008), 6(20), 3751-3761.

1.1 **Solvents:** Dichloromethane; 5 min, rt
 1.2 **Reagents:** *m*-Chloroperbenzoic acid; 21 h, rt
 1.3 **Reagents:** Sodium sulfite
Solvents: Water; rt

Experimental Protocols

Scheme 33 (1 Reaction)

Steps: 1 Yield: 100%



Suppliers (81)

Suppliers (6)

31-279-CAS-560111

Steps: 1 Yield: 100%

Anti-proliferative effect of synthesized bakuchiol analogues on cultured human tumor cell lines

By: Cha, Mi-Ran; et al

Bulletin of the Korean Chemical Society (2012), 33(7), 2378-2380.

1.1 Reagents: *m*-Chloroperbenzoic acid

Solvents: Dichloromethane; 6 h, 0 °C → rt

Scheme 34 (1 Reaction)

Steps: 1 Yield: 100%

Multi-component
structure image
available in CAS
SciFinder



Multi-component
structure image
available in CAS
SciFinder

31-279-CAS-22401650

Steps: 1 Yield: 100%

New Epoxy Thermosets Derived from a Bisimidazolium Ionic Liquid Monomer: An Experimental and Modeling Investigation

By: Radchenko, Alexei V.; et al

ACS Sustainable Chemistry & Engineering (2020), 8(32), 12208-12221.


1.1 Reagents: *m*-Chloroperbenzoic acid

Solvents: Acetonitrile; 24 h, 40 °C

Experimental Protocols

Scheme 35 (1 Reaction)

Steps: 1 Yield: 100%



Double bond geometry shown

Relative stereochemistry shown

Suppliers (56)

Suppliers (3)

31-279-CAS-12626396

Steps: 1 Yield: 100%

Terminal and Internal Olefin Epoxidation with Cobalt(II) as the Catalyst: Evidence for an Active Oxidant Co^{II}-Acylperoxo Species

By: Hyun, Min Young; et al

Journal of Organic Chemistry (2012), 77(17), 7307-7312.

1.1 Reagents: *m*-Chloroperbenzoic acid


Catalysts: Cobalt(II) perchlorate

Solvents: Acetonitrile; 10 min, rt

Experimental Protocols

Scheme 36 (1 Reaction)

Steps: 1 Yield: 100%



Suppliers (66)

Suppliers (69)

31-279-CAS-19004336

Steps: 1 Yield: 100%

- 1.1 Reagents: *m*-Chloroperbenzoic acid
 Catalysts: (SP-4-1)-[5,10,15,20-Tetrakis(2,6-dibutoxyphenyl)-21*H*,23*H*-porphinato(2-)-κ N^{21} ,κ N^{22} ,κ N^{23} ,κ N^{24}]nickel
 Solvents: Dichloromethane; 10 min, rt

Experimental Protocols


Synthesis, Characterization, and Efficient Catalytic Activities of a Nickel(II) Porphyrin: Remarkable Solvent and Substrate Effects on Participation of Multiple Active Oxidants

By: Ahn, Hye Mi; et al

Chemistry - A European Journal (2017), 23(49), 11969-11976.

Scheme 37 (1 Reaction)

Steps: 1 Yield: 100%



Suppliers (48)

Suppliers (81)

31-279-CAS-2320118

Steps: 1 Yield: 100%

- 1.1 Reagents: Hydrogen peroxide
 Catalysts: Ruthenium, chloro(2-quinolinecarboxylato-κ N^1 ,κ O^2)-(2,2':6',2''-terpyridine-κ N^1 ,κ $N^{1'}$,κ $N^{1''}$)-, (OC-6-24)-
 Solvents: Ethanol; 10 min, 298 K; 8 h, 298 K

Experimental Protocols


Isomeric Complexes of [Ru^{II}(trpy)(L)Cl] (trpy = 2,2':6',2''-Terpyridine and HL = Quinaldic Acid): Preference of Isomeric Structural Form in Catalytic Chemoselective Epoxidation Process

By: Dutta Chowdhury, Abhishek; et al

Inorganic Chemistry (2011), 50(5), 1775-1785.

Scheme 38 (1 Reaction)

Steps: 1 Yield: 100%



Suppliers (93)

Suppliers (9)

31-279-CAS-10686318

Steps: 1 Yield: 100%

- 1.1 Solvents: Chloroform; rt → 0 °C
 1.2 Reagents: *m*-Chloroperbenzoic acid; 0 °C; 0 °C → rt; rt
 1.3 Reagents: Sodium bicarbonate
 Solvents: Water; rt


The potential biologically active epoxide, hydroperoxide and endoperoxide derivatives derived from natural monoterpene β-myrcene

By: Elgendy, Eman M.

Australian Journal of Basic and Applied Sciences (2008), 2(2), 221-224.

Scheme 39 (5 Reactions)

Steps: 1 Yield: 100%



Suppliers (79)

Suppliers (70)

31-279-CAS-12308748	Steps: 1 Yield: 100%	Catalytic asymmetric epoxidation of unfunctionalized olefins using silica, LDH and resin-supported sulfonate-Mn(salen) complex By: Choudary, B. M.; et al Advanced Synthesis & Catalysis (2006), 348(4 + 5), 493-498.
1.1 Reagents: 4-Methylmorpholine, <i>m</i> -Chloroperbenzoic acid Catalysts: Manganate(2-), [[3,3'-[1,2-cyclohexanediylbis[[nitrilo-κM)methylidyne]]bis[5-(1,1-dimethylethyl)-4-(hydroxy-κO)benzenesulfonato]](4-)](2,4-pentanedionato-κO ² ,κO ⁴)-, sodium (1:2) Solvents: Acetonitrile; < 5 min, -20 °C		
31-279-CAS-8044636	Steps: 1 Yield: 100%	Catalytic asymmetric epoxidation of unfunctionalized olefins using silica, LDH and resin-supported sulfonate-Mn(salen) complex By: Choudary, B. M.; et al Advanced Synthesis & Catalysis (2006), 348(4 + 5), 493-498.
1.1 Reagents: 4-Methylmorpholine, <i>m</i> -Chloroperbenzoic acid Catalysts: Manganate(2-), [[3,3'-[1,2-cyclohexanediylbis[[nitrilo-κM)methylidyne]]bis[5-(1,1-dimethylethyl)-4-(hydroxy-κO)benzenesulfonato]](4-)](2,4-pentanedionato-κO ² ,κO ⁴)-, sodium (1:2) Solvents: Acetonitrile; 60 min, -20 °C		
31-279-CAS-14439984	Steps: 1 Yield: 100%	Catalytic asymmetric epoxidation of unfunctionalized olefins using silica, LDH and resin-supported sulfonate-Mn(salen) complex By: Choudary, B. M.; et al Advanced Synthesis & Catalysis (2006), 348(4 + 5), 493-498.
1.1 Reagents: 4-Methylmorpholine, <i>m</i> -Chloroperbenzoic acid Catalysts: Manganate(2-), [[3,3'-[1,2-cyclohexanediylbis[[nitrilo-κM)methylidyne]]bis[5-(1,1-dimethylethyl)-4-(hydroxy-κO)benzenesulfonato]](4-)](2,4-pentanedionato-κO ² ,κO ⁴)-, sodium (1:2) Solvents: Acetonitrile; < 5 min, -20 °C		
31-279-CAS-10174242	Steps: 1 Yield: 100%	Catalytic asymmetric epoxidation of unfunctionalized olefins using silica, LDH and resin-supported sulfonate-Mn(salen) complex By: Choudary, B. M.; et al Advanced Synthesis & Catalysis (2006), 348(4 + 5), 493-498.
1.1 Reagents: 4-Methylmorpholine, <i>m</i> -Chloroperbenzoic acid Catalysts: Manganate(2-), [[3,3'-[1,2-cyclohexanediylbis[[nitrilo-κM)methylidyne]]bis[5-(1,1-dimethylethyl)-4-(hydroxy-κO)benzenesulfonato]](4-)](2,4-pentanedionato-κO ² ,κO ⁴)-, sodium (1:2) Solvents: Acetonitrile; 120 min, -20 °C		
31-279-CAS-9716207	Steps: 1 Yield: 100%	Is hole transfer involved in metalloporphyrin-catalyzed epoxidation? By: Kim, Taisun; et al Journal of the American Chemical Society (1993), 115(17), 7653-64.
1.1 Reagents: Nitrogen, Phenylseleninyl benzeneseleninate Catalysts: Antimonate(1-), hexachloro-, (OC-6-11)-, salt with 4-bromo- <i>N,N</i> -bis(4-bromophenyl)benzamine (1:1) Solvents: Dichloromethane		

Scheme 40 (1 Reaction)

Steps: 1 Yield: 100%



Suppliers (63)

Suppliers (65)

31-279-CAS-18402772

Steps: 1 Yield: 100%

Synthesis of new polyfluorinated oxaziridines

1.1 Solvents: Chloroform-*d*₃; -50 °C

By: Petrov, Viacheslav; et al


1.2 Reagents: 3-Fluoro-2-(1,1,2,2,2-pentafluoroethyl)-3-(trifluoromethyl)oxaziridine; 10 min, -50 °C → 25 °C; 1 - 2 h, 25 °C

Journal of Fluorine Chemistry (2018), 207, 1-6.

Experimental Protocols

Scheme 41 (5 Reactions)

Steps: 1 Yield: 100%



Suppliers (115)

Suppliers (66)

31-279-CAS-16778979

Steps: 1 Yield: 100%

Study on the synthesis of chiral salen Mn(III) immobilized onto phenoxyl linker group modified ZPS-PVPA by axial coordination and the asymmetric epoxidation of unfunctionalized olefins

1.1 Reagents: *m*-Chloroperbenzoic acid
Catalysts: N-methylmorpholine N-oxide, *S,S*-Jacobsen catalyst (reaction products with hydroxy-containing polystyrene)
Solvents: Dichloromethane; 1 h, 0 °C

By: Zou, Xiaochuan; et al

1.2 Reagents: Sodium hydroxide
Solvents: Water; > pH 7

Huaxue Xuebao (2011), 69(4), 431-437.

31-279-CAS-3511200

Steps: 1 Yield: 100%

Catalytic asymmetric epoxidation of unfunctionalized olefins using silica, LDH and resin-supported sulfonate-Mn(salen) complex

1.1 Reagents: 4-Methylmorpholine, *m*-Chloroperbenzoic acid
Catalysts: Manganate(2-), [[3,3'-(1,2-cyclohexanediyl)bis[(nitrilo-κM)methylidyne]]bis[5-(1,1-dimethylethyl)-4-(hydroxy-κO)benzenesulfonato]](4-)](2,4-pentanedionato-κO²,κO⁴)-, sodium (1:2)
Solvents: Acetonitrile; < 5 min, -20 °C

By: Choudary, B. M.; et al

Advanced Synthesis & Catalysis (2006), 348(4 + 5), 493-498.

31-279-CAS-1365423

Steps: 1 Yield: 100%

Catalytic asymmetric epoxidation of unfunctionalized olefins using silica, LDH and resin-supported sulfonate-Mn(salen) complex

1.1 Reagents: 4-Methylmorpholine, *m*-Chloroperbenzoic acid
Catalysts: Manganate(2-), [[3,3'-(1,2-cyclohexanediyl)bis[(nitrilo-κM)methylidyne]]bis[5-(1,1-dimethylethyl)-4-(hydroxy-κO)benzenesulfonato]](4-)](2,4-pentanedionato-κO²,κO⁴)-, sodium (1:2)
Solvents: Acetonitrile; 10 min, -20 °C

By: Choudary, B. M.; et al

Advanced Synthesis & Catalysis (2006), 348(4 + 5), 493-498.

31-279-CAS-5637144 Steps: 1 Yield: 100% 1.1 Reagents: 4-Methylmorpholine, <i>m</i>-Chloroperbenzoic acid Catalysts: Manganate(2-), [[3,3'-(1,2-cyclohexanediyl)bis[(nitrilo-κM)methylidyne]]bis[5-(1,1-dimethylethyl)-4-(hydroxy-κO)benzenesulfonato]](4-)](2,4-pentanedionato-κO²,κO⁴)-, sodium (1:2) Solvents: Acetonitrile; < 5 min, -20 °C	Catalytic asymmetric epoxidation of unfunctionalized olefins using silica, LDH and resin-supported sulfonate-Mn(salen) complex By: Choudary, B. M.; et al Advanced Synthesis & Catalysis (2006), 348(4 + 5), 493-498.
31-279-CAS-14763020 Steps: 1 Yield: 100% 1.1 Reagents: 4-Methylmorpholine, <i>m</i>-Chloroperbenzoic acid Catalysts: Manganate(2-), [[3,3'-(1,2-cyclohexanediyl)bis[(nitrilo-κM)methylidyne]]bis[5-(1,1-dimethylethyl)-4-(hydroxy-κO)benzenesulfonato]](4-)](2,4-pentanedionato-κO²,κO⁴)-, sodium (1:2) Solvents: Acetonitrile; < 10 min, -20 °C	Catalytic asymmetric epoxidation of unfunctionalized olefins using silica, LDH and resin-supported sulfonate-Mn(salen) complex By: Choudary, B. M.; et al Advanced Synthesis & Catalysis (2006), 348(4 + 5), 493-498.