

a
Number of isolates
EN
39.9%
EP
60.1%
Pantoea
Enterobacter
Bacillus
Burkholderia
Klebsiella
Stenotrophomonas
Staphylococcus
Curtobacterium
Other
Genus
b
Spearman r = 0.16, P = 0.534
Isolation (%)
Amplicon (%)
c
Spearman r = 0.86, P < 0.001
Landrace relative abundance (%)
Inbred line relative abundance (%)
d
Spearman r = 0.89, P < 0.001
Epiphyte relative abundance (%)
Endophyte relative abundance (%)
e
Species
D011P03
D119N01
D018P05
D113N10
D014P08
D053P02
D112P05
f
P. dispersa
Orientation
Reverse
Forward
D011P03 (Mb)
D119N01 (Mb)
g
P. stewartii
D018P05 (Mb)
D113N10 (Mb)
h
B. gladioli
D112P05 (Mb)
D014P08 (Mb)
i
P. dispersa vs D011P03
P. stewartii vs D113N10
B. gladioli vs D112P05
Mean alignment identity (%)
j
P. dispersa
P. stewartii
B. gladioli
Count
Gaps >10 kb
Inversions
k
SNP density (SNPs/kb)
P. dispersa
P. stewartii
B. gladioli Chr1
B. gladioli Chr2
l
Ti/Tv ratio
Ti/Tv = 2
Ti/Tv = 1