

Data Dictionary for Clinical Metadata, v1 (June 2023 release)

individualID	identifying string linked to a particular individual (example: R1234567)
---------------------	--

dataContributionGroup	The research group who contributed the sample	
		Columbia
		Emory
		Mayo
		MSSM
		Rush

cohort	The initial study group population to which the individual belonged
	AA
	Banner
	Biggs Institute Brain Bank
	CLINCOR
	Emory
	LATC
	MAP
	MARS
	Mayo Clinic
	Mt Sinai Brain Bank
	NY Brain Bank
	UPenn

	ROS
	UFL

sex	Self-reported sex
	Female
	Male

race	Self-reported race
	American Indian or Alaska Native
	Asian
	Black or African American
	White
	other
	Missing or unknown

isHispanic	Self-reported indicator for Hispanic or Latino ethnicity
	True
	False
	Missing or unknown

ageDeath	Age at Death (in years)
-----------------	-------------------------

pmi	Post-mortem interval (in hours)
------------	---------------------------------

apoeGenotype	Apolipoprotein E genotype	
	Coded value	description
	22	ε2ε2
	23	ε2ε3
	24	ε2ε4
	33	ε3ε3
	34	ε3ε4
	44	ε4ε4
	missing or unknown	

amyThal	Thal Phase (a measure of amyloid deposition)	
	None	
	Phase 1	
	Phase 2	
	Phase 3	
	Phase 4	
	Phase 5	
	missing or unknown	

amyA	Indicator of Thal Phase, grouped	
	None	
	Thal Phase 1 or 2	
	Thal Phase 3	
	Thal Phase 4 or 5	

	Missing or unknown
--	--------------------

amyCerad	CERAD score, a semi-quantitative measure of neuritic plaque
	None/No AD/C0
	Sparse/Possible/C1
	Moderate/Probable/C2
	Frequent/Definite/C3
	Missing or unknown

amyAny	Indicator for presence of any amyloid by CERAD score	
	Coded value	description
	0	amyCerad = None/No AD/C0
	1	amyCerad = Sparse/Possible/C1 or Moderate/Probable/C2 or Frequent/Definite/C3
	Missing or unknown	

Braak	Braak stage, a semi-quantitative measure of neurofibrillary tangle pathology
	None
	Stage I
	Stage II
	Stage III
	Stage IV
	Stage V

	Stave VI
	Missing or unknown

bScore	Indicator of Braak Stage, grouped
	None
	Braak Stage I-II
	Braak Stage III-IV
	Braak Stage V-VI
	Missing or unknown

mayoDx	For samples from the Mayo clinic cohort, this variable is a neuropathology-based diagnosis performed by Mayo Clinic neurologists per NINCDS-ADRD criteria, under the following thresholds:	
	Coded value	Definition
	AD	Braak $\geq$ Stage IV and amyThal $\geq$ Phase 2, in absence of non-AD tauopathy.
	Control	Braak $\leq$ Stage III and amyThal $<$ Phase 2
	other	All other combinations and non-AD tauopathies
	not applicable	Participant is not from a Mayo Clinic cohort
For Mayo samples from the UFL cohort, neuropathologist-assigned diagnoses follow NIA guidelines for neuropathological diagnosis of AD ((Hyman et al., PMID:22265587 and Montine et al., PMID: 22101365). AD case status was assigned to patients with any degree of AD neuropathological diagnosis.		

reag	Modified NIA Reagan, following rubric established by Rush.	
	Coded value	definition
	High Likelihood	amyCerad = Definite/frequent/C3 and Braak $\geq$ Stage V
	Intermediate Likelihood	amyCerad = Definite/frequent/C3 and Braak = Stage I, II, III, or IV OR: amyCerad = Probable/moderate/C2 and Braak $\geq$ Stage III
	Low Likelihood	amyCerad = Probable/moderate/C2 and Braak $\leq$ Stage II OR: amyCerad = Possible/sparse or infrequent/C1 and Braak = any Stage OR: amyCerad = No AD and Braak $\geq$ Stage I
	No AD	amyCerad = No AD/None/C0 and Braak = None

ADoutcome	Derived harmonized outcomes based on CERAD neuritic plaque score and Braak stage, or NINCDS-ADRD criteria for Mayo. <i>The following criteria seek to harmonize outcomes between cohorts as closely as possible, with Mayo's neuropath-based diagnoses (as determined by Mayo specialists) set as the standard to match.</i>	
	Rush, MSSM, Columbia, and Emory derived outcome criteria:	
	Coded value	Definition
	AD	amyCerad = Definite/frequent/C3 or Probable/moderate/C2 and Braak $\geq$ Stage IV
	Control	amyCerad = Possible/sparse or

		infrequent/C1 or None/No AD/C0 and Braak $\leq$ Stage III
	Other	All other combinations
	Some patients from the UFL cohort (via Mayo) were not consistent with other MayoDx criteria (i.e. 8 individuals had Braak < 4 and a mayoDx==AD). These individuals were assigned 'Other' criteria in the ADoutcome variable to harmonize more closely with MayoDx criteria.	