

1 Section 1.7

Question 6

Question 7

Question 8

Question 9

2 Section 2.2

Question 4

Let $f(x, y, z) = z^2$. Prove that 0 is not a regular value off and yet that $f^{-1}(0)$ is a regular surface.

Solution. Note that

$$df = (f_x, f_y, f_z) = (0, 0, 2z),$$

which is not surjective only when $z = 0$. Hence, $(0, 0, 0)$ is a critical point and thus $f(0, 0, 0) = 0$ is not a regular value.

However,

$$\begin{aligned} f^{-1}(0) &= \{(x, y, z) \in \mathbb{R}^3 | z^2 = 0\} \\ &= \{(x, y, 0) | x, y \in \mathbb{R}\} \\ &= \mathbb{R}^2 \times 0. \end{aligned}$$

Hence, $f^{-1}(0)$ is homeomorphic to \mathbb{R}^2 and therefore is regular.

Question 5

Let $P = \{(x, y, z) \in \mathbb{R}^3 | x = y\}$ (a plane) and let $x : U \subset \mathbb{R}^2 \rightarrow \mathbb{R}^3$ be given by

$$x(u, v) = (u + v, u + v, uv),$$

where $U = \{(u, v) \in \mathbb{R}^2 | u > v\}$. Clearly, $x(U) \subset P$. Is x a parametrization of P ?

Solution. Yes, x is a parametrization. Clearly, x is differentiable in U with

$$dx(u, v) = \begin{pmatrix} 1 & 1 \\ 1 & 1 \\ u & v \end{pmatrix}.$$

Note that for $(u, v) \in U$, we have $u > v$. Then

$$\begin{vmatrix} 1 & 1 \\ u & v \end{vmatrix} = v - u \neq 0.$$

This implies $dx(u, v)$ is injective for all $(u, v) \in U$. Now let (a, a, b) be any point in $x(U)$. Then

$$\begin{aligned} u + v &= a, uv = b \\ \Rightarrow u(a - u) &= b \\ \Rightarrow \left(u - \frac{a}{2}\right)^2 &= \frac{a^2}{4} - b. \end{aligned}$$

Notice that here one must have $\frac{a^2}{4} - b \geq 0$ as the equations $\begin{cases} u + v = a \\ uv = b \end{cases}$ should have real solutions for $(a, a, b) \in x(U)$. Then given $u > v$, we have

$$\begin{aligned} u &= \frac{a}{2} + \sqrt{\frac{a^2}{4} - b} \\ v &= \frac{a}{2} - \sqrt{\frac{a^2}{4} - b}. \end{aligned}$$

These are the unique (u, v) solving $x(u, v) = (a, b)$, which shows x is injective. Hence, by Prop.4 x^{-1} must be continuous and we can conclude that x is indeed a parametrization.

Question 6

Give another proof of Prop. 1 by applying Prop. 2 to $h(x, y, z) = f(x, y) - z$.

Solution. Since f is differentiable in U , for any point in $U \times \mathbb{R}$, we have

$$dh = (f_x, f_y, -1),$$

which is always surjective regardless of the value of f_x, f_y . Hence, any $z_0 \in f(U)$ with $f(x_0, y_0) = z_0$, we have

$$h(x_0, y_0, z_0) = f(x_0, y_0) - z_0 = 0,$$

being a regular value. This implies that

$$\begin{aligned} h^{-1}(0) &= \{(x, y, z) \in U \times \mathbb{R} \mid h(x, y, z) = 0\} \\ &= \{(x, y, z) \in U \times \mathbb{R} \mid f(x, y) = z\} \\ &= \{(x, y, f(x, y)) \mid (x, y) \in U\} \end{aligned}$$

is a regular surface.