

Based on: T.Davidson, F.Kloosterman, M.Wilson "Hippocampal replay of extended experience", in Neuron, vol. 63, pp. 497-507, 2009

$$\begin{aligned}
Pr(x|spikes) &= \frac{Pr(spikes|x)Pr(x)}{Pr(spikes)} \\
Pr(spikes|x) &= \prod_{i=1}^N Pr(spikes_i|x) = \\
&= \prod_{i=1}^N \frac{\Delta t \tau_i(x)^{n_i}}{n_i!} \exp(-\Delta t \tau_i(x)) \\
\log(Pr(spikes|x)) &= \sum_{i=1}^N n_i \log\left(\frac{\Delta t \tau_i(x)}{n_i!}\right) - \Delta t \sum_{i=1}^N \tau_i(x)
\end{aligned}$$

$$\tau_i(x) = \begin{cases} \cos\left(\frac{2\pi}{2\varphi_{PF}}(x - \varphi_{mPF_i})\right) \tau_{avg}, & \text{if } x \in PF \\ 0, & \text{otherwise} \end{cases}$$

where:

N: $\#\{\text{neurons}\} = 4000$, Δt : length of the time bin, n_i : $\#\{\text{spikes in the bin (from the i-th cell)}\}$, φ_{PF} : length of a place field = $0.628[\text{rad}]$, φ_{mPF_i} : middle of the i-th cell's place field, $\tau_{avg} = 20[\text{Hz}]$