```
Tristan Erney
November 29<sup>th</sup>, 2021
Intro to Cryptology
Hands On Exercise 13
1)
        p = 13
       root = 2
        primititve roots = 2, 6, 7, 11
primitive_roots.py
#!/usr/bin/env python
# assume all p's are prime
def primitive_roots(p):
  roots = []
  phi = p - 1
  for i in range(2, p):
     k = 0
     for j in range(1, p):
        k = i ** j \% p
          if k == 1 and j != phi:
               break
         elif k == 1 and j == phi:
               roots.append(i)
  return roots
roots = primitive_roots(13)
print("primitive roots = {}".format(roots))
print("dlog 2, 13 (3) = {}".format(2 ** 3 % 13))
print("dlog 2, 13 (11) = {}".format(2 ** 11 % 13))
       dlog_{2,13}(3) = 2^3 \mod 13 = 8
        dlog_{2,13}(11) = 2^{11} \mod 13 = 7
2)
       6^5 \pmod{11} = 10 \pmod{11}
3)
       q = 71
        a = 7
       X_A = 5
        X_B = 12
```

a)
$$Y_A = (a)^{XA} = 7^5 \mod 71 = 51 \pmod{71}$$

b)
$$Y_B = (a)^{XB} = 7^{12} \mod 71 = 4 \pmod{71}$$

c)
$$K = (Y_A)^{XB} = 51^{12} \mod 71 = 30$$

$$K = (Y_B)^{XA} = 4^5 \mod 71 = 30$$

4)

$$q = 11$$

 $a = 2$
 $Y_A = 9$
 $Y_B = 3$

a)
$$phi(11) = 10$$

$$2^{1} = 2$$
 $2^{2} = 4$
 $2^{3} = 8$
 $2^{4} = 5$
 $2^{5} = 10$

$$2^{6} = 10$$

 $2^{6} = 9$

$$2^7 = 7$$

 $2^8 = 3$

$$2^9 = 6$$

$$2^{10} = 1$$

b)
$$Y_A = (a)^{XA} = 2^{XA} \mod 11 = 9 \pmod 11$$

$$X_A = 6$$

c)
$$K = (Y_B)^{XA} = 3^6 \mod 11 = 3 \pmod{11}$$

5)

$$a = 3$$

$$Y_A = 27$$
 $\rightarrow 3^{XA} = 27$ $\rightarrow X_A = 3$
 $Y_B = 243$ $\rightarrow 3^{XB} = 243$ $\rightarrow X_B = 5$

$$K = (Y_A)^{XB} = 27^5 = 14348907$$

 $K = (Y_B)^{XA} = 243^3 = 14348907$

6)

$$q = 71$$

 $a = 7$

a)
$$Y_B = 3$$

$$k = 2$$

$$M = 30$$

$$K = (Y_B)^k \mod q = 3^2 \mod 71 = 9 \pmod 71$$

$$C_1 = a^k = 7^2 \mod 71 = 49 \pmod 71$$

$$C_2 = KM = 9(30) \mod 71 = 57 \pmod 71$$

$$C = (49, 57)$$

b)
$$M = 30$$

$$a = 7$$

$$C = (59, C_2)$$

$$C_1 = a^k = 7^k \mod 71 = 59 \pmod 71$$

$$k = 3$$

$$K = (Y_B)^k = 3^3 \mod 71 = 27 \pmod 71$$

$$C_2 = KM = 27(30) \mod 71 = 29 \pmod 71$$