

# VHEGEN: A vibronic Hamiltonian expansion generator for trigonal and tetragonal polyatomic systems

James Brown      Robert A. Lang      Riley J. Hickman      Tao Zeng

Thank you for using **VHEGEN**, the **V**-ibronic **H**-amiltonian **E**-xpansion **GEN**-erator for trigonal and tetragonal polyatomic systems. This is a **VHEGEN** output file compiled by **pdflatex**. If the **VHEGEN** package was used in research resulting in a publication, please reference the article in *Computer Physics Communications* which describes the program ([doi here]). Additional information regarding the matrix element expansion process, including the independent matrix element eigenvalues, their root formulas and constraints, and their transformation to the real basis (if applicable), can be found in the **log** output file. For questions, bugs, or comments, please contact [jbrown88@yorku.ca](mailto:jbrown88@yorku.ca).

## Contents

<b>1</b>	<b>Vibronic interaction</b>	<b>2</b>
<b>2</b>	<b>Vibronic Hamiltonian operator in the complex <math>E</math> basis</b>	<b>2</b>
<b>3</b>	<b>Matrix element expansions in the complex <math>E</math> basis</b>	<b>2</b>
3.1	Order: 0 . . . . .	2
3.2	Order: 1 . . . . .	2
3.3	Order: 2 . . . . .	3
3.4	Order: 3 . . . . .	4
<b>4</b>	<b>Vibronic Hamiltonian operator in the real <math>E</math> basis</b>	<b>5</b>
<b>5</b>	<b>Matrix element expansions in the real <math>E</math> basis</b>	<b>5</b>
5.1	Order: 0 . . . . .	5
5.2	Order: 1 . . . . .	5
5.3	Order: 2 . . . . .	6
5.4	Order: 3 . . . . .	7

# 1 Vibronic interaction

$$E_1'' \otimes (e_1' + e_1'') \text{ in } C_{5h}$$

$$\begin{aligned} \rho_1, \phi_1, x_1, y_1 &\rightarrow e_1' \\ \rho_2, \phi_2, x_2, y_2 &\rightarrow e_1'' \end{aligned}$$

## 2 Vibronic Hamiltonian operator in the complex $E$ basis

$$\hat{H} = (|+\rangle \quad |-\rangle) \begin{pmatrix} H_{++} & H_{+-} \\ H_{-+} & H_{--} \end{pmatrix} \begin{pmatrix} \langle +| \\ \langle -| \end{pmatrix}$$

## 3 Matrix element expansions in the complex $E$ basis

### 3.1 Order: 0

Number of fitting parameters:  $H_{++}$ : 2,  $H_{+-}$ : 0.

**Polar e-coordinates:**

$$H_{++}^{(0)} = ia_{0,0,0,0}^i + a_{0,0,0,0}^r$$

$$H_{--}^{(0)} = ia_{0,0,0,0}^i + a_{0,0,0,0}^r$$

$$H_{+-}^{(0)} = 0$$

$$H_{-+}^{(0)} = 0$$

**Cartesian e-coordinates:**

$$H_{++}^{(0)} = ia_{0,0,0,0}^i + a_{0,0,0,0}^r$$

$$H_{--}^{(0)} = ia_{0,0,0,0}^i + a_{0,0,0,0}^r$$

$$H_{+-}^{(0)} = 0$$

$$H_{-+}^{(0)} = 0$$

### 3.2 Order: 1

Number of fitting parameters:  $H_{++}$ : 0,  $H_{+-}$ : 0.

**Polar e-coordinates:**

$$H_{++}^{(1)}=0$$

$$H_{--}^{(1)}=0$$

$$H_{+-}^{(1)}=0$$

$$H_{-+}^{(1)}=0$$

**Cartesian e-coordinates:**

$$H_{++}^{(1)}=0$$

$$H_{--}^{(1)}=0$$

$$H_{+-}^{(1)}=0$$

$$H_{-+}^{(1)}=0$$

### 3.3 Order: 2

Number of fitting parameters:  $H_{++}$ : 4,  $H_{+-}$ : 4.

**Polar e-coordinates:**

$$H_{++}^{(2)}=ia_{0,0,0,1}^i\rho_2^2+ia_{0,1,0,0}^i\rho_1^2+a_{0,0,0,1}^r\rho_2^2+a_{0,1,0,0}^r\rho_1^2$$

$$H_{--}^{(2)}=ia_{0,0,0,1}^i\rho_2^2+ia_{0,1,0,0}^i\rho_1^2+a_{0,0,0,1}^r\rho_2^2+a_{0,1,0,0}^r\rho_1^2$$

$$H_{+-}^{(2)}=ib_{-2,0,0,0}^i\rho_1^2\exp(-2i\phi_1)+ib_{0,0,-2,0}^i\rho_2^2\exp(-2i\phi_2)+b_{-2,0,0,0}^r\rho_1^2\exp(-2i\phi_1)+b_{0,0,-2,0}^r\rho_2^2\exp(-2i\phi_2)$$

$$H_{-+}^{(2)}=-ib_{-2,0,0,0}^i\rho_1^2\exp(2i\phi_1)-ib_{0,0,-2,0}^i\rho_2^2\exp(2i\phi_2)+b_{-2,0,0,0}^r\rho_1^2\exp(2i\phi_1)+b_{0,0,-2,0}^r\rho_2^2\exp(2i\phi_2)$$

### Cartesian e-coordinates:

$$H_{++}^{(2)} = ia_{0,0,0,1}^i(x_2^2 + y_2^2) + ia_{0,1,0,0}^i(x_1^2 + y_1^2) + a_{0,0,0,1}^r(x_2^2 + y_2^2) + a_{0,1,0,0}^r(x_1^2 + y_1^2)$$

$$H_{--}^{(2)} = ia_{0,0,0,1}^i(x_2^2 + y_2^2) + ia_{0,1,0,0}^i(x_1^2 + y_1^2) + a_{0,0,0,1}^r(x_2^2 + y_2^2) + a_{0,1,0,0}^r(x_1^2 + y_1^2)$$

$$H_{+-}^{(2)} = 2b_{-2,0,0,0}^i x_1 y_1 + ib_{-2,0,0,0}^i (x_1 - y_1)(x_1 + y_1) + 2b_{0,0,-2,0}^i x_2 y_2 + ib_{0,0,-2,0}^i (x_2 - y_2)(x_2 + y_2) - 2ib_{-2,0,0,0}^r x_1 y_1 + b_{-2,0,0,0}^r (x_1 - y_1)(x_1 + y_1) - 2ib_{0,0,-2,0}^r x_2 y_2 + b_{0,0,-2,0}^r (x_2 - y_2)(x_2 + y_2)$$

$$H_{-+}^{(2)} = 2b_{-2,0,0,0}^i x_1 y_1 - ib_{-2,0,0,0}^i (x_1 - y_1)(x_1 + y_1) + 2b_{0,0,-2,0}^i x_2 y_2 - ib_{0,0,-2,0}^i (x_2 - y_2)(x_2 + y_2) + 2ib_{-2,0,0,0}^r x_1 y_1 + b_{-2,0,0,0}^r (x_1 - y_1)(x_1 + y_1) + 2ib_{0,0,-2,0}^r x_2 y_2 + b_{0,0,-2,0}^r (x_2 - y_2)(x_2 + y_2)$$

### 3.4 Order: 3

Number of fitting parameters:  $H_{++}$ : 0,  $H_{+-}$ : 4.

### Polar e-coordinates:

$$H_{++}^{(3)} = 0$$

$$H_{--}^{(3)} = 0$$

$$H_{+-}^{(3)} = ib_{1,0,2,0}^i \rho_1 \rho_2^2 \exp(i(\phi_1 + 2\phi_2)) + ib_{3,0,0,0}^i \rho_1^3 \exp(3i\phi_1) + b_{1,0,2,0}^r \rho_1 \rho_2^2 \exp(i(\phi_1 + 2\phi_2)) + b_{3,0,0,0}^r \rho_1^3 \exp(3i\phi_1)$$

$$H_{-+}^{(3)} = -ib_{1,0,2,0}^i \rho_1 \rho_2^2 \exp(-i(\phi_1 + 2\phi_2)) - ib_{3,0,0,0}^i \rho_1^3 \exp(-3i\phi_1) + b_{1,0,2,0}^r \rho_1 \rho_2^2 \exp(-i(\phi_1 + 2\phi_2)) + b_{3,0,0,0}^r \rho_1^3 \exp(-3i\phi_1)$$

### Cartesian e-coordinates:

$$H_{++}^{(3)} = 0$$

$$H_{--}^{(3)} = 0$$

$$H_{+-}^{(3)} = ib_{1,0,2,0}^i (x_1(x_2^2 - y_2^2) - 2x_2 y_1 y_2) - b_{1,0,2,0}^i (2x_1 x_2 y_2 + y_1(x_2^2 - y_2^2)) + ib_{3,0,0,0}^i x_1(x_1^2 - 3y_1^2) - b_{3,0,0,0}^i y_1(3x_1^2 - y_1^2) + b_{1,0,2,0}^r (x_1(x_2^2 - y_2^2) - 2x_2 y_1 y_2) + ib_{1,0,2,0}^r (2x_1 x_2 y_2 + y_1(x_2^2 - y_2^2)) + b_{3,0,0,0}^r x_1(x_1^2 - 3y_1^2) + ib_{3,0,0,0}^r y_1(3x_1^2 - y_1^2)$$

$$H_{-+}^{(3)} = -ib_{1,0,2,0}^i (x_1(x_2^2 - y_2^2) - 2x_2 y_1 y_2) - b_{1,0,2,0}^i (2x_1 x_2 y_2 + y_1(x_2^2 - y_2^2)) - ib_{3,0,0,0}^i x_1(x_1^2 - 3y_1^2) - b_{3,0,0,0}^i y_1(3x_1^2 - y_1^2) + b_{1,0,2,0}^r (x_1(x_2^2 - y_2^2) - 2x_2 y_1 y_2) - ib_{1,0,2,0}^r (2x_1 x_2 y_2 + y_1(x_2^2 - y_2^2)) + b_{3,0,0,0}^r x_1(x_1^2 - 3y_1^2) - ib_{3,0,0,0}^r y_1(3x_1^2 - y_1^2)$$

## 4 Vibronic Hamiltonian operator in the real $E$ basis

$$\hat{H} = (|X\rangle \quad |Y\rangle) \begin{pmatrix} H_{XX} & H_{XY} \\ H_{YX} & H_{YY} \end{pmatrix} \begin{pmatrix} \langle X| \\ \langle Y| \end{pmatrix}$$

## 5 Matrix element expansions in the real $E$ basis

### 5.1 Order: 0

Number of fitting parameters:  $H_{XX}$ : 1 (all from  $H_{++}$ ),  $H_{XY}$ : 0,  $H_{YY}$ : 1 (all from  $H_{++}$ ).

**Polar e-coordinates:**

$$H_{XX}^{(0)} = ia_{0,0,0,0}^i + a_{0,0,0,0}^r$$

$$H_{XY}^{(0)} = 0$$

$$H_{YX}^{(0)} = 0$$

$$H_{YY}^{(0)} = ia_{0,0,0,0}^i + a_{0,0,0,0}^r$$

**Cartesian e-coordinates:**

$$H_{XX}^{(0)} = ia_{0,0,0,0}^i + a_{0,0,0,0}^r$$

$$H_{XY}^{(0)} = 0$$

$$H_{YX}^{(0)} = 0$$

$$H_{YY}^{(0)} = ia_{0,0,0,0}^i + a_{0,0,0,0}^r$$

### 5.2 Order: 1

Number of fitting parameters:  $H_{XX}$ : 0,  $H_{XY}$ : 0,  $H_{YY}$ : 0.

**Polar e-coordinates:**

$$H_{XX}^{(1)} = 0$$

$$H_{XY}^{(1)} = 0$$

$$H_{YX}^{(1)} = 0$$

$$H_{YY}^{(1)} = 0$$

**Cartesian e-coordinates:**

$$H_{XX}^{(1)}=0$$

$$H_{XY}^{(1)}=0$$

$$H_{YX}^{(1)}=0$$

$$H_{YY}^{(1)}=0$$

### 5.3 Order: 2

Number of fitting parameters:  $H_{XX}$ : 6 (2 from  $H_{++}$ , 4 from  $H_{+-}$ ),  $H_{XY}$ : 4 (all from  $H_{+-}$ ),  $H_{YY}$ : 6 (2 from  $H_{++}$ , 4 from  $H_{+-}$ ).

**Polar e-coordinates:**

$$H_{XX}^{(2)} = a_{0,0,0,1}^r \rho_2^2 + a_{0,1,0,0}^r \rho_1^2 + b_{-2,0,0,0}^i \rho_1^2 \sin(2\phi_1) + b_{0,0,-2,0}^i \rho_2^2 \sin(2\phi_2) + b_{-2,0,0,0}^r \rho_1^2 \cos(2\phi_1) + b_{0,0,-2,0}^r \rho_2^2 \cos(2\phi_2) + i(a_{0,0,0,1}^i \rho_2^2 + a_{0,1,0,0}^i \rho_1^2)$$

$$H_{XY}^{(2)} = -b_{-2,0,0,0}^i \rho_1^2 \cos(2\phi_1) - b_{0,0,-2,0}^i \rho_2^2 \cos(2\phi_2) + b_{-2,0,0,0}^r \rho_1^2 \sin(2\phi_1) + b_{0,0,-2,0}^r \rho_2^2 \sin(2\phi_2)$$

$$H_{YX}^{(2)} = -b_{-2,0,0,0}^i \rho_1^2 \cos(2\phi_1) - b_{0,0,-2,0}^i \rho_2^2 \cos(2\phi_2) + b_{-2,0,0,0}^r \rho_1^2 \sin(2\phi_1) + b_{0,0,-2,0}^r \rho_2^2 \sin(2\phi_2)$$

$$H_{YY}^{(2)} = a_{0,0,0,1}^r \rho_2^2 + a_{0,1,0,0}^r \rho_1^2 - b_{-2,0,0,0}^i \rho_1^2 \sin(2\phi_1) - b_{0,0,-2,0}^i \rho_2^2 \sin(2\phi_2) - b_{-2,0,0,0}^r \rho_1^2 \cos(2\phi_1) - b_{0,0,-2,0}^r \rho_2^2 \cos(2\phi_2) + i(a_{0,0,0,1}^i \rho_2^2 + a_{0,1,0,0}^i \rho_1^2)$$

**Cartesian e-coordinates:**

$$H_{XX}^{(2)} = a_{0,0,0,1}^r (x_2^2 + y_2^2) + a_{0,1,0,0}^r (x_1^2 + y_1^2) + 2b_{-2,0,0,0}^i x_1 y_1 + 2b_{0,0,-2,0}^i x_2 y_2 + b_{-2,0,0,0}^r (x_1 - y_1)(x_1 + y_1) + b_{0,0,-2,0}^r (x_2 - y_2)(x_2 + y_2) + i(a_{0,0,0,1}^i (x_2^2 + y_2^2) + a_{0,1,0,0}^i (x_1^2 + y_1^2))$$

$$H_{XY}^{(2)} = -b_{-2,0,0,0}^i (x_1 - y_1)(x_1 + y_1) - b_{0,0,-2,0}^i (x_2 - y_2)(x_2 + y_2) + 2b_{-2,0,0,0}^r x_1 y_1 + 2b_{0,0,-2,0}^r x_2 y_2$$

$$H_{YX}^{(2)} = -b_{-2,0,0,0}^i (x_1 - y_1)(x_1 + y_1) - b_{0,0,-2,0}^i (x_2 - y_2)(x_2 + y_2) + 2b_{-2,0,0,0}^r x_1 y_1 + 2b_{0,0,-2,0}^r x_2 y_2$$

$$H_{YY}^{(2)} = a_{0,0,0,1}^r (x_2^2 + y_2^2) + a_{0,1,0,0}^r (x_1^2 + y_1^2) - 2b_{-2,0,0,0}^i x_1 y_1 - 2b_{0,0,-2,0}^i x_2 y_2 - b_{-2,0,0,0}^r (x_1 - y_1)(x_1 + y_1) - b_{0,0,-2,0}^r (x_2 - y_2)(x_2 + y_2) + i(a_{0,0,0,1}^i (x_2^2 + y_2^2) + a_{0,1,0,0}^i (x_1^2 + y_1^2))$$

## 5.4 Order: 3

Number of fitting parameters:  $H_{XX}$ : 4 (all from  $H_{+-}$ ),  $H_{XY}$ : 4 (all from  $H_{+-}$ ),  $H_{YY}$ : 4 (all from  $H_{+-}$ ).

### Polar e-coordinates:

$$H_{XX}^{(3)} = -b_{1,0,2,0}^i \rho_1 \rho_2^2 \sin(\phi_1 + 2\phi_2) - b_{3,0,0,0}^i \rho_1^3 \sin(3\phi_1) + b_{1,0,2,0}^r \rho_1 \rho_2^2 \cos(\phi_1 + 2\phi_2) + b_{3,0,0,0}^r \rho_1^3 \cos(3\phi_1)$$

$$H_{XY}^{(3)} = -b_{1,0,2,0}^i \rho_1 \rho_2^2 \cos(\phi_1 + 2\phi_2) - b_{3,0,0,0}^i \rho_1^3 \cos(3\phi_1) - b_{1,0,2,0}^r \rho_1 \rho_2^2 \sin(\phi_1 + 2\phi_2) - b_{3,0,0,0}^r \rho_1^3 \sin(3\phi_1)$$

$$H_{YX}^{(3)} = -b_{1,0,2,0}^i \rho_1 \rho_2^2 \cos(\phi_1 + 2\phi_2) - b_{3,0,0,0}^i \rho_1^3 \cos(3\phi_1) - b_{1,0,2,0}^r \rho_1 \rho_2^2 \sin(\phi_1 + 2\phi_2) - b_{3,0,0,0}^r \rho_1^3 \sin(3\phi_1)$$

$$H_{YY}^{(3)} = b_{1,0,2,0}^i \rho_1 \rho_2^2 \sin(\phi_1 + 2\phi_2) + b_{3,0,0,0}^i \rho_1^3 \sin(3\phi_1) - b_{1,0,2,0}^r \rho_1 \rho_2^2 \cos(\phi_1 + 2\phi_2) - b_{3,0,0,0}^r \rho_1^3 \cos(3\phi_1)$$

### Cartesian e-coordinates:

$$H_{XX}^{(3)} = -b_{1,0,2,0}^i (2x_1 x_2 y_2 + y_1 (x_2^2 - y_2^2)) - b_{3,0,0,0}^i y_1 (3x_1^2 - y_1^2) + b_{1,0,2,0}^r (x_1 (x_2^2 - y_2^2) - 2x_2 y_1 y_2) + b_{3,0,0,0}^r x_1 (x_1^2 - 3y_1^2)$$

$$H_{XY}^{(3)} = -b_{1,0,2,0}^i (x_1 (x_2^2 - y_2^2) - 2x_2 y_1 y_2) - b_{3,0,0,0}^i x_1 (x_1^2 - 3y_1^2) - b_{1,0,2,0}^r (2x_1 x_2 y_2 + y_1 (x_2^2 - y_2^2)) - b_{3,0,0,0}^r y_1 (3x_1^2 - y_1^2)$$

$$H_{YX}^{(3)} = -b_{1,0,2,0}^i (x_1 (x_2^2 - y_2^2) - 2x_2 y_1 y_2) - b_{3,0,0,0}^i x_1 (x_1^2 - 3y_1^2) - b_{1,0,2,0}^r (2x_1 x_2 y_2 + y_1 (x_2^2 - y_2^2)) - b_{3,0,0,0}^r y_1 (3x_1^2 - y_1^2)$$

$$H_{YY}^{(3)} = b_{1,0,2,0}^i (2x_1 x_2 y_2 + y_1 (x_2^2 - y_2^2)) + b_{3,0,0,0}^i y_1 (3x_1^2 - y_1^2) - b_{1,0,2,0}^r (x_1 (x_2^2 - y_2^2) - 2x_2 y_1 y_2) - b_{3,0,0,0}^r x_1 (x_1^2 - 3y_1^2)$$