

EXPERIMENT NO: 13

Hypothetical using T-Test

Aim:

To test whether the average IQ score of a sample of students differs significantly from a population mean IQ score of 100.

Algorithm:

1. Null Hypothesis (H_0): The average IQ score of the sample is 100.
2. Alternative Hypothesis (H_1): The average IQ score of the sample is not 100.
3. Sample: Measure the IQ scores of 25 randomly selected students.
4. T-Test: Conduct a one-sample T-test to compare the sample mean to 100.
5. Decision Rule: Use a significance level of $\alpha = 0.05$.

Program:

```
[3]: import numpy as np
import scipy.stats as stats

[4]: np.random.seed(42)
sample_size = 25
sample_data = np.random.normal(loc=100, scale=15, size=sample_size)

[5]: population_mean = 100
sample_mean = np.mean(sample_data)
sample_std = np.std(sample_data, ddof=1)
n = len(sample_data)

[6]: t_statistic, p_value = stats.ttest_1samp(sample_data, population_mean)
alpha = 0.05

[7]: print(f"Sample Mean: {sample_mean:.2f}")
print(f"T-Statistic: {t_statistic:.4f}")
print(f"P-Value: {p_value:.4f}")

if p_value < alpha:
    print("Reject the null hypothesis: The average IQ score is significantly different from 100.")
else:
    print("Fail to reject the null hypothesis: There is no significant difference in average IQ score from 100.")

Sample Mean: 99.55
T-Statistic: -0.1577
P-Value: 0.8760
Fail to reject the null hypothesis: There is no significant difference in average IQ score from 100.
```

Result:

Thus, the Python code to perform a one-sample T-test to compare the sample IQ scores with the population mean is successfully executed.