



Università degli Studi di Cagliari

Facoltà di Scienze Matematiche, Fisiche e Naturali

Laurea Magistrale in Fisica

**Observation-Guided Simulation of the
Extragalactic Binary Gravitational Wave
Foreground in view of LISA**

Relatore:

Prof. Riccardo Murgia

Co-relatori:

Prof. Thomas Kupfer

Dott. Weitian Yu

Candidato:

Gabriele Todde

Abstract

In recent years ground-based interferometers such as LIGO and Virgo opened the high-frequency window of the rapidly evolving gravitational wave (GW) astronomy. Future space-based detectors, in particular the *Laser Interferometer Space Antenna* (LISA), will extend GW detection into the low-frequency band, where unresolved Galactic compact binaries such as double white dwarfs (DWDs) are expected to shape LISA's sensitivity curve significantly. The aim of this work is to investigate the expected GW foreground **signal** produced by *extragalactic* DWDs within 100Mpc from our Galaxy. We will do this by employing the *Compact Object Synthesis and Monte Carlo Investigation Code* (COSMIC) to generate synthetic populations representing the galaxies included in the observational survey **Gravitational Wave Galaxy Catalog** (GWGC). Each simulated binary contributes to the total GW strain, which is then combined across frequency bins and compared against the LISA sensitivity curve. **Our** results show that the total simulated extragalactic signal lies approximately one order of magnitude below LISA's detection threshold, indicating that the noise from the unresolved sources will not produce a measurable imprint on the sensitivity curve. Future developments could include the use of more advanced population synthesis codes and different prescriptions for binary evolution, or the extension of the analysis to larger observational galaxy catalogs and cosmological volumes.

Contents

Abstract	1
Introduction	5
1 Theoretical Framework	7
1.1 Double White Dwarfs (DWDs)	7
1.1.1 White dwarfs	7
1.1.2 The host galaxies	9
1.2 Gravitational Wave Theory	10
1.2.1 Gravitational waves as perturbations	10
1.2.2 The physical wave	11
1.2.3 Motion and geodesics	13
1.2.4 The quadrupole approximation	14
1.2.5 Energy carried by a gravitational wave	18
1.2.6 Evolution of a compact binary system	19
1.3 Gravitational Wave Interferometers	20
1.3.1 Michelson interferometer	21
1.3.2 The Laser Interferometer Space Antenna	24
2 Methods and Data	27
2.1 Stellar Populations Synthesis Codes	27
2.1.1 The starting point	27
2.1.2 Single star evolution	29
2.1.3 Binary star evolution	29
2.2 The COSMIC Code	30
2.2.1 Fixed population	30
2.2.2 Astrophysical population	32
2.3 The Gravitational Wave Galaxy Catalog	33
2.3.1 What it has vs what we need	33
2.3.2 Mass-luminosity relation	33
2.3.3 Mass-metallicity relation	35
3 Results and Discussion	37
3.1 Preliminary steps	37
3.1.1 Missing galaxies	37
3.1.2 The final fixed populations	41
3.2 The Signal Computation	42

CONTENTS	4
3.2.1 LISA’s frequency resolution	42
3.2.2 The final output	43
3.3 Results and Discussion	44
3.3.1 Robustness of the results	45
Conclusions and Future Perspectives	47
A	49

Introduction

In the context of the GW study, there are mainly two possible paths: the first is the **analysis of existing data** from experiments such as LIGO, Virgo and KAGRA ([Aasi et al. \[2015\]](#), Acernese et al. [2014], Akutsu et al. [2020]), with the purpose of detecting and characterizing the observable sources within their relative frequency bands; the second is the **forecasting approach**, which aims to characterize future experiments in order to better understand what types of sources they could be able to detect and how well.

One of the most important future detectors is LISA (Amaro-Seoane et al. [2023]), a NASA-ESA collaboration space-based GW detector which is scheduled to launch around 2030s (Luo et al. [2016]). LISA will be arranged as an equilateral triangle with 2.5 million kilometers long arms, placed in a heliocentric orbit, and will operate within the frequency range from approximately 0.1mHz to 1Hz. Among the typical sources that emit GW signal in this range are the **Compact Binary Coalescences (CBCs)**, and in our particular case we are interested in **Double White Dwarfs (DWDs)**.

Many studies have been made to explore the effects of different type of sources on the LISA's sensitivity curve. For example, we know that the unresolved binary Galactic systems produce a confusion GW foreground that shapes noticeably the curve close to its lowest threshold. While the Galactic contribution has been extensively investigated, less attention has been devoted to the possible extragalactic binaries foreground. Therefore, as a natural continuation, the purpose of this work is to estimate the GW foreground produced by the *extragalactic* DWDs in the nearby universe to an extension of $\sim 100\text{Mpc}$, using [the publicly available code COSMIC](#) (Breivik et al. [2020]) to generate synthetic astrophysical populations and the galaxies listed in the **Gravitational Wave Galaxy Catalog (GWGC)** (White et al. [2011]), an observational survey, as a reference. The potential impact of these sources in the tentative to characterize the expected LISA sensitivity and to avoid misinterpreting unresolved signals, are the factors that make this research so important.

In **Chapter 1: Theoretical Framework** we give a broad theoretical astrophysical framework, briefly discussing the birth and evolution of DWDs, both as single objects, and in binary systems, and the galactic environments that will host them. We also introduce the theoretical foundations of GW, starting from the linearized Einstein field equations, to derive the amplitude of the signal generated by an inspiralling binary system, and define the most important parameters. We then give a brief overview of how GW detectors work, to better understand what sources LISA will be able to see, what resolution and what sensitivity it will have and why in particular we are interested in DWDs.

In **Chapter 2: Methods and Data** we introduce the concept of stellar population syn-

thesis and the code COSMIC used for this purpose. We present its features, main parameters, and pipeline, and explain how it is used to generate full-size astrophysical populations of DWDs. In order to do this, we introduce the GWGC, and explain the procedure to generate the actual fixed populations, scale them to astrophysical full-size galaxies, and compute their GW signal. Moreover, we discuss how to take care of the lack of galaxies due to the catalog's incompleteness and the Milky Way coverage in the **Zone Of Avoidance (ZOA)**. Finally, we discuss the LISA frequency resolution problem, and how it affects the way we must process the GW signals simulated.

In **Chapter 3: Results and Discussion** we compute and show the total resulting signal on the LISA's sensitivity curve, and we demonstrate that the total simulated extragalactic signal lies at one order of magnitude below LISA's sensitivity curve (Robson et al. [2019]).

Finally, in **Chapter 4: Conclusions and Future Perspectives** we recap our work as a whole, discussing limitations, assumptions, possible extensions and follow-ups.

Chapter 1

Theoretical Framework

In this chapter we will cover all the theoretical foundations for this thesis. We will start by describing the astrophysics of DWDs and the galaxies hosting them. Next, we will go through the GW theory, starting from the linearization of Einstein's equations, up to the characteristic strain that we would expect from the DWDs. Finally, we will understand the functioning of GW detectors, using the Michelson interferometer as example, and introducing LISA's characteristics.

1.1 Double White Dwarfs (DWDs)

In this section, we will introduce the main astrophysical theory elements for the GW sources we want to simulate. First of all, we focus on white dwarfs, the compact remnants of low- and intermediate-mass stars, both as isolated objects or as components of binary systems. We then shift our discussion to galaxies, the large-scale environments in which these compact binaries form and evolve. The properties of galaxies, including their morphology, stellar populations, and dynamical structure, strongly influence the formation rates, spatial distribution, and evolution of white dwarf binaries. A clear understanding of both the compact objects themselves and their host galaxies is necessary for predicting the GW signals that can be observed.

1.1.1 White dwarfs

During its evolution, a star generates energy in its core through the nuclear fusion reactions, “burning” the elements in its core. During this process, which represents most of the star's life, the *radiation pressure* generated holds the star from the gravitational collapse. For most stars, this process begins with the fusion of hydrogen into helium, and when the core runs out of hydrogen, it contracts while the external layers expand: the luminosity is now spread in a larger surface, and since the *black body energy density* is $\propto T^4$, where T is the surface temperature, the star will appear colder and larger, as a *red giant* (Iben [1967]). If the star is sufficiently massive, core contraction raises the temperature enough to ignite helium fusion into heavier elements, leading to subsequent burning stages (Hoyle [1954], Burbidge et al. [1957]). In contrast, for medium or low massive stars ($M_* \leq 8M_\odot$), after the red giant phase the central pressure is initially insufficient to ignite heavier fusion, and the contraction continues until the *electron degeneracy*

pressure (a quantum mechanical effect arising from the Pauli exclusion principle) halts it (Bradt [2006], Chandrasekhar [1931]). In such stars, when the helium starts burning the emitted radiation is not sufficient to push back against the enormous gravitational pressure. At this point a runaway process can be initialized, in which the core keeps compressing, and heating, and the fusion rate increases. This positive feedback process induces a very fast helium burning called *helium flash*, that lifts the star back out of the degeneracy stage (Schwarzschild and Härm [1962]). The cycle of contraction and fusion continues until the core is composed of an element that cannot be fused under the prevailing conditions, either because the temperatures are too low, or because the reactions would no longer be exothermic (as in the case of iron core white dwarfs). Moreover, for low and intermediate-mass stars, intense winds and envelope ejections during late stages of the evolution can strip away the outer layers, leaving behind a naked, degenerate core. These types of medium-to-low massive star's remnant, supported entirely by electron degeneracy pressure and slowly cooling over time, are what we call **white dwarfs** (Bradt [2006]).

Binary systems

A great fraction of the white dwarfs in the universe are found DWDS systems, that show a wide range of orbital separations and periods, depending on their formation and evolutionary history (Gokhale et al. [2007]). In particular, in the Milky Way, DWDS are expected to be extremely common systems, many of which having orbital periods short enough to emit GWs in the milliHertz band (Nelemans et al. [2001]). Among the most relevant parameters for DWDS, typical values are:

- **Masses:** typically between $0.2 M_{\odot}$ and $1.2 M_{\odot}$, with the most common systems having total masses $M_{\text{tot}} \sim 0.6\text{--}1.0 M_{\odot}$.
- **Radii:** $\sim 0.01 R_{\odot}$ ($\approx 7 \times 10^3$ km), decreasing with increasing mass according to the mass-radius relation for degenerate matter.
- **Orbital periods:** between $\sim 10^4$ s (a few hours) and $\lesssim 300$ s (5 minutes) for the most compact systems.
- **Gravitational wave frequencies:** $f_{\text{GW}} = 2/P_{\text{orb}}$, hence in the range $\sim 0.1\text{--}10$ mHz for LISA-relevant binaries.

In some binaries, the DWDS are detached, and neither star fills its Roche lobe, while in **interacting systems** there might be mass transfers. In these cases, once one of the white dwarfs overflows its own Roche lobe, material flows through the inner Lagrangian point toward the companion. The nature of this transfer depends on the mass ratio: the transfer can be stable, leading to long-lived systems characterized by persistent mass accretion and bright electromagnetic counterparts, or it can become dynamically unstable, potentially resulting in the rapid merger of the two stars, which may produce a single more massive white dwarf, a neutron star, or even trigger a thermonuclear explosion if the critical conditions for a Type Ia supernova are met (Das et al. [2025], Shen and Bildsten [2009], Fink, M. et al. [2010], Shen et al. [2018]).

1.1.2 The host galaxies

Galaxies are the fundamental building blocks of the large-scale structure of the Universe. They host the vast majority of stars, stellar remnants, and interstellar matter, and therefore also the majority of DWDs, and represent the environment in which such objects are generated and evolve. At the same time, their distribution in space and their physical properties affect the detectability of the gravitational-wave signals produced by these binaries.

Morphological types

The role of galaxies as environment in which DWDs are born and evolve, and their properties are fundamental because they influence star evolution greatly. Therefore, it becomes necessary to categorize them, in order to schematize the most important differences from this perspective. The classical way to classify them is based on their morphology, as first organized by Hubble [1926] into the *Hubble sequence*. This classification broadly distinguishes between:

- **Elliptical galaxies (E):** They present smooth, featureless light profiles with nearly elliptical shape, and they are mostly populated by older stars and very little interstellar medium, and show little to no ongoing star formation. They range from nearly spherical (E0) to highly elongated ellipses (E7), depending on the eccentricity.
- **Lenticular galaxies (S0):** An intermediate class between elliptical and spiral galaxies, they present a central bulge and a disk but do not have prominent spiral arms. They also often contain older stars and have low gas content, but their flattened structure suggests a disk-like stellar kinematics.
- **Spiral galaxies (Sa, Sb, Sc, Sd):** such as our own, they are disk galaxies with spiral arms rooted in a central bulge. Early-type spirals (Sa) have tightly wound arms and larger bulges, whereas late-type spirals (Sc, Sd) have more open arms, smaller bulges, and more gas and young stars.
- **Irregular galaxies (Irr):** They lack of a defined shape, and are often rich in gas and undergoing intense star formation, possibly as a result of interactions or mergers.

For quantitative studies as the present one, morphological types can be conveniently parameterized by the **T-value** introduced in the *Third Reference Catalogue of Bright Galaxies* (de Vaucouleurs et al. 1991), which allows an easy mapping between the numerical index and the traditional Hubble sequence morphology. This system assigns values from $T = -6$ (E ellipticals) through $T \approx 0$ (S lenticulars), to $T \approx 10$ (irregulars), with intermediate integers corresponding to specific spiral subclasses (Sa, Sb, Sc, etc.). Note that the difference in the age of the stars populating the different morphological types illustrated translates in a systematic variation in the typical metallicity values, with early types generally being more metal-rich on average than late types. This metallicity fingerprint that distinguishes morphological types will be an important feature in the next chapters, where the stellar population properties will influence the compact binary formation and GW emission.

1.2 Gravitational Wave Theory

The DWDs described in the previous section are massive rotating objects, and therefore they produce ripples in spacetime that propagate as waves (Einstein [1918]). The purpose of this thesis is to analyze the GW produced by such systems, and it is thus necessary now to make a summary of the GW theory as a whole, to derive the characteristic strain that we will use to simulate the total signal. The treatment that follows primarily relies on Ferrari et al. [2020], particularly Chapters 12, 13, and 14. We will not go into the full mathematical derivations here, but they are presented in detail in the book.

1.2.1 Gravitational waves as perturbations

The typical approach to the study of GWs is to derive them as small perturbations of the foreground from the Einstein's equations:

$$G_{\mu\nu} = R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R, \quad (1.1)$$

which can be conveniently written as

$$R_{\mu\nu} = \frac{8\pi G}{c^4} \left(T_{\mu\nu} - \frac{1}{2}g_{\mu\nu}T \right), \quad (1.2)$$

where $G_{\mu\nu}$ is the Einstein tensor, $R_{\mu\nu}$ is the Ricci tensor, R is the Ricci scalar, $g_{\mu\nu}$ is the metric of spacetime, and $T_{\mu\nu}$ is the stress-energy tensor. As a foreground solution we can consider the flat spacetime described by the metric $\eta_{\mu\nu}$, to which the perturbation term appears as a fluctuation in the metric $|h_{\mu\nu}| \ll |\eta_{\mu\nu}|$: this is known as *weak field* approximation. Thus, the perturbed spacetime can be written as:

$$g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu}, \quad |h_{\mu\nu}| \ll |\eta_{\mu\nu}|.$$

With this metric, the equations 1.2 becomes

$$\{\square_F h_{\mu\nu} - \left[\frac{\partial^2}{\partial x^\lambda \partial x^\mu} h_\nu^\lambda + \frac{\partial^2}{\partial x^\lambda \partial x^\nu} h_\mu^\lambda + \frac{\partial^2}{\partial x^\nu \partial x^\mu} h_\lambda^\lambda \right]\} = -\frac{16\pi G}{c^4} \left(T_{\mu\nu} - \frac{1}{2}\eta_{\mu\nu}T \right).$$

Now, by requiring that the *weak-field* approximation remains satisfied for infinitesimal diffeomorphisms, and by choosing a coordinate system in which the *harmonic gauge condition*¹, defined as

$$\Gamma^\lambda = g^{\mu\nu} \Gamma_{\mu\nu}^\lambda = 0, \quad (1.3)$$

where $\Gamma_{\mu\nu}^\lambda$ are the *affine connections*, is satisfied, we can find that, up to first order in $h_{\mu\nu}$, the harmonic gauge condition is equivalent to

$$\frac{\partial}{\partial x^\mu} h_\rho^\mu = \frac{1}{2} \frac{\partial}{\partial x^\rho} h, \quad h = \eta^{\mu\nu} h_{\mu\nu} \equiv h_\nu^\nu. \quad (1.4)$$

¹It is an arbitrary coordinate condition which makes it possible to solve the Einstein field equations. It can be found by requiring that the linearized Einstein equations satisfy the D'Alembert equation.

After defining the *trace-reversed*² tensor as

$$\bar{h}_{\mu\nu} \equiv h_{\nu\mu} - \frac{1}{2}\eta_{\mu\nu}h,$$

we can finally write the linearized Einstein equations as

$$\begin{cases} \square \bar{h}_{\mu\nu} = -\frac{16\pi G}{c^4}T_{\mu\nu}, \\ \partial^\mu \bar{h}_{\mu\nu} = 0. \end{cases}$$

This form, and its twin with $T_{\mu\nu} = 0$, where the first equation becomes the D'Alembert equation, are relevant because they show that *a perturbation of a flat spacetime propagates as a wave travelling at the speed of light*. As in Electrodynamics, the solution of (1.5) can be written in terms of *retarded potentials*:

$$\bar{h}_{\mu\nu}(t, \mathbf{x}) = \frac{4G}{c^4} \int_V \frac{T_{\mu\nu}(t - \frac{|\mathbf{x}-\mathbf{x}'|}{c}, \mathbf{x}')}{|\mathbf{x}-\mathbf{x}'|} d^3x', \quad (1.5)$$

where V is the three dimensional source volume, $\mathbf{x}'$ is the distance of an element of the emitting source from the origin of a frame centered in the same point of the source, $\mathbf{x}$ is the distance between source and observer. It can be proved that the solutions in (1.5) automatically satisfy the harmonic gauge condition.

Harmonic gauge

It is important to notice that if the *harmonic gauge* condition is not satisfied in a reference frame, a new frame in which it is can always be found, by making an infinitesimal coordinate transformation

$$x'^\lambda = x^\lambda + \epsilon^\lambda,$$

provided that ϵ^λ satisfies the following equation:

$$\square_F \epsilon_\rho = \frac{\partial h_\rho^\beta}{\partial x^\beta} - \frac{1}{2} \frac{\partial h}{\partial x^\rho} = 0.$$

This is an inhomogeneous wave equation that can be solved to find the components ϵ_α , which identify the coordinate system in which the harmonic gauge condition is satisfied. Notice though, that the harmonic condition in (1.3) does not determine the gauge uniquely, but instead leaves some more gauge freedom to be used.

1.2.2 The physical wave

As we have seen, perturbations of flat spacetime satisfy a wave equation and a harmonic gauge condition, as in (1.5). The general solutions of these wave equation is a linear superposition of monochromatic plane waves, with a polarization tensor (or wave amplitude) $A_{\mu\nu}$ and a wave four-vector $\vec{k}$, such as

$$\square_F \bar{h}_{\mu\nu} = -A_{\mu\nu} \eta^{\alpha\beta} k_\alpha k_\beta e^{ik_\gamma x^\gamma} = 0.$$

²The name comes by noting that $\bar{h} = \eta^{\mu\nu} \bar{h}_{\mu\nu} = -h$.

Thus, neglecting the trivial solution $A_{\mu\nu} = 0$, we find

$$\eta^{\alpha\beta} k_\alpha k_\beta = 0,$$

which means that $\vec{k}$ is a null vector. If we also consider the harmonic gauge, we find a condition that imposes the orthogonality of the wave four-vector to the polarization tensor,

$$k_\mu A_\nu^\mu = 0.$$

The *wavefronts*, i.e. the spatial surfaces where $\bar{h}_{\mu\nu} = \text{const.}$ are the planes where $k_i x^i = \text{const.}$ Conventionally, k^0 is referred to as $\frac{\omega}{c}$, where ω is the frequency, thus

$$\vec{k}_0 = \left(\frac{\omega}{c}, \mathbf{k} \right),$$

where $\mathbf{k}$ is the wave three-vector orthogonal to the wavefront, and is related to the *wavelength* by $|\mathbf{k}| = 2\pi/\lambda$. Notice that, since the wave four-vector $\vec{k}$ is a null vector, it follows that

$$-(k^0)^2 + |\mathbf{k}|^2 = 0 \rightarrow \omega = ck_0 = c|\mathbf{k}|,$$

which gives the dispersion relation for a wave moving at the speed of light.

The TT gauge

In a one dimensional case, the wave equation can be written as

$$\left(-\frac{1}{c^2} \frac{\partial^2}{\partial t^2} + \frac{\partial^2}{\partial x^2} \right) \bar{h}_\nu^\mu = 0,$$

which generally has, as solution, an arbitrary function of $t \pm \frac{x}{c}$. If we consider, for example, a progressive wave $\bar{h}_\nu^\mu[f(t, x)]$, where $f(t, x) = t - \frac{x}{c}$, and apply the harmonic gauge condition, focusing only in the time-dependent part of the solution, we find the *first four* conditions:

$$\bar{h}_t^t = \bar{h}_t^x, \quad \bar{h}_x^t = \bar{h}_x^x, \quad \bar{h}_y^t = \bar{h}_y^x, \quad \bar{h}_z^t = \bar{h}_z^x. \quad (1.6)$$

As we have already discussed, there still is the freedom of making an infinitesimal coordinate change, and requiring that the harmonic gauge remains satisfied in the new coordinates generates *four more conditions*:

$$\bar{h}_x^t = \bar{h}_y^t = \bar{h}_z^t = \bar{h}_y^y + \bar{h}_z^z = 0,$$

and with the (1.6) follows

$$\bar{h}_x^x = \bar{h}_y^x = \bar{h}_z^x = \bar{h}_t^t = 0.$$

The only components which cannot be set to zero are $\bar{h}_y^z$ and $\bar{h}_y^y - \bar{h}_z^z$, because now we have completely used all the gauge freedom. It can be shown that, from all the above conditions, it follows that $\bar{h} \equiv h$, i.e. in this gauge the wave results to be *traceless*. Thus, a

plane GW propagating along the x -axis is characterized by only two non-zero functions h_{zy} and $h_{yy} = -h_{zz}$:

$$h_{\mu\nu}^{TT} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & h_{yy} & h_{yz} \\ 0 & 0 & h_{yz} & -h_{yy} \end{pmatrix}. \quad (1.7)$$

The GW only has two **physical degrees of freedom** which correspond to two polarization states. This gauge is called **TT gauge** because of the *transverse traceless* nature: $h_{\mu\nu}$ is traceless, thus $h = 0$, and transverse, since the components of $h_{\mu\nu}$ along the direction of propagation are null (in this case $h_{\mu x} = 0$).

1.2.3 Motion and geodesics

Free test particles in General Relativity move along geodesic, which means that, in terms of a world line $x^\mu(\tau)$ parametrized by proper time τ , they satisfy the *geodesic equation*:

$$\frac{d^2 x^\alpha}{d\tau^2} + \Gamma_{\mu\nu}^\alpha \frac{dx^\mu}{d\tau} \frac{dx^\nu}{d\tau} \equiv \frac{du^\alpha}{d\tau} + \Gamma_{\mu\nu}^\alpha u^\mu u^\nu = 0, \quad (1.8)$$

where $u^\alpha = \frac{dx^\alpha}{d\tau}$ is the particle velocity. By imposing the rest condition, $u^\alpha = (1, 0, 0, 0)$, to the (1.8) and considering the TT gauge component prescriptions for $h_{\mu\nu}$, we find that a particle initially at rest is not accelerated by the passage of a wave, but remains at a fixed coordinate position: the motion of a single particle is not affected by the GWs.

The geodesic deviation

If we consider the relative motion of close particles, the situation is different. This relative motion is governed by the *geodesic deviation*, which quantifies how two very close geodesics deviate from one another. Although, as we have seen, the coordinates of the two particles (and thus their difference) do not change when a GW passes, this does not hold true for their *proper distance*

$$\Delta l = \int ds \neq \text{const.}$$

This apparent contradiction arises from the fact that the coordinate difference is not a *tensorial* quantity, and therefore is not suited to describe properly a physical process. This is the reason we consider the geodesic deviation, which provides a tensorial formulation of the relative acceleration between the two particles:

$$\frac{D^2 \delta x^\alpha}{d\tau^2} \equiv (\Delta_{\vec{t}}(\Delta_{\vec{t}} \delta x))^\alpha = R_{\beta\mu\nu}^\alpha t^\beta t^\mu \delta x^\nu, \quad (1.9)$$

where

$$R_{\alpha k \lambda \mu} = \frac{1}{2}(g_{\alpha\mu, \lambda k} + g_{k\lambda, \mu\alpha} - g_{\alpha\lambda, \mu k} - g_{k\mu, \lambda\alpha}) \quad (1.10)$$

is the Riemann curvature tensor, $\vec{t}$ is the tangent vector to one of the geodesic, $\vec{\delta x}$ is a deviation vector between the two geodesics and $\frac{D}{d\tau}$ is the covariant derivative over the

proper time τ . It is important to note that in the presence of a GW the Riemann tensor is never zero in any reference frame, and consequently neither is the geodesic deviation. In contrast, the quantity $\frac{d^2\delta x^i}{d\tau^2}$ (which is not a tensorial quantity) vanishes in the TT frame.

Knowing this, in order to analyze this effect in more detail we can choose to integrate the (1.9) in a *Locally Inertial Frame*, LIF $\{\xi^\alpha\}$, centered on one of the two particles, where the metric can be approximated close to the origin as Minkowski up to quadratic, negligible, corrections:

$$ds^2 = \eta_{\alpha\beta} d\xi^\alpha d\xi^\beta + O(|\xi|^2).$$

In this frame, the Riemann tensor in (1.10), which depends on second derivatives of the metric perturbation $h_{\mu\nu}$ (with $g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu}$), is not only covariant, but actually *invariant* under infinitesimal coordinate transformations from a generic LIF to the TT frame. This is especially useful because in the TT frame $h_{\mu\nu}^{TT}$ takes a particularly simple form, as we have seen in the (1.7) example, simplifying also the expression for $R_{\alpha k \lambda \mu}^{TT}$. Assuming that the wave travels along the ξ^1 direction, the solutions for the geodesic deviation equation for two close particles are the non zero components of

$$\delta \xi^j = \delta \xi_0^j + \frac{1}{2} \eta^{ji} h_{ik}^{TT} \delta \xi_0^k,$$

where $\delta \xi_0^j$ represent the initial, constant separation between the particles. As previously discussed, the non-zero components of the strain tensor h_{ik}^{TT} correspond to two possible polarizations of the GW. Their distinct effects on a ring of close particles are visually represented in Figure 1.1.




Figure 1.1: Effect of the two GW polarizations on a ring of free-falling particles, caused by a wave moving in the direction perpendicular to the page: blue and red dots represent two opposite phases of the wave.

1.2.4 The quadrupole approximation

We have discussed the effects of the passage of a GW on the relative motion of close particles; in this section we focus on another important problem: *how such waves are*

generated. Gravitational radiation is a consequence of variations in the spacetime curvature, and it generates from time-dependent distributions of mass and momentum. But not every motion of matter can produce GW signal, as we will see: conservation laws of general relativity, impose strict constraints on the possible multipole moment that can contribute to the tensorial metric perturbations.

The weak-field, slow-motion approximation

We already introduced previously the weak-field limit, which allows us to write the metric as a perturbed Minkowski $g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu}$, with $|h_{\mu\nu}| \ll 1$, and we found the solutions as in (1.5). Until now, we have only solved these equations with a null stress-energy tensor. If we assume that the region, of size ϵ , where the source of the signal is confined, is much smaller than the wavelength of the emitted gravitational radiation, $\lambda_{GW} = \frac{2\pi c}{\omega}$, we find the following condition:

$$\frac{2\pi c}{\omega} \gg \epsilon \rightarrow \epsilon\omega \ll c \rightarrow v_{\text{typical}} \ll c.$$

The typical velocities of the system are much smaller than the speed of light: this is the so called *slow motion approximation*. If we now use the solution found using the retarded potentials in (1.5), by making use of the Fourier transforms it can be found that the GW signal emitted by the source, to leading order in the weak-field, slow-motion approximation is

$$\bar{h}_{\mu\nu}(t, r) = \frac{4G}{rc^4} \int_V T_{\mu\nu}(t - \frac{r}{c}, \mathbf{x}') d^3x. \quad (1.11)$$

The quadrupole formula

The integral in (1.11) can be simplified with some expedients. First of all, by making use of the conservation law that applies to the stress-energy tensor

$$\frac{\partial T^{\mu\nu}}{\partial x^\nu} = 0, \quad \rightarrow \quad \frac{1}{c} \frac{\partial T^{\mu 0}}{\partial t} = -\frac{\partial T^{\mu k}}{\partial x^k}, \quad \mu = 0, \dots, 3, \quad k = 1, 2, 3,$$

a smart use of the Gauss' theorem (reminding that the stress-energy tensor is null on the surface of the source), it can be found that

$$\bar{h}^{\mu\nu} = 0.$$

Secondly, we can use the **tensor-virial theorem**

$$\frac{1}{c^2} \frac{\partial^2}{\partial t^2} \int_V T^{00} x^k x^n d^3x = 2 \int_V T^{kn} d^3x, \quad k, n = 1, 2, 3,$$

in the definition of the **quadrupole moment tensor**

$$q^{kn}(t) = \frac{1}{c^2} \int_V T^{00}(t, \mathbf{x}) x^k x^n d^3x,$$

we finally find that (1.11) can be re-written as

$$\begin{aligned}\bar{h}^{\mu 0} &= 0, \\ \bar{h}^{ik}(t, r) &= \frac{2G}{c^4 r} \frac{d^2}{dt^2} q^{ik} \left(t - \frac{r}{c}\right),\end{aligned}\tag{1.12}$$

known as the **quadrupole formula**. This formulation describes the GW signal emitted by a gravitating system that evolves in time, whatever the mass-energy distribution looks like, as long as $\ddot{q}^{ik} \neq 0$. Note that in this approximation, the only component of the stress-energy tensor that generates the metric perturbation is T^{00} , and that the c^{-4} makes the characteristic strain extremely small:

$$\frac{G}{c^4} \sim 10^{-49} \frac{s^2}{g \text{ cm}}.$$

Transform to the TT gauge

The solution in (1.12) represents a spherical wave far from the source, which locally looks like a plane wave. It can also be expressed, after an infinitesimal coordinate transformation that preserves the harmonic gauge. This projection in the TT gauge is performed by defining symmetric and transverse *projector operators* as

$$P_{jk} \equiv \delta_{jk} - n_j n_k,$$

to then define the **transverse-traceless projector**

$$P_{jkmn} \equiv P_{jm} P_{kn} - \frac{1}{2} P_{jk} P_{mn},\tag{1.13}$$

which “extracts” the transverse-traceless part of a rank two tensor on the three-dimensional Euclidean space. This way we can, for example, define

$$h_{jk}^{TT} = P_{jkmn} \bar{h}_{mn},$$

that, if applied to the (1.12), gives

$$\begin{aligned}\bar{h}_{\mu 0}^{TT} &= 0, \\ \bar{h}_{ik}^{TT}(t, r) &= \frac{2G}{c^4 r} \frac{d^2}{dt^2} Q_{ik}^{TT} \left(t - \frac{r}{c}\right),\end{aligned}\tag{1.14}$$

where $Q_{jk}^{TT} = P_{jkmn} q_{mn}$ is the **transverse-traceless part of the quadrupole moment**. As we will see, to compute the “luminosity” of a GW source, it is useful to define the **reduced quadrupole moment**

$$Q_{jk} \equiv q_{jk} - \frac{1}{3} \delta_{jk} q_m^m,\tag{1.15}$$

which is traceless by definition.

Gravitational waves from a binary system

As an example, we can consider a binary system where two masses m_1 and m_2 are in a circular orbit around the common center of mass. The typical parameters are the total mass

$$M \equiv m_1 + m_2,$$

the *reduced mass*

$$\mu \equiv \frac{m_1 m_2}{M},$$

and, by placing the origin of the coordinate frame in the center of mass,

$$l_0 = r_1 + r_2, \quad r_1 = \frac{m_2}{M} l_0, \quad r_2 = \frac{m_1}{M} l_0, \quad \omega_K = \sqrt{\frac{GM}{l_0^3}}.$$

Now, after defining the coordinates of the masses (x_1, y_1) and (x_2, y_2) in an orbital plane, orthogonal to the z -axis, as functions of all the above parameters, it becomes possible to compute the 00-component of the stress-energy tensor as

$$T^{00} = c^2 \sum_{n=1}^2 m_n \delta(x - x_n) \delta(y - y_n) \delta(z),$$

and thus all the non-vanishing components of q_{jk} . From this we can find Q_{jk} , Q_{jk}^{TT} , and finally

$$h_{ij}^{TT}(t, r) = -\frac{A}{r} A_{ij}^{TT}\left(t - \frac{r}{c}\right), \quad (1.16)$$

where³

$$\begin{aligned} A &= \frac{4\mu M G^2}{l_0 c^4}, \\ A_{ij}^{TT}\left(t - \frac{r}{c}\right) &= P_{ijkl} A_{kl}\left(t - \frac{r}{c}\right), \\ A_{kl}(t) &= \begin{pmatrix} \cos 2\omega_K t & \sin 2\omega_K t & 0 \\ \sin 2\omega_K t & -\cos 2\omega_K t & 0 \\ 0 & 0 & 0 \end{pmatrix}. \end{aligned} \quad (1.17)$$

Using typical values, such as the ones from the PSR 1913+16 assuming it had a circular orbit, by defining

$$h_0 = \frac{A}{r},$$

and using (1.17), we find

$$h_0 \sim 5 \times 10^{-23}, \quad (1.18)$$

as an estimate of a typical GW signal amplitude generated by a binary system.

³This shows that a *binary system in circular orbits emits waves at twice the orbital frequency*

1.2.5 Energy carried by a gravitational wave

The stress-energy tensor conservation laws that we know in Special Relativity

$$T^{\mu\nu}_{;\nu} = 0,$$

unfortunately do not hold in General Relativity. Indeed, the Stress-energy tensor describes the energy and momentum of the sources of the gravitational field, but not of the gravitational field itself: to have a proper conservation we would need to consider both contributions. The equivalence principle states that the effect of the gravitational field vanishes in a LIF centered on a particle; if there was a tensor describing it, it would be null too in this frame. But if a tensor field is null in one frame, than it must be in all the other too: for this reason, there cannot be anything like a true gravitational field tensor.

Stress-energy pseudo-tensor

Since no tensor can represent the gravitational field itself, we are going to introduce a **stress-energy pseudo tensor**, where a pseudo-tensor is a mathematical object that behaves like a tensor only under linear coordinate transformations. It is possible to define procedures of *integration* or *averaging* involving it, which allow to construct well-defined gauge-invariant quantities. In particular, it is possible to define a *local* notion of energy and momentum in the weak field approximation, on the assumption that the characteristic length-scale λ of the perturbation is much smaller than the characteristic length-scale L of the foreground: $\lambda/L \ll 1$. In this regime, it can be shown that if averaged over multiple wavelengths λ , the stress-energy pseudo-tensor transforms as a tensor for coordinate transformations linear in h (i.e. of order $O(h)$),

$$\langle t^{\mu\nu} \rangle,$$

and is therefore *suitable to describe energy and momentum carried by the perturbation*.

Gravitational wave luminosity

In general, it can be shown that the energy flowing across a unit surface orthogonal to the direction x' per unit time is given by the $0x'$ component of the stress-energy tensor times the speed of light. Similarly, the energy flux of a GW propagating in the same direction is given by the same component of the stress-energy pseudo-tensor averaged over several wavelengths, and can be derived as

$$\frac{dE_{GW}}{dtdS} = c \langle t^{0x'} \rangle = \frac{c^3}{16\pi G} \langle (\dot{h}_+(t, x'))^2 + (\dot{h}_x(t, x'))^2 \rangle.$$

where h_+ and h_x are the analog of h_{yy} and h_{yz} in the (1.7), and represent the metric perturbation polarizations. In the TT frame, by directly substituting what we have found in (1.14), we find

$$\frac{dE_{GW}}{dtdS} = \frac{G}{8\pi c^5 r^2} \left\langle \sum_{jk} \left(\ddot{Q}_{jk}^{TT} \left(t - \frac{r}{c} \right) \right)^2 \right\rangle,$$

and after defining the **gravitational wave luminosity** as

$$L_{GW} = \frac{dE_{GW}}{dt} = \int \frac{dE_{GW}}{dtdS} dS,$$

and writing Q^{TT} as a function of (1.15), by using the TT projectors in (1.13) and their properties, it can be shown that the energy associated to a GW can be written as the **luminosity quadrupole formula**:

$$L_{GW}(t) = \frac{G}{5c^5} \left\langle \left(\ddot{Q}_{ij} \left(t - \frac{r}{c} \right) \ddot{Q}_{ij} \left(t - \frac{r}{c} \right) \right) \right\rangle, \quad (1.19)$$

as was derived by Einstein in Einstein [1918].

1.2.6 Evolution of a compact binary system

By applying what we have found in the previous subsection to real astrophysical sources of GW, in this section we will find the amplitude that we are going to use for this thesis work to generate the total GW signal. In particular, we can apply the general results we found for two objects in a circular orbit in (1.17) to compute the quadratic sum of the components of $\ddot{Q}_{ij}$ in (1.19), and find the GW luminosity of a compact binary system as

$$L_{GW} = \frac{32}{5} \frac{G^4}{c^5} \frac{\mu^2 M^3}{l_0^5}$$

Remember that this represents an average over several wavelengths, and thus, of course, over many orbital periods (since $\omega_{GW} = 2\omega_K$), which makes it applicable only for systems that do not change significantly over a large number of periods (very far from merger), an assumption that is called *adiabatic approximation*. The energy emitted in GW comes from the total energy of the system, which is in part kinetic and in part potential energy. By requiring the time derivative of the two to be equal, we can find how the binary parameters must change accordingly. For example, the distance between the two objects shortens with time as

$$l_0(t) = l_0^{in} \left(1 - \frac{t}{t_c} \right)^{1/4}, \quad \text{where} \quad t_c \equiv \frac{5}{256} \frac{c^5}{G^3} \frac{(l_0^{in})^4}{\mu M^2}. \quad (1.20)$$

Using Kepler's laws it is also possible to translate this in a time dependence on the frequency:

$$\omega_K(t) = \omega_K^{in} \left(1 - \frac{t}{t_c} \right)^{-3/8}, \quad \text{where} \quad \omega_K^{in} \equiv \frac{(GM)^{1/2}}{(l_0^{in})^{3/2}}. \quad (1.21)$$

or the period

$$P(t) = P^{in} \left(1 - \frac{t}{t_c} \right)^{3/8}.$$

Signal from inspiralling compact objects

Knowing how the orbital parameters evolve with time allows us to find the GW signal for an *inspiralling compact object*. We have already found in (1.16) the general strain solution for a binary system, where its useful to redefine the instantaneous amplitude as

$$h_0 = \frac{A}{r} = \frac{4\mu MG^2}{l_0 c^4 r},$$

where r is the distance from the source, and l_0 the distance between the two objects. Moreover, l_0 can be written in terms of the angular frequency, using Kepler's law and (1.21), to get

$$h_0(t) = \frac{4\pi^{2/3} G^{5/3} M_{ch}^{5/3}}{r c^4} \nu_{GW}^{2/3}(t),$$

where

$$M_{ch} = \mu^{3/5} M^{2/5} = \frac{(m_1 m_2)^{3/5}}{M^{1/5}},$$

is the **chirp mass**, and

$$\nu_{GW}(t) = \frac{\nu_{GW}^{in} t_c^{3/8}}{(t_c - t)^{3/8}},$$

is the GW frequency. By substituting t_c as defined in (1.20), we finally find the signal emitted during the inspiral as

$$h_{ij}^{TT}(t, r) = -h_0 \left(t - \frac{r}{c} \right) P_{ijkl} A_{kl} \left(t - \frac{r}{c} \right),$$

where

$$h_0(t) = \frac{4\pi^{2/3} G^{5/3}}{c^4} \frac{M_{ch}}{r} (M_{ch} \nu_{GW}(t))^{2/3} \quad (1.22)$$

is the wave amplitude that we are going to refer to in this thesis work, when we calculate the signal from the simulated binaries.

1.3 Gravitational Wave Interferometers

The typical GW amplitude for a binary system is very small, and this makes it quite a hard task to detect it. The most sensitive instrument capable of such measurements are the interferometric detectors, that are generally based on the same principle of the simple Michelson interferometer: by splitting a laser beam along two perpendicular arms and recombining it, they detect the small phase shifts induced by a passing GW. In this section, which is mainly based on the chapter 9 of Maggiore [2007], we explain the general working principle of interferometers using the Michelson interferometer as an example. After this, we will talk briefly about LISA's characteristics, and what makes it interesting for the purpose of this thesis.




Figure 1.2: Schematic representation of a Michelson interferometer. A laser beam is split by a half-transparent mirror into two orthogonal arms of length L_A and L_B . The beams reflected from the mirrors are recombined at the detector (screen), where, if a GW has passed, they produce an interference pattern. Variations in the optical path length of one arm (through the movable mirror) shift the observed fringes. The source for this image can be found [here](#).

1.3.1 Michelson interferometer

The Michelson interferometer is an optical instrument invented in 1887 in occasion of the Michelson-Morley experiment, with the purpose of measuring changes in the travel time of light due to its relative motion with respect to *ether*. It uses a *beam splitter* to separate a single laser beam into two independent, and perpendicular paths, that end up recombining via the superposition principle before meeting the photoelectric detector or camera, as represented in **Figure 1.2**. The laser is used to have a monochromatic light source, with a frequency ω_L , a wave-vector $k_L = \omega_L/c$ and a wavelength $\lambda_L = 2\pi/k_L$. Therefore, referring to **Figure 1.2**, we label as A and B respectively the x and y arms of the interferometer, with lengths L_A and L_B . If t_0 is the time the beam reaches the beam splitter the first time, it will come back to it from the two paths at time $t = t_0 + 2L_A/c$ and $t' = t_0 + 2L_B/c$. The detector only sees one single recombined beam, not two beams arriving at different times, so in its perspective if one had to cross a longer distance than the other, it must have started its journey before the other, meaning $t \equiv t'$, and the two

beams start at $t_0^{(A)} = t - 2L_A/c$ and $t_0^{(B)} = t - 2L_B/c$. If we set the beam splitter at the origin of the frame, this initial time difference translates in a phase difference between the two separate beams. If the phase is conserved during the free propagation, there are some factors acquired by the field during the reflections and refractions, and the overall effect gives the two electric fields that will recombine at the beam splitter as

$$\begin{cases} E_1 = -\frac{1}{2}E_0 e^{-i\omega_L t + 2ik_L L_A} \\ E_2 = -\frac{1}{2}E_0 e^{-i\omega_L t + 2ik_L L_B}, \end{cases} \quad (1.23)$$

The total electric field is the sum of the two, and after some re-writing it can be shown that the power measured by the photodetector is

$$|E_{out}|^2 = E_0^2 \sin^2[k_L(L_A - L_B)], \quad (1.24)$$

therefore any variation in the lengths of the two arms corresponds to a variation in the power measured at the photodetector. Note that, since $k_L = 2\pi/\lambda_L$, the (1.24) becomes

$$|E_{out}|^2 = \frac{1}{2}E_0^2 \left[1 - \cos\left(\frac{4\pi}{\lambda_L}\Delta L\right) \right],$$

therefore the detector measures a value depending on the cosine, which in turn depends on ΔL : if there is no GW, a null measure should be expected.

Gravitational waves and TT gauge

The TT gauge describes the position of *free falling* objects, that does not change when a GW passes. In order to replicate this condition the mirrors of the interferometers should also be “free falling”, but instead their fall is halted by suspensions. This apparent problem is solved when considering that the effects of GWs are perpendicular to their direction of motion, and therefore the mirrors need to be decoupled in the horizontal plane, rather than the vertical. For this reason, the mirrors are suspended, and as far as the motion in the horizontal plane is concerned they can be taken to be following the geodesics of the time-dependent part of the gravitational field. Thus, in the TT gauge the coordinates of the beam splitter (at the origin) and of the mirrors (at $(L_A, 0)$ and $(0, L_B)$) are not affected by the passage of a wave, but the propagation of light in the arms is. Assuming a *plus-polarized* GW moving in the z direction, perpendicular to the plane defined by the two arms, we have

$$h_+(t) = h_0 \cos \omega_{gw} t,$$

and the spacetime interval in the TT frame will be given by

$$ds^2 = -c^2 dt^2 + [1 + h_+(t)]dx^2 + [1 - h_+(t)]dy^2 + dz^2.$$

Since photons follow the null geodesic, we can set $ds = 0$, and, considering the arm in the x direction as an example, we get

$$dx = \pm c dt \left[1 - \frac{1}{2}h_+(t) \right],$$

where the sign depends on the direction of motion with respect to the origin. These can be used to write the length of the arms as the time integral from the beam splitter at t_0 , to the mirror at time t_1 , or from this time back to the beam splitter, at t_2 , in order to find the total time interval in which the light has traveled along the arms. By neglecting terms $O(h^2)$, and using some trigonometric identities, we can find the time interval for the x -axis arm as

$$t_2 - t_0 = \frac{2L_A}{c} + \frac{L_A}{c} h(t_0 + L_A/c) \frac{\sin(\omega_{gw} L_A/c)}{(\omega_{gw} L_A/c)}, \quad (1.25)$$

where

$$\text{sinc}\left(\frac{\omega_{gw} L}{c}\right) \equiv \frac{\sin(\omega_{gw} L/c)}{(\omega_{gw} L/c)} \xrightarrow{\frac{\omega_{gw} L}{c} \rightarrow 0} 1.$$

Therefore, when the period of the GW is large with respect to the time it takes the light to travel along the arm, the Δt shift in the travel time $t_2 - t_0$, with respect to the unperturbed case is simply $h(t_1)L_A/c$. In the opposite case, when $\omega_{gw} L_A/c \gg 1$, $h(t)$ will change sign many times during the travel time, making its net contribute ~ 0 . Now, from (1.25) and its y -axis counterpart, it is possible to find the two different starting times⁴ $t_0^{(x)}$ and $t_0^{(y)}$. By substituting them in the electric field expressions in (1.23) we can find

$$\begin{cases} E^{(x/y)} = \mp \frac{1}{2} E_0 e^{-i\omega_L t_0^{(x/y)}} \\ \quad = \mp \frac{1}{2} E_0 e^{-i\omega_L \left(t - \frac{L_{A/B}}{c}\right) + i\Delta\phi_{x/y}(t)}, \end{cases} \quad (1.26)$$

where

$$\Delta\phi_{x/y} = \pm h_0 \frac{\omega_L L_{A/B}}{c} \text{sinc}\left(\omega_{gw} \frac{L_{A/B}}{c}\right) \cos\left[\omega_{gw} \left(t - \frac{L_{A/B}}{c}\right)\right], \quad (1.27)$$

is the phase difference from the unperturbed case in the two arms. Generally, L_A and L_B are made to be as similar as possible, therefore we can replace them with $L = (L_A + L_B)/2$ in (1.27). We still consider the minimal differences between the two in (1.26), which can be rewritten⁵ as

$$E^{(x/y)}(t) = \mp \frac{1}{2} E_0 e^{-i\omega_L(t - 2L/c) \pm i\phi_0 + i\Delta\phi_{x/y}(t)},$$

where

$$\phi_0 = k_L(L_A - L_B),$$

is a phase parameter that the experimenter can adjust. The strategy to build a “null instrument”, which records zero signal when the GW is not passing, by setting $L_A = L_B \rightarrow \phi_0 = 0$, turns out to be failing, since it makes the instrument insensitive to calibration uncertainties. At the same time, setting the instrument to a phase where the

⁴Note that if $L_A = L_B = L$, and $\frac{\omega_L}{c} \ll 1$, the difference between the two would be

$$t_0^{(y)} - t_0^{(x)} = \frac{2L}{c} h\left(t - \frac{L}{c}\right) \text{sinc}\left(\omega_L/c\right) \approx \frac{2L}{c} h\left(t - \frac{L}{c}\right).$$

⁵By using $2L_{A/B} = 2L \pm (L_A - L_B)$.

variation is maximum, as for $\phi_0 = \pi/4$, makes the instrument too sensible, so much that it will not be able to distinguish a signal from the laser power fluctuations. Therefore, depending on how this parameter is adjusted the instrument can be more or less sensible to signal variations, and the best value tends to be between these two extremes.

1.3.2 The Laser Interferometer Space Antenna

Already existing detectors such as LIGO and Virgo (Abbott et al. [2016], Abbott et al. [2019]) have pioneered the technique we have just shown in the high-frequency band (Amaro-Seoane et al. [2023], Acernese et al. [2014]). The purpose for future space-based interferometers such as LISA is to extend it to the low-frequency regime (approximately within 0.1mHz and 1Hz), opening a new observational window on astrophysical and cosmological sources. Indeed, space-based detectors such as LISA will be sensitive to DWDs with periods of a few hours down to a few minutes (Baker et al. [2019]). At the shortest periods, these binaries are in very tight orbits, with separations of only a few 10^4 – 10^5 km, comparable to Earth’s diameter. Between the binary systems, the interacting ones can have extremely short orbital periods ($\lesssim 10$ minutes) and are strong, nearly monochromatic GW sources over LISA’s lifetime, which in this work is considered of ~ 4 yr (Baker et al. [2019]). Detached systems with periods in the mHz range are also important, as they provide long-lived, stable signals that can serve as calibration sources or “verification binaries” (Kupfer et al. [2018], Kupfer et al. [2024]) for space-based detectors. Given their abundance and GW luminosity, binary white dwarfs are expected to be the most numerous class of resolvable sources for LISA, and their unresolved population will form a significant component of the low-frequency GW foreground.

Instrument description

Basing new detectors in space allows to eliminate sources of noise such as seismic noise and gravity gradients from local movements, allowing to explore a frequency regime which contains precious information on many interesting astrophysical sources, such as DWDs, extreme mass-ratio inspirals and massive black hole mergers. In LISA’s case, the arms are not going to be physical structures, but 2.5×10^6 km separations between three identical spacecrafts at the vertices of an equilateral triangle (Baker et al. [2019]). Each one of the spacecraft will host two free-falling test masses, isolated from all non-gravitational interactions by a drag-free control system. The spacecrafts will trail by about 20° , with the triangular constellation inclined by 60° to the ecliptic plane, following the earth around the sun (Baker et al. [2019]). With this configuration, it is possible to keep the long baseline stable, while having continuous source visibility, and a good localization (granted by the triangular shape).

Frequency

LISA’s frequency band roughly covers the low range $0.1 \text{ mHz} \lesssim f \lesssim 1 \text{ Hz}$ (compared to LIGO’s frequency of 10 Hz to 10 kHz (Martynov et al. [2016])), corresponding to objects in much wider orbits and potentially much heavier than those that LIGO is searching for, opening up the to the detection of a wider range of GW sources. In this frequency range also reside most DWDs. In the low frequencies, the range is limited




Figure 1.3: LISA sensitivity curve in the characteristic strain representation. The solid green line shows the instrumental noise curve, while the dashed black line represents the LISA curve as shaped by the foreground contribution from unresolved Galactic binaries. Different astrophysical sources that emit GW signal in this band are indicated: massive black hole binaries (MBHBs), extreme mass-ratio inspiral (EMRI), stellar-mass black hole binaries, verification binaries, and Galactic binaries with signal-to-noise ratio larger than 7. The source for this image can be found [here](#) (Amaro-Seoane et al. [2023])

by the acceleration noise on the test masses: even in space, the “free” test masses are not completely isolated, but could be influenced by solar radiation and winds, and the “drag-free” micro-propellers that compensate these disturbances make noise under the 0.1 mHz (Baker et al. [2019]). Also, in order to detect very slow wave, LISA needs to observe it for at least one complete cycle, and if the wave has a period of months, the signal could become indistinguishable from a slow relative position change between the three satellites. At the high frequencies, the main limits are due to the arms length, which make LISA unable to detect waves much shorter than that scale, and the power limits of the laser, which makes the photon count Poissonian noise more relevant at high frequencies (Baker et al. [2019]). LISA’s sensitivity curve combines instrumental noise and the astrophysical foreground, including Galactic foreground and Galactic binary systems, as shown in Figure 1.3 (Amaro-Seoane et al. [2023]). As far as the frequency resolution⁶ of a coherent observation is concerned, the limits are not due to hardware, but the actual total observational time: in a Fourier analysis, this is set by the inverse of

⁶Not to be confused with the sensitivity of the instrument, which states how faint of a signal it can detect.

the total observation time:

$$\Delta f \approx \frac{1}{T_{\text{obs}}}. \quad (1.28)$$

For a nominal mission duration of $T_{\text{obs}} \simeq 4$ years, $\Delta f \approx 8 \times 10^{-9}$ Hz, allowing very fine separation of closely spaced monochromatic sources.

Why double white dwarfs

From a scientific perspective, double white dwarfs (DWDs) provide an excellent laboratory for gravitational-wave studies:

- DWDs are typically found in compact orbits, with corresponding GW frequencies that lie within LISA's observational band;
- They are among the most common compact binary systems in the universe, often representing a typical outcome of stellar binary evolution. When a large number of such sources cannot be individually resolved, their signals overlap stochastically, producing a foreground;
- In the circular orbit approximation, their strain can be described by a simple and well-defined expression [Eq. (1.22)], which can be readily simulated and used to generate coherent synthetic populations within astrophysical models;
- As previously discussed, Galactic binaries are already expected to impact LISA's sensitivity curve, making extra-galactic populations a realistic and relevant candidate to investigate.

Galactic binaries are therefore expected to generate a confusion foreground that significantly impacts the LISA sensitivity curve, primarily due to their proximity and large number (Hils et al. [1990], Nelemans et al. [2001], Amaro-Seoane et al. [2023]). The aim of this work, however, is to determine whether *extra-galactic* DWDs, despite their much greater distances, can contribute significantly to an overall detectable stochastic GW foreground in the LISA band. The combination of their abundance, characteristic signal strength, and astrophysical importance makes extra-galactic DWDs an optimal and compelling target for LISA's mission.

Chapter 2

Methods and Data

In anticipation of the launch of a space-based gravitational-wave detector, the study of the expected signal from DWDs entirely relies on simulations. To this end, binary population synthesis codes have been developed as dedicated tools, whose accuracy benefits significantly from high-quality electromagnetic observations, especially in the era of large-scale surveys such as the [Sloan Digital Sky Survey \(SDSS\)](#), the [Zwicky Transient Facility \(ZTF\)](#), the [Dark Energy Spectroscopic Instrument \(DESI\)](#), and [Euclid](#). These observations not only provide complementary insights but also serve as crucial constraints to guide and validate the simulations, ensuring that the predicted populations are consistent with the systems already observed. In this chapter, we will first discuss population synthesis codes, then introduce in detail the COSMIC code used in this work. Finally, we will present the astrophysical surveys that we adopted as reference for the simulation of our sources.

2.1 Stellar Populations Synthesis Codes

Generating a synthetic population of stars is a very complex task, that includes multiple steps, each involving important choices. *First*, we need to choose a starting point: we could start from the very beginning of star's formation and simulate all the process from the birth onward, or we could select a later phase in the star's evolution, shared from the most, in order to reduce unnecessary computational power and time consumption. If we want to simulate entire stellar populations choosing a starting point also implies selecting appropriate distributions for the main parameters that characterize the "starting point population", such as masses, metallicities, but also orbital parameters for the stars that are in binary systems, such as orbital period, distance, and eccentricity. *Second*, we should choose how the stars will evolve from the starting point, and this involves the single star evolution but also the effects that interaction with other stars in binary systems have on it. *Finally*, we have to decide when we want to stop the simulation, choosing an endpoint that aligns with the needs of this study.

2.1.1 The starting point

As we know, in the Hertzsprung-Russel diagram, shown in [Figure 2.1](#), which plots the *luminosity* of the stars in solar units vs their surface temperature in Kelvin, most of the

stars appear distributed in the **main sequence (MS)**, a continuous and distinctive band. A star's position on this band is determined by its initial mass, and a good rule of thumb is that the most massive stars are hotter, more luminous, and evolve more quickly, while the lower-mass stars burn their fuel more slowly, and remain on the MS longer.




Figure 2.1: Hertzsprung–Russell (H–R) diagram showing the relationship between stellar luminosity (in solar units) and surface temperature (in Kelvin), that indicates the main stellar evolutionary stages: the main sequence, where most stars lie, the giant and supergiant branches, and the white dwarf region. The source of the image is [here](#).

Since almost all stars go through a phase in the MS, and evolve from there differently, in this work, the chosen starting point for stellar evolution is the Zero Age Main Sequence (ZAMS). At this stage, stars have just begun hydrogen burning in their cores, marking the start of their stable main sequence phase. This allows to bypass the early phases of star formation, which are much less relevant to the GE sources of interest, while

still capturing the essential evolutionary processes that lead to the formation of compact objects.

2.1.2 Single star evolution

Simulating the evolution of a single star is in itself a very complex matter, and the only way to make it computationally feasible in the context of large-scale population synthesis is to approximate the evolution for a wide range of mass M and metallicity Z . In fact, detailed evolution codes can require substantial computational time even for the evolution of a single star, which is not practical when generating a full-scale astrophysical population containing millions of stars. Also, in order to make population synthesis statistically robust a large enough number of stars of a certain type should be evolved in order to overcome stochastic noise (in particular, the Poisson noise for n simulations of a particular type of star, implies an error that grows as $\sqrt{n}$). A winning strategy, adopted by several population synthesis frameworks, is to pre-generate a large grid of detailed stellar evolution models, and use them to derive a number of interpolation formulae as functions that approximate stellar properties as a function of age, mass and metallicity. In Hurley et al. [2000] this method is implemented through the development of a set of **Single Star Evolution (SSE)** formulae, with the result of a very compact, efficient and adaptable code, which makes it perfect for the integration of binary-star interactions. The work presented in Hurley et al. [2000] therefore serves as the theoretical and computational foundation for many complex stellar population synthesis codes, including the one used in this thesis. It takes care of the single-star evolution from ZAMS through all the possible evolutionary outcomes, depending on the star's initial conditions.

2.1.3 Binary star evolution

While the evolution of single stars already represents a challenge, the inclusion of binary interactions introduces a much higher level of complexity. In such systems, the evolution of each star is strongly influenced by its companion through a variety of processes, such as mass transfer and accretion, common envelope evolution, collisions, supernova kicks, tidal effects, angular momentum loss, and mergers. These interactions can drastically alter the final outcomes, and are essential for modeling the formation of compact binaries that are potential GW targets for LISA. To efficiently model binary evolution within the framework of stellar population synthesis, the work of Hurley et al. [2002] extends the SSE formalism by introducing a set of prescriptions for binary interactions, and updating the treatment of processes such as Roche lobe overflow, common envelope evolution and coalescence by collision, leading to the development of the **Binary Star Evolution (BSE)** algorithm. This code includes the interpolation-based approach used in SSE for single-star evolution, but adds a comprehensive treatment of binary-specific processes, enabling the simulation of a wide range of binary configurations but keeping the affordable computational requirements of SSE. The BSE algorithm tracks the joint evolution of both stars in a binary system, taking into account their initial parameters, such as masses, orbital period, eccentricity, and metallicity, and updates these properties dynamically as the system evolves. The flexibility and speed of the BSE code make it a key component in many modern population synthesis tools, including the one used in this thesis, which we will now introduce.

2.2 The COSMIC Code

For the purposes of this work, we employ a community-developed binary population synthesis (BPS) python-based code, called the **Compact Object Synthesis and Monte Carlo Investigation Code (COSMIC)**, whose «*primary purpose is to generate synthetic populations with an adaptive size based on how the shape of binary parameter distributions change as the number of simulated binaries increases*». COSMIC's binary evolution is built upon BSE, incorporating extensive modifications in order to include updated physical prescriptions¹. It includes all necessary tools to generate a population, from the generation of initial conditions, to scaling the simulated systems to full-size astrophysical populations. The code is presented in Breivik et al. [2020], where it is described in full detail and used, as a proof of concept, to simulate the Galactic population of compact binaries and their associated GW signal. In the following section we will see the main features of the code, and explain what makes it the right choice for this thesis work.

2.2.1 Fixed population

A fundamental concept in COSMIC, which is the key to the code's efficiency, is the idea of *fixed population*. This refers to a relatively small sample of just enough binaries to capture, in a statistically meaningful way, the underlying shape of the parameter distribution functions of the target population, as determined by the user specified Star Formation History (SFH) and evolution model. This is achieved following an iterative process designed to reach a convergence with respect to a defined matching condition, and consists of five key steps:

1. The user selects a binary evolution model and SFH;
2. Based on the SFH and the chosen initial parameter distribution, an initial population is generated;
3. The population evolves for a user specified number of steps, according to the selected evolution model;
4. If it is the first iteration, half of the simulated systems is compared with the total population. In the following steps, the population from the previous one gets compared to the population containing both the current and previous iterations. In any case, the comparison is done in order to check if the matching condition has been achieved;
5. Once the parameter distributions of the population have converged, the corresponding population is called *fixed population*, which represents the statistical features of a binary evolution model.

¹Beyond the various improvements in modeling astrophysical processes in binary evolution, the main strength of COSMIC, when compared with other BSE-based codes, lies in its efficient use of Monte Carlo techniques, which make large-scale binary population simulations highly efficient. From this point onward, whenever we refer to *sampling*, we specifically mean this Monte Carlo approach. It is worth noting, however, that other recent codes exist, such as Stellar EVolution for N-body (SEVN, Spera et al. [2015]), which, instead of relying on the SSE and BSE algorithms, couples the PAdova and TRieste Stellar Evolution Code (PARSEC, Bressan et al. [2012]) stellar evolution tracks with up-to-date prescriptions for supernova explosions.

In practice, the fixed population is the converged, computationally efficient representation of the systems that we want to simulate, embedded in a complete small-scale synthetic galaxy that also contains other stellar components. The output is stored in a data frame, which separates the full galaxy properties from the fixed population ones. The last step required to construct a full size galaxy is to scale the fixed population (by mass or by number of stars) with a re-sampling approach with replacement, allowing to extrapolate a larger final population that preserves the statistical properties encoded in the fixed population.

Initialization

The fixed population is generated from an initial collection of binaries sampled from distribution functions to assign to each binary an initial value of metallicity (Z), primary star mass (m), mass ratio (q), orbital separation (a), eccentricity (e), and birth time (T_0) according to the selected SFH. In COSMIC the user can choose between different binary parameter distributions, and different parameters can be treated independently. Moreover, COSMIC allows a complete personalization of the initial population through a number of other parameters, including different time-steps to control the binary physics, metallicity, stellar winds, common envelope phase, natal kicks, remnant mass, remnant spin, GW orbital decay, mass transfer, tides, and particular specifications for different kinds of stellar objects, mixing variables, and magnetic braking. In this work all the parameters were left default, but one: we tweaked the metallicity value, in order to differentiate fixed populations describing the parameter distributions for galaxies of different types. We will go more into detail on this topic in the next chapters.

Convergence

The number of simulated systems in the fixed population ideally describes the final parameter distribution functions while being low enough to keep the code efficient. Since every population depends on a different binary evolution model, to quantify this number a *discrete match criteria* is developed, based on the work Chatziioannou et al. [2017]. Independently generated histograms for each parameter are used to track their distribution as successive populations are generated and cumulatively added to the fixed population. The physical limits of the simulated systems are then enforced by taking the logistic transform, and finally the match is defined as:

$$match = \frac{\sum_{k=1}^N P_{k,i} P_{k,i+1}}{\sqrt{\sum_{k=1}^N (P_{k,i} P_{k,i}) \sum_{k=1}^N (P_{k,i} P_{k,i+1})}},$$

where $P_{k,i}$ is the probability for the k th bin, for the i th iteration. For how it is defined, the match value shifts between 0 and 1, and tends to unity as the parameter distributions converge to a distinct shape.

The output

Since COSMIC uses BSE as its core binary evolution algorithm, the output of COSMIC follows most of the same conventions as BSE. The *kstar values* (e.g. the number that

represent a specific stellar type) and evolution stages are nearly identical to their BSE counterparts, and the exact references can be found in **Table A.1** in the **Appendix**. In order to generate a fixed population, COSMIC can be ran through a one-line command directly on the terminal, specifying a parameter file, the *kstar* values for the primary and secondary star, the maximum number of systems to evolve, every how many systems to check in, in order to track the distributions of the parameters, and how many processors to use. The final output is in an *hdf5* file containing several data frames, that keep track of various important quantities during the evolution: the total number of stars and total mass of the entire population, the number of binaries, the convergence, and so on. The *conv* data frame contains all the information about the final fixed population, and thus is the one that we will use the most: from it we can extract all the parameter distributions of the fixed population, such as the orbital parameters, and the individual star information. The parameter distributions of a fixed population of binary white dwarfs with a default metallicity value set at 0.020 are shown in **Figure 2.2**.




Figure 2.2: Distributions of the parameters of a fixed population with metallicity $Z = 0.020$ composed of He+He, CO+He and CO+CO binary white dwarfs. This includes the mass, radius of primary and secondary stars, the radius ratios between the two, the orbital period, average separation, and evolution time.

2.2.2 Astrophysical population

Once the convergence criteria are achieved, an astrophysical population can be sampled. The number of sources in the astrophysical population $N_{astro,tot}$ can be found by upscaling the size of the fixed population, N_{fixed} , by the ratio of the mass of the astrophysical population, M_{astro} , to the mass of all the stars in the whole small-scale galaxy in which the fixed population is embedded, $M_{fixed,stars}$, as follows:

$$N_{astro} = N_{fixed} \frac{M_{astro,tot}}{M_{fixed,stars}}, \quad (2.1)$$

or by the ratio of the number of stars in the astrophysical population, $N_{astro,tot}$, to the total number of stars in the whole small-scale galaxy in which the fixed population is

embedded, $N_{fixed,tot}$,

$$N_{astro} = N_{fixed} \frac{N_{astro,tot}}{N_{fixed,stars}}.$$

Thus, to create a full-scale astrophysical population we need a *reference population* from which we can extract either the total mass or the total number of stars, to then use to scale up our fixed population. As we will now see, the chosen reference for our purpose is a catalog that reports many key galactic parameters in it, which will allow us to proceed using the method in (2.1).

2.3 The Gravitational Wave Galaxy Catalog

As we have seen, the goal of this work is to simulate the GW foreground produced by compact binaries in the *local universe*, by generating the sources using COSMIC. To replicate the existing, observed galaxies in the vicinity of the Milky Way and simulate their stellar content we rely on the dataset provided in White et al. [2011], the GWGC. This catalog includes a list of 53,255 galaxies within $100Mpc$ from earth, containing information on sky position, distance, blue magnitude, major and minor diameters, position angle, and galaxy type, currently used for follow-up searches of electromagnetic counterparts from GW searches.

2.3.1 What it has vs what we need

In principle, we could generate a separate fixed population for each galaxy in the GWGC and scale it individually. However, this is simply not practical because of the computational power and time it would require², and therefore we should find a strategy to group them in a few, representative, categories. As we will show in this section, many of the information in the GWGC can be used to infer the missing astrophysical quantities we need for population synthesis. Ultimately, we will find that metallicity is the most suitable parameter for grouping galaxies. To get there, we follow a chain of empirical relations, starting from the galaxy morphological type, through a luminosity to mass, and then a mass to metallicity relation. This process will enable us to provide COSMIC with the necessary input, found in a consistent and astrophysically motivated way.

2.3.2 Mass-luminosity relation

As previously discussed, in order to scale a fixed population to the size of a specific galaxy, we need either its total stellar mass or the number of stars. In particular, Faber and Gallagher [1979] presents a luminosity-to-mass relation that depends on the morphological type. This allows to use the blue magnitude and the T-type provided in GWGC to estimate each galaxy's stellar mass.

²A single fixed population can require up to $\sim 4h$ to be generated **on a commercial laptop with 12-core CPU**, and the GWGC contains $\sim 50,000$ galaxies: generating a fixed population per each galaxy would require a computational time of $\sim 20yr$.

Magnitude to luminosity

To compute the stellar mass, the luminosity-to-mass ratio presented in Faber and Gallagher [1979] applies to the absolute blue luminosity of each galaxy. GWGC provides the absolute blue magnitude, thus we have to convert it into blue luminosity using the Sun's blue-band absolute magnitude as a reference (typically $m_{B,\odot} = 5.48$). Starting from the magnitude definition,

$$m_{B,gal} - m_{B,\odot} = -2.5 \log_{10} \left(\frac{L_{B,gal}}{L_{B,\odot}} \right),$$

which, in stellar units, we get to the following conversion:

$$L_{B,gal} = 10^{-0.4(m_{B,gal} - m_{B,\odot})} L_{B,\odot}, \quad (2.2)$$

where $m_{B,gal}$ and $m_{B,\odot} \approx 5.48$ are the absolute blue magnitudes of the specific galaxy and the sun.

Galaxy morphological types

Although the GWGC denotes galaxy morphology using the de Vaucouleurs et al. [1991] T-type scale, the luminosity-to-mass relations used in Faber and Gallagher [1979] are defined in terms of Hubble morphological classes. This classification is crucial because different morphological types exhibit significantly different stellar populations and star formation histories, which affect both luminosity and mass content. For example, elliptical galaxies generally have lower luminosity-to-mass ratios than spirals due to their older stellar populations and lack of ongoing star formation. Fortunately, a direct correspondence exists between T-type values and Hubble types, presented in De Vaucouleurs [1974], and thanks to the results in Faber and Gallagher [1979] we find the following correspondences:

T-Value	Hubble Class	$\frac{M}{L_B}$
-6.00 to -4.01	<i>E</i>	8.5
-4.00 to -2.01	<i>SO</i> ⁻	9.5
-2.00 to -0.99	<i>SO</i> ⁺ - <i>S</i> _a	6.2
1.00 to 3.99	<i>S</i> _{ab} - <i>S</i> _{bc}	6.5
4.00 to 4.99	<i>S</i> _{bc} - <i>S</i> _c	4.7
5.00 to 5.99	<i>S</i> _{cd} - <i>S</i> _d	3.9
6.00 to 10.00	<i>S</i> _{dm} - <i>Irr</i>	8.5

Table 2.1: The first two columns of this table present a conversion between the Hubble class and the T-value classifications for the different galactic morphological types. This was done using the conversion table presented in De Vaucouleurs [1974]. The last column associates at each galaxy type a mass-to-luminosity ratio, found using the work in Faber and Gallagher [1979].

The luminosity resulting from (2.2), expressed in units of solar blue luminosity $L_{B,\odot}$,

can then be multiplied by the appropriate M/L_B ratio, found in Table 2.1, to yield the total stellar mass for the galaxy

$$M_{gal} = \frac{M}{L_B} L_{B,gal}$$

2.3.3 Mass-metallicity relation

Now that we have estimated the stellar mass of each galaxy, we can also derive the corresponding metallicity. This is made possible by the results of Tremonti et al. [2004], where an empirical relation between stellar mass and gas-phase oxygen abundance (a proxy for galaxy metallicity) is established. The study, based on a sample of over 50,000 star-forming galaxies observed by the Sloan Digital Sky Survey, spanning three orders of magnitude in stellar mass and a factor of ten in metallicity, reveals a tight correlation between the two, which can be written as follows:

$$Z' = -1.492 + 1.847 \times \log(M_*) - 0.08026 \times [\log(M_*)]^2, \quad (2.3)$$

where M_* is the galactic stellar mass, and

$$Z' = 12 + \log(O/H) \quad (2.4)$$

is the metallicity written in terms of the oxygen abundance³.

On metallicity units

Now, in these units the sun's metallicity is $Z'_\odot = 8.69$ (Allende Prieto et al. [2001]), whereas in the units used by COSMIC it is $Z_\odot = 0.0134$ (Asplund et al. [2009])⁴. We can easily find a conversion between the two unit systems by assuming that the rate of oxygen abundance in the galaxy and the one in the sun is equal to the rate of the metallicities in COSMIC's units,

$$\frac{(O/H)_{gal}}{(O/H)_\odot} = \frac{Z_{gal}}{Z_\odot}, \quad (2.5)$$

where (O/H) can be found inverting (2.4). At this point, knowing that $Z'_\odot = 8.69$, we can:

- Compute the Z'_{gal} value for each galaxy in GWGC using (2.3);
- Write $(O/H)_\odot$ and $(O/H)_{gal}$ in terms of the corresponding Z' value using (2.4): $(O/H) = 10^{(Z'-12)}$ and $(O/H)_\odot \approx 0.32$.

Finally, finding Z_{gal} from (2.5), we can write the metallicity units conversion as:

$$Z_{gal} = 27.3 \times 10^{(Z'-12)}, \quad (2.6)$$

where the "27.3" coefficient mainly depends on the $Z_\odot$ value. This relation allows us to assign a realistic metallicity estimate to each GWGC galaxy in COSMIC's units, which we can now use to group them into representative subsets. This way, we can account for the significant impact of metallicity on stellar wind strength, remnant masses, and binary evolution outcomes, that greatly affects the expected GW signal.

³All the metallicity values with an apostrophe are intended as in these units.

⁴All the metallicity values without the apostrophe are intended as in these units.

Chapter 3

Results and Discussion

In the previous chapters, we have laid the theoretical and methodological foundations of our investigation: from the GWs theory, to the astrophysical context of WDs and their host galaxies, the principles of GW detection, and finally the construction of the simulated populations used in this work. In this chapter, we turn to the quantitative generation of the corresponding GW signal. We begin by completing the list of galaxies that we use as a reference for the simulations, by taking into account for the incomplete entries of the GWGC and the galaxies obscured by the ZOA. Afterwards, we generate the fixed populations that are used to simulate all the galaxies, by choosing the corresponding metallicity values based on the range covered by the galaxies in the final catalog. After discussing LISA's frequency resolution, and understanding how to take it into account, we are able to apply the computational pipeline described in Chapter 3 to our final galaxy catalog, extracting the GW signal from the simulated populations of binary white dwarfs. The resulting spectra are then compared to the LISA sensitivity curve to verify the detectability of this source of an extragalactic foreground. Finally, we discuss the frequency distribution and amplitude characteristics of the simulated signals, as well as the physical reasons underlying our results, and their implications for future space-based GW detectors.

3.1 Preliminary steps

Before computing the total extragalactic DWDs gravitational foreground, we **should** ensure that our list of galaxies is truly representative of the total population of DWDs in the nearby universe. This means we **should** consider that the final catalog that we use to compute the GW signal is incomplete, analysing all the causes of lack of galaxies. Finally, we group the final galaxies in terms of metallicity, and generate the fixed populations that we will use to reproduce the DWDs hosted in them.

3.1.1 Missing galaxies

The GWGC is far from being a complete list of the galaxies in the local Universe, and there are two main factors contributing to this: observational limitations due to the ZOA and limitations of the information that the catalog presents.

GWGC incompleteness and non-galactic objects

Not all the objects listed in the GWGC are suitable for our purposes: in addition to galaxies, the catalog also includes globular clusters, which can be identified through their T-type values; moreover, not all galaxies in the catalog contain all the information required to perform the procedures described in the previous sections. To construct a working sample, we applied a series of masks to clean the catalog:

- Only keep entries classified as galaxies;
- Require that the distance, T-type, and absolute blue magnitude are provided;

After this selection, the sample we are left with scaled down in size, from the original 53,255 entries to 20,246 galaxies suitable for population synthesis.

Zone Of Avoidance

Since the GWGC is an observation-based catalog derived from optical surveys, we **should** account for the ZOA, the region of the sky obscured by the Milky Way’s interstellar dust and stellar crowding, which impedes the detection of foreground galaxies. The extent of the ZOA depends on the wavelength of observation: infrared surveys penetrate deeper through the dust, while optical surveys, such as those used for GWGC (the reported magnitude for these galaxies is in the B-band), are more strongly affected. In the optical band, the ZOA is expected to cover about $\sim 25\%$ of the sky, as reported in Kraan-Korteweg and Lahav [2000]. For this work, we estimated the ZOA directly from the spatial distribution of galaxies in GWGC, by plotting their positions in Galactic coordinates, and highlighting the empty regions of the sky. This way, the ZOA appears clearly as a horizontal band centered around the Galactic plane (see **Figure 3.1**). To quantify its extent, we used a Python package that handles pixelated data on the sphere based on the Hierarchical Equal Area isoLatitude Pixelization, **HEALPix** (Gorski et al. [2005]), called **healpy**, to divide the sky map into n_{pix} equal-area pixels, and focused on the band $|b| \leq 15^\circ$ in Galactic latitude. Counting the empty pixels in this band and dividing by n_{pix} gives an estimated ZOA coverage of $\sim 20\%$ – 25% of the sky, depending on the chosen pixel resolution. This estimate is fully consistent with the literature Kraan-Korteweg and Lahav [2000], so in our analysis we will assume that GWGC effectively covers $\sim 80\%$ of the sky.

Filling the gap

In order to compensate for the missing galaxies, we have made two physically motivated assumptions.

1. Cosmological Principle:

The GWGC lists all the observed galaxies within 100Mpc. On such scales, the Universe can be assumed to be statistically *homogeneous* and *isotropic*. This assumption is supported from observations of the *Cosmic Microwave foreground* (CMB), where we know that the correlation function of the temperature fluctuations across the sky peaks at scales of $\sim 100\text{Mpc}$. Under this assumption, the selected galaxies of the GWGC **should** be statistically representative of the overall population in the same volume, and therefore of the missing galaxies too.




Figure 3.1: Visual representation of the regions of the sky not covered by any of the 53,255 galaxies in the GWGC, plotted in galactic coordinates using `healpy`, a Python package that handles pixelated data on the sphere based on the Hierarchical Equal Area isoLatitude Pixelization (**HEALPix**). This library allows the user to choose the number of pixels, and thus their size, through a power-of-2 parameter in the `nside2npix` function: the **larger** the parameter, the smaller the pixel and the **larger** their number. The gray dots in the figure represent the empty sky pixels, while the white areas correspond to pixels containing at least one galaxy. The central band where the Milky Way lies, appears significantly emptier than the rest of the sky. To estimate this ZOA, we computed the percentage of empty pixels in the band $|b| \leq 15^\circ$ in Galactic latitude. Choosing the parameter to be too small, the produced pixels would be too **large**, and the empty area would be underestimated. **larger** values would make the estimation only more precise, and by selecting parameters starting from $2^6 = 64$, it's possible to estimate roughly $\sim 20\%$ – 25% of empty sky, fully consistent with the value reported in Kraan-Korteweg and Lahav [2000]. The 22.07% estimation in the title is found using 64 as parameter value.

2. Distance:

The angular position is irrelevant in the computation of the GW signal: only the distance of the source matters. Therefore, missing galaxies from the unexplored volume covered by the Milky Way can be statistically represented by the galaxies from our retained sample.

Following these assumptions, we “fill in” the missing fraction of the sky and the dropped entries of the catalog by sampling with replacement from the cleaned GWGC, just like we did to extract an astrophysical population from the fixed one, preserving its parameter distributions. This ensures that the statistical properties of the synthetic galaxy distribution match those of the observed portion while restoring the full-sky coverage. This is shown in a graphical representation of the distributions of distances, blue band magnitudes and T-types, in **Figure 3.2**: the distributions have been scaled up without visible changes.




Figure 3.2: Distributions of T-type, distance and B-band magnitude of the galaxies in the GWGC before and after compensating for the lack of galaxies due to the catalog’s incompleteness and the ZOA. The three graphs *above* show the mentioned distributions for the 20,246 galaxies of the original GWGC after we dropped those missing the key parameters we needed; the three graphs *below* show the same distributions for the 66,568 galaxies we are left with after taking into account both the catalog’s incompleteness and the ZOA using a sampling with replacement method. The parameters distributions appear to be indistinguishable in shape: the scaling up in size left the properties of the catalog unchanged.

The total number of galaxies that we have after taking into consideration both the dropped incomplete entries of the original GWGC and the galaxies covered by the ZOA, has grown to 66,568.

3.1.2 The final fixed populations

As we already discussed, the metallicity of a galaxy has a significant impact on the stellar evolution and, thus, on the GW signal. This, plus the fact that metallicity is also closely linked to the galaxy types, led us to choose it as characterizing parameter for the fixed populations that we will generate to represent the galaxy in the catalog. In particular, we the final catalog covers a range from a minimum of $Z_{min} = 0.00748$ to a maximum of $Z_{max} = 0.03948$. We decided to group all the values in between in ten equally wide bins, represented in **Figure 3.3**.




Figure 3.3: Metallicity distribution of the final galaxy catalog. The metallicity of the galaxies has been shown to correlate with their morphological type in Faber and Gallagher [1979]. Therefore, the entries of our final catalog can be grouped into 10 bins, each associated to a metallicity value (the middle value of the corresponding bin, shown in the x -axis label of the graph), and a corresponding fixed population.

Each one of the bins center values, shown in the x label, is linked to a different synthetic fixed population generated with COSMIC. The simulations were done generating the fixed populations "*the cosmic way*", via terminal, specifying $Nstep = 5000$ and $Niter = 1000000000$ and $kstar$ values 10, 11 and 12, corresponding to He, CO, and Ne white dwarfs¹, assuming the sun's metallicity to be $Z_{\odot} = 0.0134$ and leaving all the remaining COSMIC's specifiable parameters unchanged from the default values.

¹All the $kstar$ values and the corresponding astrophysical objects are shown in **Table A.1**, in the **Appendix**.

3.2 Th Signal Computation

Generating the DWDs using COSMIC gives us the possibility to compute the GW contribution of each one of them, assuming their distance to be the same as the hosting galaxy's one, provided from the galaxy catalog, and with an extreme precision in frequency. Before scaling up the fixed population, and computing the GW signal of each binary, and confronting it with LISA's sensitivity curve, we **should** then consider LISA's limited *frequency resolution*, and understand how to process the signals from indistinguishable sources.

3.2.1 LISA's frequency resolution

The duration of LISA's mission directly impacts its frequency resolution: the longer the observation time, the finer the binning in frequency². In this work, we assume an observation time of:

$$T_{obs} = 4yr = 126,144,000s$$

which corresponds to a minimum frequency resolution of

$$\delta\nu_{LISA,min} \approx \frac{1}{T_{obs}} \sim 8 \times 10^{-9} Hz. \quad (3.1)$$

This fundamental limit comes from the Fourier transform properties, and determines the smallest frequency interval over which the signals can be resolved. **Even with such fine resolution, the orbital period of the simulated binaries leads to a frequency precision so high that LISA cannot distinguish sources within the same frequency bin.** To model this, we group the resulting GW signals in bins of size $\delta\nu_{LISA,min}$, and sum all the signals that fall within each bin.

Signal accumulation

In principle, over the observation time T_{obs} , the orbital parameters of the sources may slowly evolve, causing a drift in the GW frequency. Integrating the resulting signal during this period is no easy task. However, for the slowly evolving white dwarf binaries we are considering, the change in such short time-frames is negligible, so we can treat their signals as effectively monochromatic within the observation window. Thus, we can define a commonly used variable to simplify our job, which is called **Amplitude Spectral Density (ASD)**:

$$ASD = h\sqrt{T_{obs}}, \quad (3.2)$$

where h is the GW strain of the binary, computed in (1.22), and T_{obs} is the mission duration for LISA. This quantity represents the characteristic strain amplitude of the GW signal per square root of frequency, scaled by the observation time, and its very convenient for comparing the signals in frequency space. Correspondingly, we can define the **Power Spectral Density (PSD)** as the square of the ASD

$$PSD = ASD^2, \quad (3.3)$$

²Not to be confused with the sensitivity of the instrument, which tells how faint of a GW strain it is able to detect.

which properly represents the signal power per frequency unit. Since power contributions from independent sources add linearly, the total PSD in a given frequency bin is simply found by summing the power of all the sources within it:

$$PSD_{tot,bin_1} = PSD_{*1,bin_1} + PSD_{*2,bin_1} + \dots \quad (3.4)$$

This way, we effectively accounted for LISA's frequency resolution limit. The reason for introducing this quantities is that GW from different binaries are uncorrelated, so their strains add incoherently. This means that we cannot simply sum the strain amplitudes directly. Instead, the PSD provides a way to combine signals that ensures a correct representation of the combined GW foreground.

3.2.2 The final output

In order to compute the total extragalactic DWDs gravitational foreground, we **should** apply an iterative process to each galaxy in the final catalog, which consists of the following steps:

- Compute the right N_{astro} using the (2.1)
- Scale the fixed population accordingly with a sampling with replacement method, creating the full-size galaxy simulation.

At this point for each binary system in this astrophysical population we:

- Compute the GW signal using (1.22)³ and the corresponding PSD using (3.2) and (3.3);
- Sum the total PSD of the binaries whose frequencies fall within the same LISA's frequency bin, using (3.4);
- Plot the total binned signals on LISA's curve.

LISA's sensitivity curve

The LISA curve is constantly updated when new GW components that can affect it are found, meaning that there are many different versions around. The one we chose to adopt is the widely used version presented in Robson et al. [2019], which is written as:

$$S_n = \frac{10}{3L^2} \left(P_{OMS}(f) + \frac{4P_{acc}(f)}{(2\pi f)^4} \right) \left(1 + \frac{6}{10} \left(\frac{f}{f_*} \right)^2 \right) + S_c(f),$$

where $L = 2.5Gm$ is the LISA arm length, $f_* = 19.09mHz$ and $P_{OMS}(f)$, $P_{acc}(f)$ and $S_c(f)$ are the expressions for the single-link optical metrology noise, the single test mass acceleration noise and the total Michaelson-style LISA data channel. The equations for these factors, as well as more detail on the curve, can be found in the article.

³Note here that all the binaries within the same galaxy are given the same distance from us. This approximation is considered to be valid since the distance of the galaxy is much much **larger** than its size, and thus then the maximum possible distance between two binaries within the same galaxy.

3.3 Results and Discussion

The main result of this work is the spectral distribution of the final computed signal plotted over the LISA's sensitivity curve (Robson et al. [2019]) as blue dots in Figure 3.4. Every dot in this graph represents a frequency bin, whose width is given by (3.1), where




Figure 3.4: The blue dots represent the total GW signal of the simulated DWDs hosted by the 66,568 galaxies of the estimated potential gravitational wave emitting galaxies, whereas the orange dots represent the GW signal considering only the 20,246 of the GWGC after dropping the galaxies with incomplete information, and without considering the ZOA. In Both cases, the PSDs are binned in $\sim 8 \times 10^{-9} \text{ Hz}$ large frequency bins, representing LISA's spectral resolution for an observational time $T_{obs} = 4 \text{ yr}$. Every dot represents the GW signal of a frequency bin, computed by summing the Power Spectral Density (PSD) of every galaxy that it contained, using (3.4). The dots, plotted on LISA's sensitivity curve, as found in Robson et al. [2019], are about one order of magnitude below the minimum threshold of this curve and a slight difference in height is visible between the blue and the orange dot's distributions, showing the effect of considering a significantly different number of sources. This results suggest that we **should** not expect LISA to be able to detect the unresolved extragalactic DWDs confusion foreground, at least under the assumptions and approximations adopted in this work, and suggests that the results do not depend significantly from the number of galaxies.

the GW signal of all the binaries within it has been summed using (3.4). The resulting computed signal is about one order of magnitude smaller than the minimum detectable strain for LISA across the whole band. This indicates that the stochastic contribution

from unresolved extragalactic DWDs will not have a measurable impact on LISA's sensitivity curve, at least under the assumptions of our population model and scaling relations.

Distribution of the sources The simulated extragalactic binaries foreground populates primarily the low-frequency range ($f \lesssim 2$ mHz), with the number of systems per frequency bin increasing steeply at higher frequencies. This is evident for at least two reasons:

1. The dots are more sparse at lower frequencies, and increasingly densely packed towards higher frequencies, showing that more empty bins appear to be at the lower end of the spectrum, and the more populated ones at the higher end;
2. The sources lay on a slope, with the PSD values steadily increasing with the frequency, showing that at higher frequencies, the occupied frequency bins tend to be more and more populated, and therefore the total PSD they contain higher and higher in magnitude.

A possible physical explanation for this phenomenon is that the time evolution of the orbital parameters is initially very slow, allowing the orbital frequencies to distribute over a wider range of bins. As the binary systems evolve they get more compact, and the parameters change more rapidly, increasing the likelihood that the systems occupy very close frequency bins. Moreover, the spectral shape of the resulting foreground is located at frequencies similar to the expected Galactic foreground ones, corresponding to the slight bump at the left of the lowest point in the graph (this is better shown in Figure 1.3), but with a significantly lower amplitude due to the much larger average distances. Indeed, as shown in (1.22), the GW amplitude scales inversely with the distance from the source, so even relatively nearby galaxies contribute far less per binary than the Milky Way population. The total number of galaxies in the local Universe, even if large, is not sufficient to compensate for this distance-induced amplitude suppression.

3.3.1 Robustness of the results

The simulated GW signal, represented by the blue dots, would be detectable by LISA if it fell above the LISA curve, meaning that the total PSD in a certain frequency bin is higher than the minimum threshold of LISA's detectability. Therefore, the dots can get close to, or above the sensitivity curve either by containing extremely close and strong sources, by containing the contribution of many farther, and fainter sources, or a combination of both. In principle, the results we presented could depend on the assumptions that we made, and the methods that we used.

The sources

The results could be affected by the way we represented the sources, via GWGC. Indeed, this is the input we give our pipeline to produce, as an output, what should be the total GW confusion foreground from all the unresolved DWDs in the local universe. For example, if we only used the GWGC catalog after dropping the incomplete voices, and without considering the ZOA, the total GW signal produced by the remaining 20,246 galaxies would be as represented in the orange dots in Figure 3.4. Although a slight

difference in height with the results obtained for all the 66,568 galaxies, presented as blue dots in the same figure, is noticeable, it is important to note that it is caused by considering less than a third of the sources. Therefore, even if it is possible to consider different and more complete catalogs, and the ZOA could be estimated with better precision, the results we found do not depend significantly on this. In addition to this, the single DWDs were simulated using COSMIC, which represent an arbitrary choice between a number of different population synthesis codes: a different code might produce different sources parameter distributions, and could be an interesting extension of the present work.

The GW signal

The main assumptions that we made to obtain the results are surely how we computed the GW characteristic strain, and the way we integrate the GW signal coming from unresolved sources within the same frequency bin. The GW signal computed in (1.22) is the standard characteristic strain predicted from the General Relativity for an inspiraling binary system, and is therefore the most precise way to represent the DWDs signal. Furthermore, the use of ASD and PSD is unavoidable, since it is the only way to confront signals that LISA could not resolve in frequency with its sensitivity curve. Interestingly, assuming a longer observational time would guarantee LISA a better frequency resolution, and each frequency bin would appear to be less populated. Since, as we said, less populated bins generally contain lower total PSD, the better the LISA's frequency resolution, the lower the sensitivity to the noise that the unresolved sources produce, and therefore the lower LISA's sensitivity curve needs to be to detect it. Overall, the methods we used guarantee a pretty stable and reproducible output, given the same input.

Conclusions and Future Perspectives

In this thesis we investigated the contribution of extragalactic DWDs within the nearby universe ($\sim 100\text{Mpc}$ from us) to the low-frequency GW foreground, with a focus on their potential detectability by the space-based interferometer LISA. We began by discussing the theoretical GW framework, and the astrophysical elements that are relevant to this problem, including a brief description of the formation and evolution of white dwarfs, both as isolated stars and in binary systems, and the galactic environments in which they reside and evolve. We discussed the fundamentals of interferometric detection, and the specific capabilities of LISA as a space-based detector. On this theoretical basis, we developed a computational pipeline to simulate synthetic populations of extragalactic DWDs using the population synthesis code COSMIC. We then scaled them according to the stellar masses of host galaxies, using the astrophysical observations galaxy survey GWGC as reference, and computed their GW strain and power spectral density. The simulated signals were finally binned according to LISA's frequency resolution and compared with its sensitivity curve.

The main original result of this study is that the predicted GW signal from the extragalactic DWDs population lies approximately one order of magnitude below LISA's detection threshold⁴ across the considered frequency band. Therefore, although, as we know, DWDs are expected to constitute a dominant foreground from Galactic sources, able to shape LISA's curve substantially, we do not expect the extragalactic component to measurably affect LISA's future observations at all.

While the present analysis points towards a negligible extragalactic contribution for LISA, it provides a quantitative baseline that can be refined in some directions. We questioned the robustness of our results and found that significant variations in the most relevant steps leading to the output are either inapplicable, as in the computation of the strain within General Relativity, or produce changes in the output so small that they can be considered negligible, as shown considering the output of a much smaller number of sources.

Future extensions of this work could incorporate more detailed galaxy catalogs with redshift-dependent star formation histories and metallicity evolution, and that take in account more precisely the missing sources blocked from our Galaxy in the ZOA. A less predictable, and possibly more relevant improvement, could regard the choice of binary population synthesis models: for example, by including different prescriptions also on other parameters, and not only focusing on the metallicity, such as mass transfer prescriptions or alternative common-envelope outcomes. Moreover, since population

⁴For the LISA curve presented in Robson et al. [2019]

synthesis is a very active and rapidly evolving research field, testing alternative codes, possibly based on different Single Star and Binary Star evolution models, could help quantify the model dependence of our predictions and reveal systematic uncertainties.

Finally, although the extragalactic DWD foreground may be undetectable for LISA, this work shows how future space-based detectors with longer baselines or improved sensitivity in the sub-millihertz regime could, in principle, probe this population, offering a complementary view to Galactic studies and providing constraints on the binary evolution across cosmic time. In summary, this work has explored the extragalactic DWDs component of the GW landscape, reinforcing the understanding that the GW sky at low frequencies will be dominated by Galactic sources, but also highlighting that the extragalactic foreground, though faint, remains a scientifically interesting target for the next generation of detectors.

Appendix A

Table A.1: In this table are presented all the possible kstar values that the user can choose in COSMIC, and the corresponding astrophysical object and evolutionary state.

kstar value	evolutionary state
0	Main Sequence (MS), $< 0.7M_{\odot}$
1	MS, $> 0.7M_{\odot}$
2	Hertzsprung Gap
3	First Giant Branch
4	Core Helium Burning
5	Early Asymptotic Giant Branch (AGB)
6	Thermally Pulsing AGB
7	Naked Helium Star MS
8	Naked Helium Star Hertzsprung Gap
9	Naked Helium Star Giant Branch
10	Helium White Dwarf
11	Carbon/Oxygen White Dwarf
12	Oxygen/Neon White Dwarf
13	Neutron Star
14	Black Hole
15	Massless Remnant

Bibliography

- J Aasi, B P Abbott, R Abbott, T Abbott, M R Abernathy, K Ackley, C Adams, T Adams, P Adesso, R X Adhikari, V Adya, C Affeldt, N Aggarwal, O D Aguiar, A Ain, P Ajith, A Alesic, B Allen, D Amariutei, S B Anderson, W G Anderson, K Arai, M C Araya, C Arceneaux, J S Areeda, G Ashton, S Ast, S M Aston, P Aufmuth, C Aulbert, B E Aylott, S Babak, P T Baker, S W Ballmer, J C Barayoga, M Barbet, S Barclay, B C Barish, D Barker, B Barr, L Barsotti, J Bartlett, M A Barton, I Bartos, R Bassiri, J C Batch, C Baune, B Behnke, A S Bell, C Bell, M Benacquista, J Bergman, G Bergmann, C P L Berry, J Betzwieser, S Bhagwat, R Bhandare, I A Bilenko, G Billingsley, J Birch, S Biscans, C Biwer, J K Blackburn, L Blackburn, C D Blair, D Blair, O Bock, T P Bodiya, P Bojtos, C Bond, R Bork, M Born, Sukanta Bose, P R Brady, V B Braginsky, J E Brau, D O Bridges, M Brinkmann, A F Brooks, D A Brown, D D Brown, N M Brown, S Buchman, A Buikema, A Buonanno, L Cadonati, J Calderón Bustillo, J B Camp, K C Cannon, J Cao, C D Capano, S Caride, S Caudill, M Cavaglià, C Cepeda, R Chakraborty, T Chalermongsak, S J Chamberlin, S Chao, P Charlton, Y Chen, H S Cho, M Cho, J H Chow, N Christensen, Q Chu, S Chung, G Ciani, F Clara, J A Clark, C Collette, L Cominsky, M Constanicio, D Cook, T R Corbitt, N Cornish, A Corsi, C A Costa, M W Coughlin, S Countryman, P Couvares, D M Coward, M J Cowart, D C Coyne, R Coyne, K Craig, J D E Creighton, T D Creighton, J Cripe, S G Crowder, A Cumming, L Cunningham, C Cutler, K Dahl, T Dal Canton, M Damjanic, S L Danilishin, K Danzmann, L Darte, I Dave, H Daveloza, G S Davies, E J Daw, D DeBra, W Del Pozzo, T Denker, T Dent, V Dergachev, R T DeRosa, R DeSalvo, S Dhurandhar, M D'iaz, I Di Palma, G Dojcinoski, E Dominguez, F Donovan, K L Dooley, S Doravari, R Douglas, T P Downes, J C Driggers, Z Du, S Dwyer, T Eberle, T Edo, M Edwards, M Edwards, A Effler, H.-B Eggenstein, P Ehrens, J Eichholz, S S Eichenberry, R Essick, T Etzel, M Evans, T Evans, M Factourovich, S Fairhurst, X Fan, Q Fang, B Farr, W M Farr, M Favata, M Fays, H Fehrmann, M M Fejer, D Feldbaum, E C Ferreira, R P Fisher, Z Frei, A Freise, R Frey, T T Fricke, P Fritschel, V V Frolov, S Fuentes-Tapia, P Fulda, M Fyffe, J R Gair, S Gaonkar, N Gehrels, L Á Gergely, J A Giaime, K D Giardina, J Gleason, E Goetz, R Goetz, L Gondan, G González, N Gordon, M L Gorodetsky, S Gossan, S Goßler, C Gräf, P B Graff, A Grant, S Gras, C Gray, R J S Greenhalgh, A M Gretarsson, H Grote, S Grunewald, C J Guido, X Guo, K Gushwa, E K Gustafson, R Gustafson, J Hacker, E D Hall, G Hammond, M Hanke, J Hanks, C Hanna, M D Hannam, J Hanson, T Hardwick, G M Harry, I W Harry, M Hart, M T Hartman, C-J Haster, K Haughian, S Hee, M Heintze, G Heinzl, M Hendry, I S Heng, A W Heptonstall, M Heurs, M Hewitson, S Hild, D Hoak, K A Hodge, S E Hollitt, K Holt, P Hopkins, D J Hosken,

- J Hough, E Houston, E J Howell, Y M Hu, E Huerta, B Hughey, S Husa, S H Hutner, M Huynh, T Huynh-Dinh, A Idrisy, N Indik, D R Ingram, R Inta, G Islas, J C Isler, T Isogai, B R Iyer, K Izumi, M Jacobson, H Jang, S Jawahar, Y Ji, F Jiménez-Forteza, W W Johnson, D I Jones, R Jones, L Ju, K Haris, V Kalogera, S Kandhasamy, G Kang, J B Kanner, E Katsavounidis, W Katzman, H Kaufer, S Kaufer, T Kaur, K Kawabe, F Kawazoe, G M Keiser, D Keitel, D B Kelley, W Kells, D G Keppel, J S Key, A Khalaidovski, F Y Khalili, E A Khazanov, C Kim, K Kim, N G Kim, N Kim, Y.-M Kim, E J King, P J King, D L Kinzel, J S Kissel, S Klimenko, J Kline, S Koehlenbeck, K Kokeyama, V Kondrashov, M Korobko, W Z Korth, D B Kozak, V Kringel, B Krishnan, C Krueger, G Kuehn, A Kumar, P Kumar, L Kuo, M Landry, B Lantz, S Larson, P D Lasky, A Lazzarini, C Lazzaro, J Le, P Leaci, S Leavey, E O Lebigot, C H Lee, H K Lee, H M Lee, J R Leong, Y Levin, B Levine, J Lewis, T G F Li, K Libbrecht, A Libson, A C Lin, T B Littenberg, N A Lockerbie, V Lockett, J Logue, A L Lombardi, M Lormand, J Lough, M J Lubinski, H Lück, A P Lundgren, R Lynch, Y Ma, J Macarthur, T MacDonald, B Machenschalk, M MacInnis, D M Macleod, F Magaña-Sandoval, R Magee, M Mageswaran, C Maglione, K Mailand, I Mandel, V Mandic, V Mangano, G L Mansell, S Márka, Z Márka, A Markosyan, E Maros, I W Martin, R M Martin, D Martynov, J N Marx, K Mason, T J Massinger, F Matichard, L Matone, N Mavalvala, N Mazumder, G Mazzolo, R McCarthy, D E McClelland, S McCormick, S C McGuire, G McIntyre, J McIver, K McLin, S McWilliams, G D Meadors, M Meinders, A Melatos, G Mendell, R A Mercer, S Meshkov, C Messenger, P M Meyers, H Miao, H Middleton, E E Mikhailov, A Miller, J Miller, M Millhouse, J Ming, S Mirshekari, C Mishra, S Mitra, V P Mitrofanov, G Mitselmakher, R Mitelman, B Moe, S D Mohanty, S R P Mohapatra, B Moore, D Moraru, G Moreno, S R Morriss, K Mossavi, C M Mow-Lowry, C L Mueller, G Mueller, S Mukherjee, A Mullavey, J Munch, D Murphy, P G Murray, A Mytidis, T Nash, R K Nayak, V Nacula, K Nedkova, G Newton, T Nguyen, A B Nielsen, S Nissanke, A H Nitz, D Nolting, M E N Normandin, L K Nuttall, E Ochsner, J O'Dell, E Oelker, G H Ogin, J J Oh, S H Oh, F Ohme, P Oppermann, R Oram, B O'Reilly, W Ortega, R O'Shaughnessy, C Osthelder, C D Ott, D J Ottaway, R S Ottens, H Overmier, B J Owen, C Padilla, A Pai, S Pai, O Palashov, A Pal-Singh, H Pan, C Pankow, F Pannarale, B C Pant, M A Papa, H Paris, Z Patrick, M Pedraza, L Pekowsky, A Pele, S Penn, A Perreca, M Phelps, V Pierro, I M Pinto, M Pitkin, J Poeld, A Post, A Poteomkin, J Powell, J Prasad, V Predoi, S Premachandra, T Prestegard, L R Price, M Principe, S Privitera, R Prix, L Prokhorov, O Puncken, M Pürner, J Qin, V Quetschke, E Quintero, G Quiroga, R Quitzow-James, F J Raab, D S Rabeling, H Radkins, P Raffai, S Raja, G Rajalakshmi, M Rakhmanov, K Ramirez, V Raymond, C M Reed, S Reid, D H Reitze, O Reula, K Riles, N A Robertson, R Robie, J G Rollins, V Roma, J D Romano, G Romanov, J H Romie, S Rowan, A Rüdigger, K Ryan, S Sachdev, T Sadecki, L Sadeghian, M Saleem, F Salemi, L Sammut, V Sandberg, J R Sanders, V Sannibale, I Santiago-Prieto, B S Sathyaprakash, P R Saulson, R Savage, A Sawadsky, J Scheuer, R Schilling, P Schmidt, R Schnabel, R M S Schofield, E Schreiber, D Schuette, B F Schutz, J Scott, S M Scott, D Sellers, A S Sengupta, A Sergeev, G Serna, A Seigny, D A Shaddock, M S Shahriar, M Shaltev, Z Shao, B Shapiro, P Shawhan, D H Shoemaker, T L Sidery, X Siemens, D Sigg, A D Silva, D Simakov, A Singer, L Singer, R Singh, A M Sintès, B J J Slagmolen, J R Smith,

- M R Smith, R J E Smith, N D Smith-Lefebvre, E J Son, B Sorazu, T Souradeep, A Staley, J Stebbins, M Steinke, J Steinlechner, S Steinlechner, D Steinmeyer, B C Stephens, S Steplewski, S Stevenson, R Stone, K A Strain, S Strigin, R Sturani, A L Stuver, T Z Summerscales, P J Sutton, M Szczepanczyk, G Szeifert, D Talukder, D B Tanner, M Tápai, S P Tarabrin, A Taracchini, R Taylor, G Tellez, T Theeg, M P Thirugnanasambandam, M Thomas, P Thomas, K A Thorne, K S Thorne, E Thrane, V Tiwari, C Tomlinson, C V Torres, C I Torrie, G Traylor, M Tse, D Tshilumba, D Ugolini, C S Unnikrishnan, A L Urban, S A Usman, H Vahlbruch, G Vajente, G Valdes, M Vallisneri, A A van Veggel, S Vass, R Vaulin, A Vecchio, J Veitch, P J Veitch, K Venkateswara, R Vincent-Finley, S Vitale, T Vo, C Vorvick, W D Vousden, S P Vyatchanin, A R Wade, L Wade, M Wade, M Walker, L Wallace, S Walsh, H Wang, M Wang, X Wang, R L Ward, J Warner, M Was, B Weaver, M Weinert, A J Weinstein, R Weiss, T Welborn, L Wen, P Wessels, T Westphal, K Wette, J T Whelan, S E Whitcomb, D J White, B F Whiting, C Wilkinson, L Williams, R Williams, A R Williamson, J L Willis, B Willke, M Wimmer, W Winkler, C C Wipf, H Wittel, G Woan, J Worden, S Xie, J Yablon, I Yakushin, W Yam, H Yamamoto, C C Yancey, Q Yang, M Zanolin, Fan Zhang, L Zhang, M Zhang, Y Zhang, C Zhao, M Zhou, X J Zhu, M E Zucker, S Zuraw, and J Zweizig. Advanced ligo. *Classical and Quantum Gravity*, 32(7):074001, mar 2015. doi: 10.1088/0264-9381/32/7/074001. URL <https://dx.doi.org/10.1088/0264-9381/32/7/074001>.
- B. P. Abbott, R. Abbott, T. D. Abbott, M. R. Abernathy, F. Acernese, K. Ackley, C. Adams, T. Adams, P. Addesso, R. X. Adhikari, V. B. Adya, C. Affeldt, M. Agathos, K. Agatsuma, N. Aggarwal, O. D. Aguiar, L. Aiello, A. Ain, P. Ajith, B. Allen, A. Allocca, P. A. Altin, S. B. Anderson, W. G. Anderson, K. Arai, M. A. Arain, M. C. Araya, C. C. Arceneaux, J. S. Areeda, N. Arnaud, K. G. Arun, S. Ascenzi, G. Ashton, M. Ast, S. M. Aston, P. Astone, P. Aufmuth, C. Aulbert, S. Babak, P. Bacon, M. K. M. Bader, P. T. Baker, F. Baldaccini, G. Ballardín, S. W. Ballmer, J. C. Barayoga, S. E. Barclay, B. C. Barish, D. Barker, F. Barone, B. Barr, L. Barsotti, M. Barsuglia, D. Barta, J. Bartlett, M. A. Barton, I. Bartos, R. Bassiri, A. Basti, J. C. Batch, C. Baune, V. Bavigadda, M. Bazzan, B. Behnke, M. Bejger, C. Belczynski, A. S. Bell, C. J. Bell, B. K. Berger, J. Bergman, G. Bergmann, C. P. L. Berry, D. Bersanetti, A. Bertolini, J. Betzwieser, S. Bhagwat, R. Bhandare, I. A. Bilenko, G. Billingsley, J. Birch, I. A. Birney, O. Birnholtz, S. Biscans, A. Bisht, M. Bitossi, C. Biwer, M. A. Bizouard, J. K. Blackburn, C. D. Blair, D. G. Blair, R. M. Blair, S. Bloemen, O. Bock, T. P. Bodiya, M. Boer, G. Bogaert, C. Bogan, A. Bohe, P. Bojtos, C. Bond, F. Bondu, R. Bonnand, B. A. Boom, R. Bork, V. Boschi, S. Bose, Y. Bouffanais, A. Bozzi, C. Bradaschia, P. R. Brady, V. B. Braginsky, M. Branchesi, J. E. Brau, T. Briant, A. Brillet, M. Brinkmann, V. Brisson, P. Brockill, A. F. Brooks, D. A. Brown, D. D. Brown, N. M. Brown, C. C. Buchanan, A. Buikema, T. Bulik, H. J. Bulten, A. Buonanno, D. Buskulic, C. Buy, R. L. Byer, M. Cabero, L. Cadonati, G. Cagnoli, C. Cahillane, J. Calderón Bustillo, T. Callister, E. Calloni, J. B. Camp, K. C. Cannon, J. Cao, C. D. Capano, E. Capocasa, F. Carbognani, S. Caride, J. Casanueva Diaz, C. Casentini, S. Caudill, M. Cavaglià, F. Cavalier, R. Cavalieri, G. Cella, C. B. Cepeda, L. Cerboni Baiardi, G. Cerretani, E. Cesarini, R. Chakraborty, T. Chalermsoongsak, S. J. Chamberlin, M. Chan, S. Chao, P. Charlton, E. Chassande-Mottin, H. Y. Chen, Y. Chen, C. Cheng, A. Chincarini, A. Chi-

- ummo, H. S. Cho, M. Cho, J. H. Chow, N. Christensen, Q. Chu, S. Chua, S. Chung, G. Ciani, F. Clara, J. A. Clark, F. Cleva, E. Coccia, P.-F. Cohadon, A. Colla, C. G. Collette, L. Cominsky, M. Constancio, A. Conte, L. Conti, D. Cook, T. R. Corbitt, N. Cornish, A. Corsi, S. Cortese, C. A. Costa, M. W. Coughlin, S. B. Coughlin, J.-P. Coulon, S. T. Countryman, P. Couvares, E. E. Cowan, D. M. Coward, M. J. Cowart, D. C. Coyne, R. Coyne, K. Craig, J. D. E. Creighton, T. D. Creighton, J. Cripe, S. G. Crowder, A. M. Cruise, A. Cumming, L. Cunningham, E. Cuoco, T. Dal Canton, S. L. Danilishin, S. D'Antonio, K. Danzmann, N. S. Darman, C. F. Da Silva Costa, V. Dattilo, I. Dave, H. P. Daveloza, M. Davier, G. S. Davies, E. J. Daw, R. Day, S. De, D. DeBra, G. Debreczeni, J. Degallaix, M. De Laurentis, S. Deléglise, W. Del Pozzo, T. Denker, T. Dent, H. Dereli, V. Dergachev, R. T. DeRosa, R. De Rosa, R. DeSalvo, S. Dhurandhar, M. C. Díaz, L. Di Fiore, M. Di Giovanni, A. Di Lieto, S. Di Pace, I. Di Palma, A. Di Virgilio, G. Dojcinoski, V. Dolique, F. Donovan, K. L. Dooley, S. Doravari, R. Douglas, T. P. Downes, M. Drago, R. W. P. Drever, J. C. Driggers, Z. Du, M. Ducrot, S. E. Dwyer, T. B. Edo, M. C. Edwards, A. Effler, H.-B. Eggenstein, P. Ehrens, J. Eichholz, S. S. Eikenberry, W. Engels, R. C. Essick, T. Etzel, M. Evans, T. M. Evans, R. Everett, M. Factourovich, V. Fafone, H. Fair, S. Fairhurst, X. Fan, Q. Fang, S. Farinon, B. Farr, W. M. Farr, M. Favata, M. Fays, H. Fehrmann, M. M. Fejer, D. Feldbaum, I. Ferrante, E. C. Ferreira, F. Ferrini, F. Fidecaro, L. S. Finn, I. Fiori, D. Fiorucci, R. P. Fisher, R. Flaminio, M. Fletcher, H. Fong, J.-D. Fournier, S. Franco, S. Frasca, F. Frasconi, M. Frede, Z. Frei, A. Freise, R. Frey, V. Frey, T. T. Fricke, P. Fritschel, V. V. Frolov, P. Fulda, M. Fyffe, H. A. G. Gabard, J. R. Gair, L. Gammaitoni, S. G. Gaonkar, F. Garufi, A. Gatto, G. Gaur, N. Gehrels, G. Gemme, B. Gendre, E. Genin, A. Gennai, J. George, L. Gergely, V. Germain, Abhirup Ghosh, Archisman Ghosh, S. Ghosh, J. A. Giaime, K. D. Giardina, A. Giazotto, K. Gill, A. Glaefke, J. R. Gleason, E. Goetz, R. Goetz, L. Gondan, G. González, J. M. Gonzalez Castro, A. Gopakumar, N. A. Gordon, M. L. Gorodetsky, S. E. Gossan, M. Gosselin, R. Gouaty, C. Graef, P. B. Graff, M. Granata, A. Grant, S. Gras, C. Gray, G. Greco, A. C. Green, R. J. S. Greenhalgh, P. Groot, H. Grote, S. Grunewald, G. M. Guidi, X. Guo, A. Gupta, M. K. Gupta, K. E. Gushwa, E. K. Gustafson, R. Gustafson, J. J. Hacker, B. R. Hall, E. D. Hall, G. Hammond, M. Haney, M. M. Hanke, J. Hanks, C. Hanna, M. D. Hannam, J. Hanson, T. Hardwick, J. Harms, G. M. Harry, I. W. Harry, M. J. Hart, M. T. Hartman, C.-J. Haster, K. Haughian, J. Healy, J. Heefner, A. Heidmann, M. C. Heintze, G. Heinzl, H. Heitmann, P. Hello, G. Hemming, M. Hendry, I. S. Heng, J. Hennig, A. W. Heptonstall, M. Heurs, S. Hild, D. Hoak, K. A. Hodge, D. Hoffman, S. E. Hollitt, K. Holt, D. E. Holz, P. Hopkins, D. J. Hosken, J. Hough, E. A. Houston, E. J. Howell, Y. M. Hu, S. Huang, E. A. Huerta, D. Huet, B. Hughey, S. Husa, S. H. Huttner, T. Huynh-Dinh, A. Idrisy, N. Indik, D. R. Ingram, R. Inta, H. N. Isa, J.-M. Isac, M. Isi, G. Islas, T. Isogai, B. R. Iyer, K. Izumi, M. B. Jacobson, T. Jacqmin, H. Jang, K. Jani, P. Jaranowski, S. Jawahar, F. Jiménez-Forteza, W. W. Johnson, N. K. Johnson-McDaniel, D. I. Jones, R. Jones, R. J. G. Jonker, L. Ju, K. Haris, C. V. Kalaghatgi, V. Kalogera, S. Kandhasamy, G. Kang, J. B. Kanner, S. Karki, M. Kasprzack, E. Katsavounidis, W. Katzman, S. Kaufer, T. Kaur, K. Kawabe, F. Kawazoe, F. Kéfélian, M. S. Kehl, D. Keitel, D. B. Kelley, W. Kells, R. Kennedy, D. G. Keppel, J. S. Key, A. Khalaidovski, F. Y. Khalili, I. Khan, S. Khan,

- Z. Khan, E. A. Khazanov, N. Kijbunchoo, C. Kim, J. Kim, K. Kim, Nam-Gyu Kim, Namjun Kim, Y.-M. Kim, E. J. King, P. J. King, D. L. Kinzel, J. S. Kissel, L. Kleybolte, S. Klimenko, S. M. Koehlenbeck, K. Kokeyama, S. Koley, V. Kondrashov, A. Kontos, S. Koranda, M. Korobko, W. Z. Korth, I. Kowalska, D. B. Kozak, V. Kringel, B. Krishnan, A. Królak, C. Krueger, G. Kuehn, P. Kumar, R. Kumar, L. Kuo, A. Kutynia, P. Kwee, B. D. Lackey, M. Landry, J. Lange, B. Lantz, P. D. Lasky, A. Lazzarini, C. Lazzaro, P. Leaci, S. Leavey, E. O. Lebigot, C. H. Lee, H. K. Lee, H. M. Lee, K. Lee, A. Lenon, M. Leonardi, J. R. Leong, N. Leroy, N. Letendre, Y. Levin, B. M. Levine, T. G. F. Li, A. Libson, T. B. Littenberg, N. A. Lockerbie, J. Logue, A. L. Lombardi, L. T. London, J. E. Lord, M. Lorenzini, V. Lorette, M. Lormand, G. Losurdo, J. D. Lough, C. O. Lousto, G. Lovelace, H. Lück, A. P. Lundgren, J. Luo, R. Lynch, Y. Ma, T. MacDonald, B. Machenschalk, M. MacInnis, D. M. Macleod, F. Magaña Sandoval, R. M. Magee, M. Mageswaran, E. Majorana, I. Maksimovic, V. Malvezzi, N. Man, I. Mandel, V. Mandic, V. Mangano, G. L. Mansell, M. Manske, M. Mantovani, F. Marchesoni, F. Marion, S. Márka, Z. Márka, A. S. Markosyan, E. Maros, F. Martelli, L. Martellini, I. W. Martin, R. M. Martin, D. V. Martynov, J. N. Marx, K. Mason, A. Masserot, T. J. Massinger, M. Masso-Reid, F. Matichard, L. Matone, N. Mavalvala, N. Mazumder, G. Mazzolo, R. McCarthy, D. E. McClelland, S. McCormick, S. C. McGuire, G. McIntyre, J. McIver, D. J. McManus, S. T. McWilliams, D. Meacher, G. D. Meadors, J. Meidam, A. Melatos, G. Mendell, D. Mendoza-Gandara, R. A. Mercer, E. Merilh, M. Merzougui, S. Meshkov, C. Messenger, C. Messick, P. M. Meyers, F. Mezzani, H. Miao, C. Michel, H. Middleton, E. E. Mikhailov, L. Milano, J. Miller, M. Millhouse, Y. Minenkov, J. Ming, S. Mirshekari, C. Mishra, S. Mitra, V. P. Mitrofanov, G. Mitselmakher, R. Mittleman, A. Moggi, M. Mohan, S. R. P. Mohapatra, M. Montani, B. C. Moore, C. J. Moore, D. Moraru, G. Moreno, S. R. Morris, K. Mossavi, B. Mours, C. M. Mow-Lowry, C. L. Mueller, G. Mueller, A. W. Muir, Arunava Mukherjee, D. Mukherjee, S. Mukherjee, N. Mukund, A. Mullavey, J. Munch, D. J. Murphy, P. G. Murray, A. Mytidis, I. Nardecchia, L. Naticchioni, R. K. Nayak, V. Necula, K. Nedkova, G. Nelemans, M. Neri, A. Neunzert, G. Newton, T. T. Nguyen, A. B. Nielsen, S. Nissanke, A. Nitz, F. Nocera, D. Nolting, M. E. N. Normandin, L. K. Nuttall, J. Oberling, E. Ochsner, J. O'Dell, E. Oelker, G. H. Ogín, J. J. Oh, S. H. Oh, F. Ohme, M. Oliver, P. Oppermann, Richard J. Oram, B. O'Reilly, R. O'Shaughnessy, C. D. Ott, D. J. Ottaway, R. S. Ottens, H. Overmeyer, B. J. Owen, A. Pai, S. A. Pai, J. R. Palamos, O. Palashov, C. Palomba, A. Pal Singh, H. Pan, Y. Pan, C. Pankow, F. Pannarale, B. C. Pant, F. Paoletti, A. Paoli, M. A. Papa, H. R. Paris, W. Parker, D. Pascucci, A. Pasqualetti, R. Passaquieti, D. Passuello, B. Patricelli, Z. Patrick, B. L. Pearlstone, M. Pedraza, R. Pedurand, L. Pekowsky, A. Pele, S. Penn, A. Perreca, H. P. Pfeiffer, M. Phelps, O. Piccinni, M. Pichot, M. Pickenpack, F. Piergiovanni, V. Pierro, G. Pillant, L. Pinard, I. M. Pinto, M. Pitkin, J. H. Poeld, R. Poggiani, P. Popolizio, A. Post, J. Powell, J. Prasad, V. Predoi, S. S. Premachandra, T. Prestegard, L. R. Price, M. Prijatelj, M. Principe, S. Privitera, R. Prix, G. A. Prodi, L. Prokhorov, O. Puncken, M. Punturo, P. Puppo, M. Pürrer, H. Qi, J. Qin, V. Quetschke, E. A. Quintero, R. Quitzow-James, F. J. Raab, D. S. Rabeling, H. Radkins, P. Raffai, S. Raja, M. Rakhmanov, C. R. Ramet, P. Rapagnani, V. Raymond, M. Razzano, V. Re, J. Read, C. M. Reed, T. Regimbau,

- L. Rei, S. Reid, D. H. Reitze, H. Rew, S. D. Reyes, F. Ricci, K. Riles, N. A. Robertson, R. Robie, F. Robinet, A. Rocchi, L. Rolland, J. G. Rollins, V. J. Roma, J. D. Romano, R. Romano, G. Romanov, J. H. Romie, D. Rosińska, S. Rowan, A. Rüdigger, P. Ruggi, K. Ryan, S. Sachdev, T. Sadecki, L. Sadeghian, L. Salconi, M. Saleem, F. Salemi, A. Samajdar, L. Sammut, L. M. Sampson, E. J. Sanchez, V. Sandberg, B. Sandeen, G. H. Sanders, J. R. Sanders, B. Sassolas, B. S. Sathyaprakash, P. R. Saulson, O. Sauter, R. L. Savage, A. Sawadsky, P. Schale, R. Schilling, J. Schmidt, P. Schmidt, R. Schnabel, R. M. S. Schofield, A. Schönbeck, E. Schreiber, D. Schuette, B. F. Schutz, J. Scott, S. M. Scott, D. Sellers, A. S. Sengupta, D. Sentenac, V. Sequino, A. Sergeev, G. Serna, Y. Setyawati, A. Seigny, D. A. Shaddock, T. Shaffer, S. Shah, M. S. Shahriar, M. Shaltev, Z. Shao, B. Shapiro, P. Shawhan, A. Sheperd, D. H. Shoemaker, D. M. Shoemaker, K. Siellez, X. Siemens, D. Sigg, A. D. Silva, D. Simakov, A. Singer, L. P. Singer, A. Singh, R. Singh, A. Singhal, A. M. Sintès, B. J. J. Slagmolen, J. R. Smith, M. R. Smith, N. D. Smith, R. J. E. Smith, E. J. Son, B. Sorazu, F. Sorrentino, T. Souradeep, A. K. Srivastava, A. Staley, M. Steinke, J. Steinlechner, S. Steinlechner, D. Steinmeyer, B. C. Stephens, S. P. Stevenson, R. Stone, K. A. Strain, N. Straniero, G. Stratta, N. A. Strauss, S. Strigin, R. Sturani, A. L. Stuver, T. Z. Summerscales, L. Sun, P. J. Sutton, B. L. Swinkels, M. J. Szczepańczyk, M. Tacca, D. Talukder, D. B. Tanner, M. Tápai, S. P. Tarabrin, A. Taracchini, R. Taylor, T. Theeg, M. P. Thirugnanasambandam, E. G. Thomas, M. Thomas, P. Thomas, K. A. Thorne, K. S. Thorne, E. Thrane, S. Tiwari, V. Tiwari, K. V. Tokmakov, C. Tomlinson, M. Tonelli, C. V. Torres, C. I. Torrie, D. Töyrä, F. Travasso, G. Traylor, D. Trifirò, M. C. Tringali, L. Trozzo, M. Tse, M. Turconi, D. Tuyenbayev, D. Ugolini, C. S. Unnikrishnan, A. L. Urban, S. A. Usman, H. Vahlbruch, G. Vajente, G. Valdes, M. Vallisneri, N. van Bakel, M. van Beuzekom, J. F. J. van den Brand, C. Van Den Broeck, D. C. Vander-Hyde, L. van der Schaaf, J. V. van Heijningen, A. A. van Veggel, M. Vardaro, S. Vass, M. Vasúth, R. Vaulin, A. Vecchio, G. Vedovato, J. Veitch, P. J. Veitch, K. Venkateswara, D. Verkindt, F. Vetrano, A. Viceré, S. Vinciguerra, D. J. Vine, J.-Y. Vinet, S. Vitale, T. Vo, H. Vocca, C. Vorvick, D. Voss, W. D. Vousden, S. P. Vyatchanin, A. R. Wade, L. E. Wade, M. Wade, S. J. Waldman, M. Walker, L. Wallace, S. Walsh, G. Wang, H. Wang, M. Wang, X. Wang, Y. Wang, H. Ward, R. L. Ward, J. Warner, M. Was, B. Weaver, L.-W. Wei, M. Weinert, A. J. Weinstein, R. Weiss, T. Welborn, L. Wen, P. Weßels, T. Westphal, K. Wette, J. T. Whelan, S. E. Whitcomb, D. J. White, B. F. Whiting, K. Wiesner, C. Wilkinson, P. A. Willems, L. Williams, R. D. Williams, A. R. Williamson, J. L. Willis, B. Willke, M. H. Wimmer, L. Winkelmann, W. Winkler, C. C. Wipf, A. G. Wiseman, H. Wittel, G. Woan, J. Worden, J. L. Wright, G. Wu, J. Yablon, I. Yakushin, W. Yam, H. Yamamoto, C. C. Yancey, M. J. Yap, H. Yu, M. Yvert, A. Zadrožny, L. Zangrando, M. Zanolin, J.-P. Zendri, M. Zevin, F. Zhang, L. Zhang, M. Zhang, Y. Zhang, C. Zhao, M. Zhou, Z. Zhou, X. J. Zhu, M. E. Zucker, S. E. Zuraw, and J. Zweizig. Observation of gravitational waves from a binary black hole merger. *Phys. Rev. Lett.*, 116:061102, Feb 2016. doi: 10.1103/PhysRevLett.116.061102. URL <https://link.aps.org/doi/10.1103/PhysRevLett.116.061102>.
- B. P. Abbott, R. Abbott, T. D. Abbott, S. Abraham, F. Acernese, K. Ackley, C. Adams, R. X. Adhikari, V. B. Adya, C. Affeldt, M. Agathos, K. Agatsuma, N. Aggarwal, O. D. Aguiar, L. Aiello, A. Ain, P. Ajith, G. Allen, A. Allocca, M. A. Aloy, P. A.

- Altin, A. Amato, A. Ananyeva, S.B. Anderson, W.G. Anderson, S.V. Angelova, S. Antier, S. Appert, K. Arai, M.C. Araya, J.S. Areeda, M. Arène, N. Arnaud, K.G. Arun, S. Ascenzi, G. Ashton, S.M. Aston, P. Astone, F. Aubin, P. Aufmuth, K. AultONeal, C. Austin, V. Avendano, A. Avila-Alvarez, S. Babak, P. Bacon, F. Badaracco, M.K.M. Bader, S. Bae, P.T. Baker, F. Baldaccini, G. Ballard, S.W. Ballmer, S. Banagiri, J.C. Barayoga, S.E. Barclay, B.C. Barish, D. Barker, K. Barrett, S. Barnum, F. Barone, B. Barr, L. Barsotti, M. Barsuglia, D. Barta, J. Bartlett, I. Bartos, R. Bassiri, A. Basti, M. Bawaj, J.C. Bayley, M. Bazzan, B. Bécsy, M. Beger, I. Belahcene, A.S. Bell, D. Beniwal, B.K. Berger, G. Bergmann, S. Bernuzzi, J.J. Bero, C.P.L. Berry, D. Bersanetti, A. Bertolini, J. Betzwieser, R. Bhandare, J. Bidler, I.A. Bilenko, S.A. Bilgili, G. Billingsley, J. Birch, I.A. Birney, O. Birnholtz, S. Biscans, S. Biscoveanu, A. Bisht, M. Bitossi, M.A. Bizouard, J.K. Blackburn, J. Blackman, C.D. Blair, D.G. Blair, R.M. Blair, S. Bloemen, N. Bode, M. Boer, Y. Boetzel, G. Bogaert, F. Bondu, E. Bonilla, R. Bonnand, P. Booker, B.A. Boom, C.D. Booth, R. Bork, V. Boschi, S. Bose, K. Bossie, V. Bossilkov, J. Bosveld, Y. Bouffanais, A. Bozzi, C. Bradaschia, P.R. Brady, A. Bramley, M. Branchesi, J.E. Brau, T. Briant, J.H. Briggs, F. Brighenti, A. Brillet, M. Brinkmann, V. Brisson, P. Brockill, A.F. Brooks, D.D. Brown, S. Brunett, A. Buikema, T. Bulik, H.J. Bulten, A. Buonanno, D. Buskulic, M.J. Bustamante Rosell, C. Buy, R.L. Byer, M. Cabero, L. Cadonati, G. Cagnoli, C. Cahillane, J. Calderón Bustillo, T.A. Callister, E. Calloni, J.B. Camp, W.A. Campbell, M. Canepa, K.C. Cannon, H. Cao, J. Cao, E. Capocasa, F. Carbognani, S. Caride, M.F. Carney, G. Carullo, J. Casanueva Diaz, C. Casentini, S. Caudill, M. Cavaglià, F. Cavalier, R. Cavalieri, G. Cella, P. Cerdá-Durán, G. Cerretani, E. Cesarini, O. Chaibi, K. Chakravarti, S.J. Chamberlin, M. Chan, S. Chao, P. Charlton, E.A. Chase, E. Chassande-Mottin, D. Chatterjee, M. Chaturvedi, K. Chatziioannou, B.D. Cheeseboro, H.Y. Chen, X. Chen, Y. Chen, H.-P. Cheng, C.K. Cheong, H.Y. Chia, A. Chincarini, A. Chiummo, G. Cho, H.S. Cho, M. Cho, N. Christensen, Q. Chu, S. Chua, K.W. Chung, S. Chung, G. Ciani, A.A. Ciobanu, R. Ciolfi, F. Cipriano, A. Cirone, F. Clara, J.A. Clark, P. Clearwater, F. Cleva, C. Cocchieri, E. Coccia, P.-F. Cohadon, D. Cohen, R. Colgan, M. Colleoni, C.G. Collette, C. Collins, L.R. Cominsky, M. Constancio, L. Conti, S.J. Cooper, P. Corban, T.R. Corbitt, I. Cordero-Carrión, K.R. Corley, N. Cornish, A. Corsi, S. Cortese, C.A. Costa, R. Cotesta, M.W. Coughlin, S.B. Coughlin, J.-P. Coulon, S.T. Countryman, P. Couvares, P.B. Covas, E.E. Cowan, D.M. Coward, M.J. Cowart, D.C. Coyne, R. Coyne, J.D.E. Creighton, T.D. Creighton, J. Cripe, M. Croquette, S.G. Crowder, T.J. Cullen, A. Cumming, L. Cunningham, E. Cuoco, T. Dal Canton, G. Dálya, S.L. Danilishin, S. D'Antonio, K. Danzmann, A. Dasgupta, C.F. Da Silva Costa, L.E.H. Datrier, V. Dattilo, I. Dave, M. Davier, D. Davis, E.J. Daw, D. DeBra, M. Deenadayalan, J. Degallaix, M. De Laurentis, S. Deléglise, W. Del Pozzo, L.M. DeMarchi, N. Demos, T. Dent, R. De Pietri, J. Derby, R. De Rosa, C. De Rossi, R. DeSalvo, O. de Varona, S. Dhurandhar, M.C. Díaz, T. Dietrich, L. Di Fiore, M. Di Giovanni, T. Di Girolamo, A. Di Lieto, B. Ding, S. Di Pace, I. Di Palma, F. Di Renzo, A. Dmitriev, Z. Doctor, F. Donovan, K.L. Dooley, S. Doravari, I. Dorrington, T.P. Downes, M. Drago, J.C. Driggers, Z. Du, J.-G. Ducoin, P. Dupej, S.E. Dwyer, P.J. Easter, T.B. Edo, M.C. Edwards, A. Effler, P. Ehrens, J. Eichholz, S.S. Eiken-

- berry, M. Eisenmann, R. A. Eisenstein, R. C. Essick, H. Estelles, D. Estevez, Z. B. Etienne, T. Etzel, M. Evans, T. M. Evans, V. Fafone, H. Fair, S. Fairhurst, X. Fan, S. Farinon, B. Farr, W. M. Farr, E. J. Fauchon-Jones, M. Favata, M. Fays, M. Fazio, C. Fee, J. Feicht, M. M. Fejer, F. Feng, A. Fernandez-Galiana, I. Ferrante, E. C. Ferreira, T. A. Ferreira, F. Ferrini, F. Fidecaro, I. Fiori, D. Fiorucci, M. Fishbach, R. P. Fisher, J. M. Fishner, M. Fitz-Axen, R. Flaminio, M. Fletcher, E. Flynn, H. Fong, J. A. Font, P. W. F. Forsyth, J.-D. Fournier, S. Frasca, F. Frasconi, Z. Frei, A. Freise, R. Frey, V. Frey, P. Fritschel, V. V. Frolov, P. Fulda, M. Fyffe, H. A. Gabbard, B. U. Gadre, S. M. Gaebel, J. R. Gair, L. Gammaitoni, M. R. Ganija, S. G. Gaonkar, A. Garcia, C. García-Quirós, F. Garufi, B. Gateley, S. Gaudio, G. Gaur, V. Gayathri, G. Gemme, E. Genin, A. Gennai, D. George, J. George, L. Gergely, V. Germain, S. Ghonge, Abhirup Ghosh, Archisman Ghosh, S. Ghosh, B. Giacomazzo, J. A. Giaime, K. D. Giardina, A. Giazotto, K. Gill, G. Giordano, L. Glover, P. Godwin, E. Goetz, R. Goetz, B. Goncharov, G. González, J. M. Gonzalez Castro, A. Gopakumar, M. L. Gorodetsky, S. E. Gossan, M. Gosselin, R. Gouaty, A. Grado, C. Graef, M. Granata, A. Grant, S. Gras, P. Grassia, C. Gray, R. Gray, G. Greco, A. C. Green, R. Green, E. M. Gretarsson, P. Groot, H. Grote, S. Grunewald, P. Gruning, G. M. Guidi, H. K. Gulati, Y. Guo, A. Gupta, M. K. Gupta, E. K. Gustafson, R. Gustafson, L. Haegel, O. Halim, B. R. Hall, E. D. Hall, E. Z. Hamilton, G. Hammond, M. Haney, M. M. Hanke, J. Hanks, C. Hanna, M. D. Hannam, O. A. Hannuksela, J. Hanson, T. Hardwick, K. Haris, J. Harms, G. M. Harry, I. W. Harry, C.-J. Haster, K. Haughian, F. J. Hayes, J. Healy, A. Heidmann, M. C. Heintze, H. Heitmann, P. Hello, G. Hemming, M. Hendry, I. S. Heng, J. Hennig, A. W. Heptonstall, Francisco Hernandez Vivanco, M. Heurs, S. Hild, T. Hinderer, D. Hoak, S. Hochheim, D. Hofman, A. M. Holgado, N. A. Holland, K. Holt, D. E. Holz, P. Hopkins, C. Horst, J. Hough, E. J. Howell, C. G. Hoy, A. Hreibi, Y. Huang, E. A. Huerta, D. Huet, B. Hughey, M. Hulko, S. Husa, S. H. Huttner, T. Huynh-Dinh, B. Idzkowski, A. Iess, C. Ingram, R. Inta, G. Intini, B. Irwin, H. N. Isa, J.-M. Isac, M. Isi, B. R. Iyer, K. Izumi, T. Jacqmin, S. J. Jadhav, K. Jani, N. N. Janthapur, P. Jaranowski, A. C. Jenkins, J. Jiang, D. S. Johnson, N. K. Johnson-McDaniel, A. W. Jones, D. I. Jones, R. Jones, R. J. G. Jonker, L. Ju, J. Junker, C. V. Kalaghatgi, V. Kalogera, B. Kamai, S. Kandhasamy, G. Kang, J. B. Kanner, S. J. Kapadia, S. Karki, K. S. Karvinen, R. Kashyap, M. Kasprzak, S. Katsanevas, E. Katsavounidis, W. Katzman, S. Kaufer, K. Kawabe, N. V. Keerthana, F. Kéfélian, D. Keitel, R. Kennedy, J. S. Key, F. Y. Khalili, H. Khan, I. Khan, S. Khan, Z. Khan, E. A. Khazanov, M. Khursheed, N. Kijbunchoo, Chunglee Kim, J. C. Kim, K. Kim, W. Kim, W. S. Kim, Y.-M. Kim, C. Kimball, E. J. King, P. J. King, M. Kinley-Hanlon, R. Kirchhoff, J. S. Kissel, L. Kleybolte, J. H. Klika, S. Klimenko, T. D. Knowles, P. Koch, S. M. Koehlenbeck, G. Koekoek, S. Koley, V. Kondrashov, A. Kontos, N. Koper, M. Korobko, W. Z. Korth, I. Kowalska, D. B. Kozak, V. Kringel, N. Krishnendu, A. Królak, G. Kuehn, A. Kumar, P. Kumar, R. Kumar, S. Kumar, L. Kuo, A. Kutynia, S. Kwang, B. D. Lackey, K. H. Lai, T. L. Lam, M. Landry, B. B. Lane, R. N. Lang, J. Lange, B. Lantz, R. K. Lanza, A. Lartaux-Vollard, P. D. Lasky, M. Laxen, A. Lazzarini, C. Lazzaro, P. Leaci, S. Leavey, Y. K. Lecoeuche, C. H. Lee, H. K. Lee, H. M. Lee, H. W. Lee, J. Lee, K. Lee, J. Lehmann, A. Lenon, N. Leroy, N. Letendre, Y. Levin, J. Li, K. J. L. Li, T. G. F. Li, X. Li, F. Lin, F. Linde, S. D. Linker, T. B.

- Littenberg, J. Liu, X. Liu, R. K. L. Lo, N. A. Lockerbie, L. T. London, A. Longo, M. Lorenzini, V. Lorient, M. Lormand, G. Losurdo, J. D. Lough, C. O. Lousto, G. Lovelace, M. E. Lower, H. Lück, D. Lumaca, A. P. Lundgren, R. Lynch, Y. Ma, R. Macas, S. Macfoy, M. MacInnis, D. M. Macleod, A. Macquet, F. Magaña-Sandoval, L. Magaña Zertuche, R. M. Magee, E. Majorana, I. Maksimovic, A. Malik, N. Man, V. Mandic, V. Mangano, G. L. Mansell, M. Manske, M. Mantovani, F. Marchesoni, F. Marion, S. Márka, Z. Márka, C. Markakis, A. S. Markosyan, A. Markowitz, E. Maros, A. Marquina, S. Marsat, F. Martelli, I. W. Martin, R. M. Martin, D. V. Martynov, K. Mason, E. Massera, A. Masserot, T. J. Massinger, M. Masso-Reid, S. Mastrogiovanni, A. Matas, F. Matichard, L. Matone, N. Mavalvala, N. Mazumder, J. J. McCann, R. McCarthy, D. E. McClelland, S. McCormick, L. McCuller, S. C. McGuire, J. McIver, D. J. McManus, T. McRae, S. T. McWilliams, D. Meacher, G. D. Meadors, M. Mehmet, A. K. Mehta, J. Meidam, A. Melatos, G. Mendell, R. A. Mercer, L. Mereni, E. L. Merilh, M. Merzougui, S. Meshkov, C. Messenger, C. Messick, R. Metzдорff, P. M. Meyers, H. Miao, C. Michel, H. Middleton, E. E. Mikhailov, L. Milano, A. L. Miller, A. Miller, M. Millhouse, J. C. Mills, M. C. Milovich-Goff, O. Minazzoli, Y. Minenkov, A. Mishkin, C. Mishra, T. Mistry, S. Mitra, V. P. Mitrofanov, G. Mitselmakher, R. Mittleman, G. Mo, D. Moffa, K. Mogushi, S. R. P. Mohapatra, M. Montani, C. J. Moore, D. Moraru, G. Moreno, S. Morisaki, B. Mours, C. M. Mow-Lowry, Arunava Mukherjee, D. Mukherjee, S. Mukherjee, N. Mukund, A. Mullavey, J. Munch, E. A. Muñoz, M. Muratore, P. G. Murray, A. Nagar, I. Nardecchia, L. Naticchioni, R. K. Nayak, J. Neilson, G. Nelemans, T. J. N. Nelson, M. Nery, A. Neunzert, K. Y. Ng, S. Ng, P. Nguyen, D. Nichols, A. B. Nielsen, S. Nissanke, A. Nitz, F. Nocera, C. North, L. K. Nuttall, M. Obergaullinger, J. Oberling, B. D. O'Brien, G. D. O'Dea, G. H. Ogin, J. J. Oh, S. H. Oh, F. Ohme, H. Ohta, M. A. Okada, M. Oliver, P. Oppermann, Richard J. Oram, B. O'Reilly, R. G. Ormiston, L. F. Ortega, R. O'Shaughnessy, S. Ossokine, D. J. Ottaway, H. Overmire, B. J. Owen, A. E. Pace, G. Pagano, M. A. Page, A. Pai, S. A. Pai, J. R. Palamos, O. Palashov, C. Palomba, A. Pal-Singh, Huang-Wei Pan, B. Pang, P. T. H. Pang, C. Pankow, F. Pannarale, B. C. Pant, F. Paoletti, A. Paoli, M. A. Papa, A. Parida, W. Parker, D. Pascucci, A. Pasqualetti, R. Passaquietti, D. Pasuello, M. Patil, B. Patricelli, B. L. Pearlstone, C. Pedersen, M. Pedraza, R. Pedurand, A. Pele, S. Penn, A. Perego, C. J. Perez, A. Perreca, H. P. Pfeiffer, M. Phelps, K. S. Phukon, O. J. Piccinni, M. Pichot, F. Piergiovanni, G. Pillant, L. Pinard, M. Pirello, M. Pitkin, R. Poggiani, D. Y. T. Pong, S. Ponrathnam, P. Popolizio, E. K. Porter, J. Powell, A. K. Prajapati, J. Prasad, K. Prasai, R. Prasanna, G. Pratten, T. Prestegard, S. Privitera, G. A. Prodi, L. G. Prokhorov, O. Puncken, M. Punturo, P. Puppo, M. Pürner, H. Qi, V. Quetschke, P. J. Quinonez, E. A. Quintero, R. Quitzow-James, F. J. Raab, H. Radkins, N. Radulescu, P. Raffai, S. Raja, C. Rajan, B. Rajbhandari, M. Rakhmanov, K. E. Ramirez, A. Ramos-Buades, Javed Rana, K. Rao, P. Rapagnani, V. Raymond, M. Razzano, J. Read, T. Regimbau, L. Rei, S. Reid, D. H. Reitze, W. Ren, F. Ricci, C. J. Richardson, J. W. Richardson, P. M. Ricker, G. M. Riemschneider, K. Riles, M. Rizzo, N. A. Robertson, R. Robie, F. Robinet, A. Rocchi, L. Rolland, J. G. Rollins, V. J. Roma, M. Romanelli, R. Romano, C. L. Romel, J. H. Romie, K. Rose, D. Rosińska, S. G. Rosofsky, M. P. Ross, S. Rowan, A. Rüdiger, P. Ruggi, G. Rutins, K. Ryan, S. Sachdev, T. Sadecki, M. Sakellari-

adou, O. Salafia, L. Salconi, M. Saleem, F. Salemi, A. Samajdar, L. Sammut, E. J. Sanchez, L. E. Sanchez, N. Sanchis-Gual, V. Sandberg, J. R. Sanders, K. A. Santiago, N. Sarin, B. Sassolas, B. S. Sathyaprakash, P. R. Saulson, O. Sauter, R. L. Savage, P. Schale, M. Scheel, J. Scheuer, P. Schmidt, R. Schnabel, R. M. S. Schofield, A. Schönbeck, E. Schreiber, B. W. Schulte, B. F. Schutz, S. G. Schwalbe, J. Scott, S. M. Scott, E. Seidel, D. Sellers, A. S. Sengupta, N. Sennett, D. Sentenac, V. Sequino, A. Sergeev, Y. Setyawati, D. A. Shaddock, T. Shaffer, M. S. Shahriar, M. B. Shaner, L. Shao, P. Sharma, P. Shawhan, H. Shen, R. Shink, D. H. Shoemaker, D. M. Shoemaker, S. ShyamSundar, K. Siellez, M. Sieniawska, D. Sigg, A. D. Silva, L. P. Singer, N. Singh, A. Singhal, A. M. Sintès, S. Sitmukhambetov, V. Skliris, B. J. J. Slagmolen, T. J. Slaven-Blair, J. R. Smith, R. J. E. Smith, S. Somala, E. J. Son, B. Sorazu, F. Sorrentino, T. Souradeep, E. Sowell, A. P. Spencer, A. K. Srivastava, V. Srivastava, K. Staats, C. Stachie, M. Standke, D. A. Steer, M. Steinke, J. Steinlechner, S. Steinlechner, D. Steinmeyer, S. P. Stevenson, D. Stocks, R. Stone, D. J. Stops, K. A. Strain, G. Stratta, S. E. Strigin, A. Strunk, R. Sturani, A. L. Stuver, V. Sudhir, T. Z. Summerscales, L. Sun, S. Sunil, J. Suresh, P. J. Sutton, B. L. Swinkels, M. J. Szczepańczyk, M. Tacca, S. C. Tait, C. Talbot, D. Talukder, D. B. Tanner, M. Tápai, A. Taracchini, J. D. Tasson, R. Taylor, F. Thies, M. Thomas, P. Thomas, S. R. Thondapu, K. A. Thorne, E. Thrane, Shubhanshu Tiwari, Srishti Tiwari, V. Tiwari, K. Toland, M. Tonelli, Z. Tornasi, A. Torres-Forné, C. I. Torrie, D. Töyrä, F. Travasso, G. Traylor, M. C. Tringali, A. Trovato, L. Trozzo, R. Trudeau, K. W. Tsang, M. Tse, R. Tso, L. Tsukada, D. Tsuna, D. Tuyenbayev, K. Ueno, D. Ugolini, C. S. Unnikrishnan, A. L. Urban, S. A. Usman, H. Vahlbruch, G. Vajente, G. Valdes, N. van Bakel, M. van Beuzekom, J. F. J. van den Brand, C. Van Den Broeck, D. C. Vander-Hyde, J. V. van Heijningen, L. van der Schaaf, A. A. van Veggel, M. Vardaro, V. Varma, S. Vass, M. Vasúth, A. Vecchio, G. Vedovato, J. Veitch, P. J. Veitch, K. Venkateswara, G. Venugopalan, D. Verkindt, F. Vetrano, A. Viceré, A. D. Viets, D. J. Vine, J.-Y. Vinet, S. Vitale, T. Vo, H. Vocca, C. Vorvick, S. P. Vyatchanin, A. R. Wade, L. E. Wade, M. Wade, R. Walet, M. Walker, L. Wallace, S. Walsh, G. Wang, H. Wang, J. Z. Wang, W. H. Wang, Y. F. Wang, R. L. Ward, Z. A. Warden, J. Warner, M. Was, J. Watchi, B. Weaver, L.-W. Wei, M. Weinert, A. J. Weinstein, R. Weiss, F. Wellmann, L. Wen, E. K. Wessel, P. Weßels, J. W. Westhouse, K. Wette, J. T. Whelan, L. V. White, B. F. Whiting, C. Whittle, D. M. Wilken, D. Williams, A. R. Williamson, J. L. Willis, B. Willke, M. H. Wimmer, W. Winkler, C. C. Wipf, H. Wittel, G. Woan, J. Woehler, J. K. Wofford, J. Worden, J. L. Wright, D. S. Wu, D. M. Wysocki, L. Xiao, H. Yamamoto, C. C. Yancey, L. Yang, M. J. Yap, M. Yazback, D. W. Yeeles, Hang Yu, Haocun Yu, S. H. R. Yuen, M. Yvert, A. K. Zadrożny, M. Zanolin, F. Zappa, T. Zelenova, J.-P. Zendri, M. Zevin, J. Zhang, L. Zhang, T. Zhang, C. Zhao, M. Zhou, Z. Zhou, X. J. Zhu, A. B. Zimmerman, Y. Zlochower, M. E. Zucker, and J. Zweizig.

Gwtc-1: A gravitational-wave transient catalog of compact binary mergers observed by ligo and virgo during the first and second observing runs. *Physical Review X*, 9 (3), September 2019. ISSN 2160-3308. doi: 10.1103/physrevx.9.031040. URL <http://dx.doi.org/10.1103/PhysRevX.9.031040>.

F Acernese, M Agathos, K Agatsuma, D Aisa, N Allemandou, A Allocca, J Amarni, P Astone, G Balestri, G Ballardin, F Barone, J-P Baronick, M Barsuglia, A Basti, F Basti, Th S Bauer, V Bavigadda, M Bejger, M G Beker, C Belczynski, D Bersanetti,

- A Bertolini, M Bitossi, M A Bizouard, S Bloemen, M Blom, M Boer, G Bogaert, D Bondi, F Bondu, L Bonelli, R Bonnard, V Boschi, L Bosi, T Bouedo, C Bradaschia, M Branchesi, T Briant, A Brillet, V Brisson, T Bulik, H J Bulten, D Buskalic, C Buy, G Cagnoli, E Calloni, C Campeggi, B Canuel, F Carbognani, F Cavalier, R Cavalieri, G Cella, E Cesarini, E Chassande Mottin, A Chincarini, A Chiummo, S Chua, F Cleva, E Coccia, P-F Cohadon, A Colla, M Colombini, A Conte, J-P Coulon, E Cuoco, A Dalmaz, S D'Antonio, V Dattilo, M Davier, R Day, G Debreczeni, J Degallaix, S Deléglise, W Del Pozzo, H Dereli, R De Rosa, L Di Fiore, A Di Lieto, A Di Virgilio, M Doets, V Dolique, M Drago, M Ducrot, G Endrőczy, V Fafone, S Farinon, I Ferrante, F Ferrini, F Fidecaro, I Fiori, R Flaminio, J-D Fournier, S Franco, S Frasca, F Frasconi, L Gammaitoni, F Garufi, M Gaspard, A Gatto, G Gemme, B Gendre, E Genin, A Gennai, S Ghosh, L Giacobone, A Giazotto, R Gouaty, M Granata, G Greco, P Groot, G M Guidi, J Harms, A Heidmann, H Heitmann, P Hello, G Hemming, E Hennes, D Hofman, P Jaranowski, R J G Jonker, M Kasprzak, F Kéfélian, I Kowalska, M Kraan, A Królak, A Kutynia, C Lazzaro, M Leonardi, N Leroy, N Letendre, T G F Li, B Lieunard, M Lorenzini, V Lorient, G Losurdo, C Magazzù, E Majorana, I Maksimovic, V Malvezzi, N Man, V Mangano, M Mantovani, F Marchesoni, F Marion, J Marque, F Martelli, L Martellini, A Masserot, D Meacher, J Meidam, F Mezzani, C Michel, L Milano, Y Minenkov, A Moggi, M Mohan, M Montani, N Morgado, B Mours, F Mul, M F Nagy, I Nardecchia, L Naticchioni, G Nelemans, I Neri, M Neri, F Nocera, E Pacaud, C Palomba, F Paoletti, A Paoli, A Pasqualetti, R Passaquieti, D Passuello, M Perciballi, S Petit, M Pichot, F Piergiovanni, G Pillant, A Piluso, L Pinard, R Poggiani, M Prijatelj, G A Prodi, M Punturo, P Puppo, D S Rabeling, I Rácz, P Rappagnani, M Razzano, V Re, T Regimbau, F Ricci, F Robinet, A Rocchi, L Rolland, R Romano, D Rosińska, P Ruggi, E Saracco, B Sassolas, F Schimmel, D Sentenac, V Sequino, S Shah, K Siellez, N Straniero, B Swinkels, M Tacca, M Tonelli, F Travasso, M Turconi, G Vajente, N van Bakel, M van Beuzekom, J F J van den Brand, C Van Den Broeck, M V van der Sluys, J van Heijningen, M Vasúth, G Vedovato, J Veitch, D Verkindt, F Vetrano, A Viceré, J-Y Vinet, G Visser, H Vocca, R Ward, M Was, L-W Wei, M Yvert, A Zadrożny, and J-P Zendri. Advanced virgo: a second-generation interferometric gravitational wave detector. *Classical and Quantum Gravity*, 32(2):024001, dec 2014. doi: 10.1088/0264-9381/32/2/024001. URL <https://dx.doi.org/10.1088/0264-9381/32/2/024001>.
- T Akutsu, M Ando, K Arai, Y Arai, S Araki, A Araya, N Aritomi, Y Aso, S Bae, Y Bae, L Baiotti, R Bajpai, M A Barton, K Cannon, E Capocasa, M Chan, C Chen, K Chen, Y Chen, H Chu, Y K Chu, S Eguchi, Y Enomoto, R Flaminio, Y Fujii, M Fukunaga, M Fukushima, G Ge, A Hagiwara, S Haino, K Hasegawa, H Hayakawa, K Hayama, Y Himemoto, Y Hiranuma, N Hirata, E Hirose, Z Hong, B H Hsieh, C Z Huang, P Huang, Y Huang, B Ikenoue, S Imam, K Inayoshi, Y Inoue, K Ioka, Y Itoh, K Izumi, K Jung, P Jung, T Kajita, M Kamiizumi, N Kanda, G Kang, K Kawaguchi, N Kawai, T Kawasaki, C Kim, J C Kim, W S Kim, Y M Kim, N Kimura, N Kita, H Kitazawa, Y Kojima, K Kokeyama, K Komori, A K H Kong, K Kotake, C Kozakai, R Kozu, R Kumar, J Kume, C Kuo, H S Kuo, S Kuroyanagi, K Kusayanagi, K Kwak, H K Lee, H W Lee, R Lee, M Leonardi, L C C Lin, C Y Lin, F L Lin, G C Liu, L W Luo, M Marchio, Y Michimura, N Mio,

- O Miyakawa, A Miyamoto, Y Miyazaki, K Miyo, S Miyoki, S Morisaki, Y Moriwaki, K Nagano, S Nagano, K Nakamura, H Nakano, M Nakano, R Nakashima, T Narikawa, R Negishi, W T Ni, A Nishizawa, Y Obuchi, W Ogaki, J J Oh, S H Oh, M Ohashi, N Ohishi, M Ohkawa, K Okutomi, K Oohara, C P Ooi, S Os-hino, K Pan, H Pang, J Park, F E Peña Arellano, I Pinto, N Sago, S Saito, Y Saito, K Sakai, Y Sakai, Y Sakuno, S Sato, T Sato, T Sawada, T Sekiguchi, Y Sekiguchi, S Shibagaki, R Shimizu, T Shimoda, K Shimode, H Shinkai, T Shishido, A Shoda, K Somiya, E J Son, H Sotani, R Sugimoto, T Suzuki, T Suzuki, H Tagoshi, H Takahashi, R Takahashi, A Takamori, S Takano, H Takeda, M Takeda, H Tanaka, K Tanaka, K Tanaka, T Tanaka, T Tanaka, S Tanioka, E N Tapia San Martin, S Telada, T Tomaru, Y Tomigami, T Tomura, F Travasso, L Trozzo, T Tsang, K Tsubono, S Tsuchida, T Tsuzuki, D Tuyenbayev, N Uchikata, T Uchiyama, A Ueda, T Uehara, K Ueno, G Ueshima, F Uraguchi, T Ushiba, M H P M van Putten, H Vocca, J Wang, C Wu, H Wu, S Wu, W-R Xu, T Yamada, K Yamamoto, K Yamamoto, T Yamamoto, K Yokogawa, J Yokoyama, T Yokozawa, T Yoshioka, H Yuzurihara, S Zeidler, Y Zhao, and Z H Zhu. Overview of kagra: Detector design and construction history. *Progress of Theoretical and Experimental Physics*, 2021(5):05A101, 08 2020. ISSN 2050-3911. doi: 10.1093/ptep/ptaa125. URL <https://doi.org/10.1093/ptep/ptaa125>.
- Carlos Allende Prieto, David L. Lambert, and Martin Asplund. The Forbidden Abundance of Oxygen in the Sun. *ApJ*, 556(1):L63–L66, July 2001. doi: 10.1086/322874.
- Pau Amaro-Seoane, Jeff Andrews, Manuel Arca Sedda, Abbas Askar, Quentin Baghi, Razvan Balasov, Imre Bartos, Simone S. Bavera, Jillian Bellovary, Christopher P. L. Berry, Emanuele Berti, Stefano Bianchi, Laura Blecha, Stéphane Blondin, Tamara Bogdanović, Samuel Boissier, Matteo Bonetti, Silvia Bonoli, Elisa Bortolas, Katelyn Breivik, Pedro R. Capelo, Laurentiu Caramete, Federico Cattorini, Maria Charisi, Sylvain Chaty, Xian Chen, Martyna Chruślińska, Alvin J. K. Chua, Ross Church, Monica Colpi, Daniel D’Orazio, Camilla Danielski, Melvyn B. Davies, Pratika Dayal, Alessandra De Rosa, Andrea Derdzinski, Kyriakos Destounis, Massimo Dotti, Ioana Duțan, Irina Dvorkin, Gaia Fajj, Thierry Foglizzo, Saavik Ford, Jean-Baptiste Fouvry, Alessia Franchini, Tassos Fragos, Chris Fryer, Massimo Gaspari, Davide Gerosa, Luca Graziani, Paul Groot, Melanie Habouzit, Daryl Haggard, Zoltan Haiman, Wen-Biao Han, Alina Istrate, Peter H. Johansson, Fazeel Mahmood Khan, Tomas Kimpson, Kostas Kokkotas, Albert Kong, Valeriya Korol, Kyle Kremer, Thomas Kupfer, Astrid Lamberts, Shane Larson, Mike Lau, Dongliang Liu, Nicole Lloyd-Ronning, Giuseppe Lodato, Alessandro Lupi, Chung-Pei Ma, Tomas Mac-carone, Ilya Mandel, Alberto Mangiagli, Michela Mapelli, Stéphane Mathis, Lucio Mayer, Sean McGee, Berry McKernan, M. Coleman Miller, David F. Mota, Matthew Mumpower, Syeda S. Nasim, Gijs Nelemans, Scott Noble, Fabio Pacucci, Francesca Panessa, Vasileios Paschalidis, Hugo Pfister, Delphine Porquet, John Quenby, Angelo Ricarte, Friedrich K. Röpke, John Regan, Stephan Rosswog, Ashley Ruiters, Milton Ruiz, Jessie Runnoe, Raffaella Schneider, Jeremy Schnittman, Amy Secunda, Alberto Sesana, Naoki Seto, Lijing Shao, Stuart Shapiro, Carlos Sopena, Nicholas C. Stone, Arthur Suvorov, Nicola Tamanini, Tomas Tamfal, Thomas Tauris, Karel Temmink, John Tomsick, Silvia Toonen, Alejandro Torres-Orjuela, Martina Toscani, Antonios

- Tsokaros, Caner Unal, Verónica Vázquez-Aceves, Rosa Valiante, Maurice van Putten, Jan van Roestel, Christian Vignali, Marta Volonteri, Kinwah Wu, Ziri Younsi, Shenghua Yu, Silvia Zane, Lorenz Zwick, Fabio Antonini, Vishal Baibhav, Enrico Barausse, Alexander Bonilla Rivera, Marica Branchesi, Graziella Branduardi-Raymont, Kevin Burdge, Srija Chakraborty, Jorge Cuadra, Kristen Dage, Benjamin Davis, Selma E. de Mink, Roberto Decarli, Daniela Doneva, Stephanie Escoffier, Poshak Gandhi, Francesco Haardt, Carlos O. Lousto, Samaya Nissanke, Jason Nordhaus, Richard O’Shaughnessy, Simon Portegies Zwart, Adam Pound, Fabian Schussler, Olga Sergijenko, Alessandro Spallicci, Daniele Vernieri, and Alejandro Vigna-Gómez. Astrophysics with the Laser Interferometer Space Antenna. *Living Reviews in Relativity*, 26(1):2, December 2023. doi: 10.1007/s41114-022-00041-y.
- Martin Asplund, Nicolas Grevesse, A. Jacques Sauval, and Pat Scott. The Chemical Composition of the Sun. *ARA&A*, 47(1):481–522, September 2009. doi: 10.1146/annurev.astro.46.060407.145222.
- John Baker, Jillian Bellovary, Peter L. Bender, Emanuele Berti, Robert Caldwell, Jordan Camp, John W. Conklin, Neil Cornish, Curt Cutler, Ryan DeRosa, Michael Eracleous, Elizabeth C. Ferrara, Samuel Francis, Martin Hewitson, Kelly Holley-Bockelmann, Ann Hornschemeier, Craig Hogan, Brittany Kamai, Bernard J. Kelly, Joey Shapiro Key, Shane L. Larson, Jeff Livas, Sridhar Manthripragada, Kirk McKenzie, Sean T. McWilliams, Guido Mueller, Priyamvada Natarajan, Kenji Numata, Norman Rioux, Shannon R. Sankar, Jeremy Schnittman, David Shoemaker, Deirdre Shoemaker, Jacob Slutsky, Robert Spero, Robin Stebbins, Ira Thorpe, Michele Valisneri, Brent Ware, Peter Wass, Anthony Yu, and John Ziemer. The laser interferometer space antenna: Unveiling the millihertz gravitational wave sky. 2019. URL <https://arxiv.org/abs/1907.06482>.
- Hale Bradt. *Astrophysics processes*. Cambridge University Press, 2006.
- Katelyn Breivik, Scott Coughlin, Michael Zevin, Carl L. Rodriguez, Kyle Kremer, Claire S. Ye, Jeff J. Andrews, Michael Kurkowski, Matthew C. Digman, Shane L. Larson, and Frederic A. Rasio. COSMIC Variance in Binary Population Synthesis. *ApJ*, 898(1):71, July 2020. doi: 10.3847/1538-4357/ab9d85.
- Alessandro Bressan, Paola Marigo, Léo. Girardi, Bernardo Salasnich, Claudia Dal Cero, Stefano Rubele, and Ambra Nanni. PARSEC: stellar tracks and isochrones with the PAdova and TRieste Stellar Evolution Code. *MNRAS*, 427(1):127–145, November 2012. doi: 10.1111/j.1365-2966.2012.21948.x.
- E. Margaret Burbidge, G. R. Burbidge, William A. Fowler, and F. Hoyle. Synthesis of the elements in stars. *Rev. Mod. Phys.*, 29:547–650, Oct 1957. doi: 10.1103/RevModPhys.29.547. URL <https://link.aps.org/doi/10.1103/RevModPhys.29.547>.
- S. Chandrasekhar. The Maximum Mass of Ideal White Dwarfs. *ApJ*, 74:81, July 1931. doi: 10.1086/143324.

- Katerina Chatziioannou, James Alexander Clark, Andreas Bauswein, Margaret Millhouse, Tyson B. Littenberg, and Neil Cornish. Inferring the post-merger gravitational wave emission from binary neutron star coalescences. *Phys. Rev. D*, 96(12):124035, December 2017. doi: 10.1103/PhysRevD.96.124035.
- Priyam Das, Ivo R. Seitenzahl, Ashley J. Ruiter, Friedrich K. Röpke, Rüdiger Pakmor, Frédéric P. A. Vogt, Christine E. Collins, Parviz Ghavamian, Stuart A. Sim, Brian J. Williams, Stefan Taubenberger, J. Martin Laming, Janette Suherli, Ralph Sutherland, and Nicolás Rodríguez-Segovia. Calcium in a supernova remnant as a fingerprint of a sub-chandrasekhar-mass explosion. *Nature Astronomy*, July 2025. ISSN 2397-3366. doi: 10.1038/s41550-025-02589-5. URL <https://doi.org/10.1038/s41550-025-02589-5>.
- G. De Vaucouleurs. Structure, dynamics and statistical properties of galaxies. *Symposium - International Astronomical Union*, 58:1–53, 1974. doi: 10.1017/S007418090002430X.
- Gerard de Vaucouleurs, Antoinette de Vaucouleurs, Herold G. Corwin, Jr., Ronald J. Buta, Georges Paturel, and Pascal Fouque. *Third Reference Catalogue of Bright Galaxies*. 1991.
- Albert Einstein. Über Gravitationswellen. *Sitzungsberichte der Königlich Preussischen Akademie der Wissenschaften*, pages 154–167, January 1918.
- S. M. Faber and J. S. Gallagher. Masses and mass-to-light ratios of galaxies. *ARA&A*, 17:135–187, January 1979. doi: 10.1146/annurev.aa.17.090179.001031.
- V. Ferrari, L. Gualtieri, and P. Pani. *General Relativity and its Applications: Black Holes, Compact Stars and Gravitational Waves*. CRC Press, 1st edition, 2020. doi: 10.1201/9780429491405. URL <https://doi.org/10.1201/9780429491405>.
- Fink, M., Röpke, F. K., Hillebrandt, W., Seitenzahl, I. R., Sim, S. A., and Kromer, M. Double-detonation sub-chandrasekhar supernovae: can minimum helium shell masses detonate the core? *A&A*, 514:A53, 2010. doi: 10.1051/0004-6361/200913892. URL <https://doi.org/10.1051/0004-6361/200913892>.
- Vayujeet Gokhale, Xiao Meng Peng, and Juhan Frank. Evolution of Close White Dwarf Binaries. *ApJ*, 655(2):1010–1024, February 2007. doi: 10.1086/510119.
- K. M. Gorski, E. Hivon, A. J. Banday, B. D. Wandelt, F. K. Hansen, M. Reinecke, and M. Bartelmann. Healpix: A framework for high-resolution discretization and fast analysis of data distributed on the sphere. *The Astrophysical Journal*, 622(2):759–771, April 2005. ISSN 1538-4357. doi: 10.1086/427976. URL <http://dx.doi.org/10.1086/427976>.
- D Hils, P L Bender, and R F Webbink. Gravitational radiation from the galaxy. *Astrophysical Journal; (USA)*, 360, 09 1990. ISSN ISSN 0004-637X. doi: 10.1086/169098. URL <https://www.osti.gov/biblio/6282329>.

- F. Hoyle. On Nuclear Reactions Occuring in Very Hot Stars. 1. The Synthesis of Elements from Carbon to Nickel. *Astrophys. J. Suppl.*, 1:121–146, 1954. doi: 10.1086/190005.
- E. P. Hubble. Extragalactic nebulae. *ApJ*, 64:321–369, December 1926. doi: 10.1086/143018.
- Jarrold R. Hurley, Onno R. Pols, and Christopher A. Tout. Comprehensive analytic formulae for stellar evolution as a function of mass and metallicity. *MNRAS*, 315(3): 543–569, July 2000. doi: 10.1046/j.1365-8711.2000.03426.x.
- Jarrold R. Hurley, Christopher A. Tout, and Onno R. Pols. Evolution of binary stars and the effect of tides on binary populations. *MNRAS*, 329(4):897–928, February 2002. doi: 10.1046/j.1365-8711.2002.05038.x.
- Icko Iben, Jr. Stellar Evolution.VI. Evolution from the Main Sequence to the Red-Giant Branch for Stars of Mass $1 M_{\odot}$, $1.25 M_{\odot}$, and $1.5 M_{\odot}$. *ApJ*, 147:624, February 1967. doi: 10.1086/149040.
- Renée C. Kraan-Korteweg and Ofer Lahav. The Universe behind the Milky Way. *A&A Rev.*, 10(3):211–261, January 2000. doi: 10.1007/s001590000011.
- T Kupfer, V Korol, S Shah, G Nelemans, T R Marsh, G Ramsay, P J Groot, D T H Steeghs, and E M Rossi. Lisa verification binaries with updated distances from gaia data release 2. *Monthly Notices of the Royal Astronomical Society*, 480(1):302–309, 06 2018. ISSN 0035-8711. doi: 10.1093/mnras/sty1545. URL <https://doi.org/10.1093/mnras/sty1545>.
- Thomas Kupfer, Valeriya Korol, Tyson B. Littenberg, Sweta Shah, Etienne Savalle, Paul J. Groot, Thomas R. Marsh, Maude Le Jeune, Gijs Nelemans, Anna F. Pala, Antoine Petiteau, Gavin Ramsay, Danny Steeghs, and Stanislav Babak. Lisa galactic binaries with astrometry from gaia dr3. *The Astrophysical Journal*, 963(2):100, mar 2024. doi: 10.3847/1538-4357/ad2068. URL <https://dx.doi.org/10.3847/1538-4357/ad2068>.
- Jun Luo, Li-Sheng Chen, Hui-Zong Duan, Yun-Gui Gong, Shoucun Hu, Jianghui Ji, Qi Liu, Jianwei Mei, Vadim Milyukov, Mikhail Sazhin, Cheng-Gang Shao, Viktor T Toth, Hai-Bo Tu, Yamin Wang, Yan Wang, Hsien-Chi Yeh, Ming-Sheng Zhan, Yonghe Zhang, Vladimir Zharov, and Ze-Bing Zhou. Tianqin: a space-borne gravitational wave detector. *Classical and Quantum Gravity*, 33(3):035010, January 2016. ISSN 1361-6382. doi: 10.1088/0264-9381/33/3/035010. URL <http://dx.doi.org/10.1088/0264-9381/33/3/035010>.
- Michele Maggiore. *Gravitational Waves: Volume 1: Theory and Experiments*. Oxford University Press, 10 2007. ISBN 9780198570745. doi: 10.1093/acprof:oso/9780198570745.001.0001. URL <https://doi.org/10.1093/acprof:oso/9780198570745.001.0001>.
- D. V. Martynov, E. D. Hall, B. P. Abbott, R. Abbott, T. D. Abbott, C. Adams, R. X. Adhikari, R. A. Anderson, S. B. Anderson, K. Arai, M. A. Arain, S. M. Aston,

L. Austin, S. W. Ballmer, M. Barbet, D. Barker, B. Barr, L. Barsotti, J. Bartlett, M. A. Barton, I. Bartos, J. C. Batch, A. S. Bell, I. Belopolski, J. Bergman, J. Betzwieser, G. Billingsley, J. Birch, S. Biscans, C. Biwer, E. Black, C. D. Blair, C. Bogan, C. Bond, R. Bork, D. O. Bridges, A. F. Brooks, D. D. Brown, L. Carbone, C. Celerier, G. Ciani, F. Clara, D. Cook, S. T. Countryman, M. J. Cowart, D. C. Coyne, A. Cumming, L. Cunningham, M. Damjanic, R. Dannenberg, K. Danzmann, C. F. Da Silva Costa, E. J. Daw, D. DeBra, R. T. DeRosa, R. DeSalvo, K. L. Doolley, S. Doravari, J. C. Driggers, S. E. Dwyer, A. Effler, T. Etzel, M. Evans, T. M. Evans, M. Factourovich, H. Fair, D. Feldbaum, R. P. Fisher, S. Foley, M. Frede, A. Freise, P. Fritschel, V. V. Frolov, P. Fulda, M. Fyffe, V. Galdi, J. A. Giaime, K. D. Giardina, J. R. Gleason, R. Goetz, S. Gras, C. Gray, R. J. S. Greenhalgh, H. Grote, C. J. Guido, K. E. Gushwa, E. K. Gustafson, R. Gustafson, G. Hammond, J. Hanks, J. Hanson, T. Hardwick, G. M. Harry, K. Haughian, J. Heefner, M. C. Heintze, A. W. Heptonstall, D. Hoak, J. Hough, A. Ivanov, K. Izumi, M. Jacobson, E. James, R. Jones, S. Kandhasamy, S. Karki, M. Kasprzack, S. Kaufer, K. Kawabe, W. Kells, N. Kijbunchoo, E. J. King, P. J. King, D. L. Kinzel, J. S. Kissel, K. Kokeyama, W. Z. Korth, G. Kuehn, P. Kwee, M. Landry, B. Lantz, A. Le Roux, B. M. Levine, J. B. Lewis, V. Lhuillier, N. A. Lockerbie, M. Lormand, M. J. Lubinski, A. P. Lundgren, T. MacDonald, M. MacInnis, D. M. Macleod, M. Mageswaran, K. Maitland, S. Márka, Z. Márka, A. S. Markosyan, E. Maros, I. W. Martin, R. M. Martin, J. N. Marx, K. Mason, T. J. Massinger, F. Matichard, N. Mavalvala, R. McCarthy, D. E. McClelland, S. McCormick, G. McIntyre, J. McIver, E. L. Merilh, M. S. Meyer, P. M. Meyers, J. Miller, R. Mittleman, G. Moreno, C. L. Mueller, G. Mueller, A. Mullavey, J. Munch, P. G. Murray, L. K. Nuttall, J. Oberling, J. O'Dell, P. Oppermann, Richard J. Oram, B. O'Reilly, C. Osthelder, D. J. Ottaway, H. Overmier, J. R. Palamos, H. R. Paris, W. Parker, Z. Patrick, A. Pele, S. Penn, M. Phelps, M. Pickenpack, V. Pierro, I. Pinto, J. Poeld, M. Principe, L. Prokhorov, O. Puncken, V. Quetschke, E. A. Quintero, F. J. Raab, H. Radkins, P. Raffai, C. R. Ramet, C. M. Reed, S. Reid, D. H. Reitze, N. A. Robertson, J. G. Rollins, V. J. Roma, J. H. Romie, S. Rowan, K. Ryan, T. Sadecki, E. J. Sanchez, V. Sandberg, V. Sannibale, R. L. Savage, R. M. S. Schofield, B. Schultz, P. Schwinberg, D. Sellers, A. Seigny, D. A. Shaddock, Z. Shao, B. Shapiro, P. Shawhan, D. H. Shoemaker, D. Sigg, B. J. J. Slagmolen, J. R. Smith, M. R. Smith, N. D. Smith-Lefebvre, B. Sorazu, A. Staley, A. J. Stein, A. Stochino, K. A. Strain, R. Taylor, M. Thomas, P. Thomas, K. A. Thorne, E. Thrane, K. V. Tokmakov, C. I. Torrie, G. Traylor, G. Vajente, G. Valdes, A. A. van Veggel, M. Vargas, A. Vecchio, P. J. Veitch, K. Venkateswara, T. Vo, C. Vorvick, S. J. Waldman, M. Walker, R. L. Ward, J. Warner, B. Weaver, R. Weiss, T. Welborn, P. Weßels, C. Wilkinson, P. A. Willems, L. Williams, B. Willke, I. Wilmot, L. Winkelmann, C. C. Wipf, J. Worden, G. Wu, H. Yamamoto, C. C. Yancey, H. Yu, L. Zhang, M. E. Zucker, and J. Zweizig. Sensitivity of the advanced ligo detectors at the beginning of gravitational wave astronomy. *Physical Review D*, 93(11), June 2016. ISSN 2470-0029. doi: 10.1103/physrevd.93.112004. URL <http://dx.doi.org/10.1103/PhysRevD.93.112004>.

G. Nelemans, L. R. Yungelson, and S. F. Portegies Zwart. The gravitational wave signal from the galactic disk population of binaries containing two compact objects. *Astronomy & Astrophysics*, 375(3):890–898, September 2001. ISSN 1432-0746. doi:

- 10.1051/0004-6361:20010683. URL <http://dx.doi.org/10.1051/0004-6361:20010683>.
- Travis Robson, Neil J Cornish, and Chang Liu. The construction and use of lisa sensitivity curves. *Classical and Quantum Gravity*, 36(10):105011, April 2019. ISSN 1361-6382. doi: 10.1088/1361-6382/ab1101. URL <http://dx.doi.org/10.1088/1361-6382/ab1101>.
- M. Schwarzschild and R. Härm. Red Giants of Population II. II. *ApJ*, 136:158, July 1962. doi: 10.1086/147361.
- Ken J. Shen and Lars Bildsten. Unstable helium shell burning on accreting white dwarfs. *The Astrophysical Journal*, 699(2):1365–1373, June 2009. ISSN 1538-4357. doi: 10.1088/0004-637x/699/2/1365. URL <http://dx.doi.org/10.1088/0004-637x/699/2/1365>.
- Ken J. Shen, Douglas Boubert, Boris T. Gänsicke, Saurabh W. Jha, Jennifer E. Andrews, Laura Chomiuk, Ryan J. Foley, Morgan Fraser, Mariusz Gromadzki, James Guillochon, Marissa M. Kotze, Kate Maguire, Matthew R. Siebert, Nathan Smith, Jay Strader, Carles Badenes, Wolfgang E. Kerzendorf, Detlev Koester, Markus Kromer, Broxton Miles, Rüdiger Pakmor, Josiah Schwab, Odette Toloza, Silvia Toonen, Dean M. Townsley, and Brian J. Williams. Three hypervelocity white dwarfs in gaia dr2: Evidence for dynamically driven double-degenerate double-detonation type ia supernovae. *The Astrophysical Journal*, 865(1):15, sep 2018. doi: 10.3847/1538-4357/aad55b. URL <https://dx.doi.org/10.3847/1538-4357/aad55b>.
- Mario Spera, Michela Mapelli, and Alessandro Bressan. The mass spectrum of compact remnants from the PARSEC stellar evolution tracks. *MNRAS*, 451(4):4086–4103, August 2015. doi: 10.1093/mnras/stv1161.
- Christy A. Tremonti, Timothy M. Heckman, Guinevere Kauffmann, Jarle Brinchmann, Stéphane Charlot, Simon D. M. White, Mark Seibert, Eric W. Peng, David J. Schlegel, Alan Uomoto, Masataka Fukugita, and Jon Brinkmann. The Origin of the Mass-Metallicity Relation: Insights from 53,000 Star-forming Galaxies in the Sloan Digital Sky Survey. *ApJ*, 613(2):898–913, October 2004. doi: 10.1086/423264.
- Darren J White, E J Daw, and V S Dhillon. A list of galaxies for gravitational wave searches. *Classical and Quantum Gravity*, 28(8):085016, March 2011. ISSN 1361-6382. doi: 10.1088/0264-9381/28/8/085016. URL <http://dx.doi.org/10.1088/0264-9381/28/8/085016>.