

PrEP

PREGNANT & BREASTFEEDING WOMEN

know the facts

PrEP is a safe HIV prevention method that HIV-negative people can use to prevent HIV:

- ♥ The pills need to be taken daily to help prevent HIV.
- ♥ Oral PrEP has been shown to reduce the chances of HIV infection by more than 90%.
- ♥ You have to take the pills every day, for as long as you need it.
- ♥ PrEP is only for people who are HIV-negative.

PrEP during pregnancy:

- ♥ If you are pregnant and have sex without a condom, your chances of getting HIV is much higher.
- ♥ If you test positive for HIV, you will receive ARV treatment - this prevents your baby from getting HIV.
- ♥ Using PrEP before, during and after pregnancy can prevent HIV.

What is the difference between PrEP, PEP, and ART?

All three use antiretrovirals in different combinations for different purposes:

♥ **PrEP** is when ARVs are taken **before** exposure to HIV, to **prevent** getting HIV.

♥ **PEP** is when ARVs are taken **after** exposure to HIV, to **prevent** HIV (within 72 hours and taken for 28 days only).

♥ **ART** is when ARVs are used to treat a person living with HIV, and is taken lifelong.

PrEP ♥ IS CHOICE
#ICHOOSEME



Version: Factsheet PrEP_PBFP_Oct 2025



www.myprep.co.za



@MyPrEPSouthAfrica



health

Department:
Health
REPUBLIC OF SOUTH AFRICA

PrEP

Pre
(before)

Exposure
(coming into
contact with HIV)

Prophylaxis
(a medicine to
prevent infection)

Where can I get PrEP...

PrEP is now available in all public primary health care clinics. Visit your nearest clinic if you are interested in using PrEP. For more information about PrEP, please visit myprep.co.za.

**Find your
nearest
clinic with
this code:**



Starting PrEP...

First visit:

- Health check, including screening for HIV and STIs, supported by counselling.
- Get your oral PrEP supply for a month.
- As oral PrEP builds up in your body, use a condom or abstain from sex for the first 7 days of taking oral PrEP.
- After 7 days, you need to continue taking oral PrEP daily for as long as you need it.

Month 1 visit:

- Health check, including screening for HIV and STIs, supported by counselling.
- Get your 3-month supply of oral PrEP pills.

Every 3 months:

- Health check, including screening for HIV and STIs, supported by counselling.
- Every 3 months, you return for an HIV test and a 3-month supply of oral PrEP.

Oral PrEP works best when taken daily and used with a condom.

Decide if PrEP is for you:

- ♥ PrEP is safe for you and your unborn baby and child, while you are pregnant and/or breastfeeding.
- ♥ PrEP can protect you from HIV.
- ♥ PrEP is easy to take, just one pill a day.
- ♥ You can take PrEP without anybody else knowing.
- ♥ You can take PrEP if you and your partner who is living with HIV want to have a baby.
- ♥ You can continue taking PrEP even when you are breastfeeding.

**Don't forget
your ante-
natal care
visits**

**PrEP is one of many
options for HIV
prevention. You can
also try:**

- ♥ Condoms
- ♥ Other PrEP options which may be available at your clinic, ask your provider
- ♥ Counselling
- ♥ PEP
- ♥ Treatment for STIs
- ♥ Male medical circumcision
- ♥ ART for partners living with HIV
- ♥ Regular HIV testing for you and your partner

You can take care of yourself and your baby, choose PrEP!

Use a
condom

If you test
negative for HIV,
you can
use PrEP!

Ask your
partner to
test for
HIV

Start and
continue with
ARVs if you
test positive
for HIV

Encourage your
partner living
with HIV to take
ARVs daily

