

### 1 START with an HIV test: follow national HTS guidelines

**Negative**

No signs of acute HIV infection or exposure in the past 72 hours

**Positive**

Post-test counselling + **initiate** immediate ART as per HIV treatment guideline

Assess for PEP and refer for further management

### 2 Assess the need for HIV prevention

**Counsel for HIV prevention**

**CLIENT REQUESTS PrEP**

**OR**

**CLIENT MAY BENEFIT FROM PrEP**



### 3 PrEP choice counselling

Consider client needs and preferences

Offer different PrEP options available

**If client chooses lenacapavir**

If client chooses other PrEP method, assess as per guidelines (oral PrEP)

### 4 Assess for eligibility AND screen

**Individuals eligible for lenacapavir:**

HIV-negative test

Weight ≥ 35 kg

No signs of acute HIV infection

No allergies/hypersensitivity/contraindications to lenacapavir

**Also check (at initiation and every follow-up visit):**

Pregnancy assessment (for risk-benefit discussion)

TB and TB treatment

Other medications for possible drug-drug interactions

Confirm that client wants to proceed and can commit to 6 monthly schedule.

### 5 Lenacapavir initiation

**If client is eligible**

Counsel about lenacapavir use

Administer two injections

Observe client taking two oral tablets

Provide two tablets for next day

Provide dates for 4 week and 26 week follow-up appointments

Assist the client to enter the appointment date on their mobile phone calendar

### 6 Follow-up

**4 Weeks (1 month) after initiation**

Assess for side effects and perform HIV test

**26 weeks (6 months) after initiation**

Perform HIV test and follow-up injections

**then Every 26 weeks (6 months)**

Perform HIV test and follow-up injections

Confirm contact details for client  
Explain how to reschedule an appointment

**At every visit:**

- HIV test
- Promote condom use
- Assess for side effects
- STI and contraceptive services
- Pregnancy screening
- Behavioural counselling
- Appointment reminder

**Check time since last injection**

**24-28 weeks:** Continue with next injections.

**More than 28 weeks:** Restart initiation process.